

ASTHMA FACTS

Asthma continues to be a serious public health problem. According to the Centers for Disease Control and Prevention:

- Almost 25 million people, including almost 7 million children, have asthma.^{1,2}
 - * Asthma prevalence is higher among persons with family income below the poverty level.
- Almost 13 million people report having an asthma attack in the past year.³
- Asthma accounts for over 15 million physician office and hospital outpatient department visits,⁴ and nearly 2 million emergency department visits each year.³
- African Americans continue to have higher rates of asthma emergency department visits, hospitalizations, and deaths than do Caucasians:
 - * The rate of emergency department visits is 330% higher.³
 - * The hospitalization rate is 220% higher.³
 - * The asthma death rate is 190% higher.³
- Approximately 3 million Hispanics in the U.S. have asthma and Puerto Ricans are disproportionately impacted:
 - * The rate of asthma among Puerto Ricans is 113% higher than non-Hispanic white people and 50% higher than non-Hispanic black people.³
 - * The prevalence of asthma attacks is highest among Puerto Ricans.³

Asthma in Children

- * Asthma is one of the most common serious chronic diseases of childhood.
- * Asthma is the third-ranking cause of hospitalization among children under 15.⁵
- * An average of one out of every 10 school-aged children has asthma.⁶
- * 10.5 million school days are missed each year due to asthma.⁷

The Cost of Asthma

- * Annual expenditures for health and lost productivity due to asthma are estimated at over \$20 billion, according to the National Heart, Lung, and Blood Institute.⁸

Environmental Factors

- Indoor and outdoor environmental factors can trigger asthma attacks: dust mites, molds, cockroaches, pet dander, and secondhand smoke.

ASTHMA FACTS

Asthma Can be Controlled

- With a plan that includes medical treatment and control of environmental triggers, people with asthma can lead healthy, active lives.

Asthma and the Environment

Research by EPA and others has shown that:

- Dust mites, molds, cockroaches, pet dander, and secondhand smoke trigger asthma attacks.
- Exposure to secondhand smoke can cause asthma in pre-school aged children.
- Exposure to dust mites can cause asthma.
- Ozone and particle pollution can cause asthma attacks.
 - * When ozone levels are high, more people with asthma have attacks that require a doctor's attention.
 - * Ozone makes people more sensitive to asthma triggers such as pet dander, pollen, dust mites, and mold.

Learn more at www.epa.gov/asthma

References

1. Summary Health Statistics for U.S. Adults: National Health Interview Survey, 2006, Tables 3 and 4
2. Summary Health Statistics for U.S. Children: National Health Interview Survey, 2008, Table 1. http://www.cdc.gov/nchs/data/series/sr_10/sr10_244.pdf
3. <http://www.cdc.gov/nchs/data/nhsr/nhsr032.pdf>
4. National Ambulatory Medical Care Survey: 2006 Summary, Table 12
5. DeFrances CJ, Cullen KA, Kozak LJ. National Hospital Discharge Society: 2005 Annual Summary with detailed Diagnosis and Procedure Data. National Center for Health Statistics. Vital Health Statistics 12 (165); 2007
6. American Lung Association, Epidemiology and Statistics Unit, Research Program Services. Trends in Asthma Morbidity and Mortality. February 2010. <http://www.lungusa.org/finding-cures/our-research/trend-reports/asthma-trend-report.pdf>
7. Akinbami LJ. The State of Childhood Asthma, United States, 1980-2005, Advance Data from Vital and Health Statistics: no 381, Revised December 29, 2006, Hyattsville, MD: National Center for Health Statistics, 2006 (NHIS 2003 absenteeism)
8. National Heart, Lung, and Blood Institute Chartbook on Cardiovascular, Lung and Blood Diseases, U.S. Department of Health and Human Services, National Institute of Health, 2009. (http://www.nhlbi.nih.gov/resources/docs/2009_ChartBook.pdf)