



U.S. Environmental Protection Agency



U.S. Maritime Administration

This document is one section from the “National Guidance: Best Management Practices for Preparing Vessels Intended to Create Artificial Reefs,” published in May 2006. The reference number is EPA 842-B-06-002. You can find the entire document at <http://www.epa.gov/owow/oceans/habitat/artificialreefs/index.html>.

National Guidance: Best Management Practices for Preparing Vessels Intended to Create Artificial Reefs

Appendix C

May 2006

Appendix C

Information related to materials found on scuttled vessels that may have potentially hazardous effects on the marine environment*

*The text provided in this appendix is an excerpt from the 2005 “Policy Statement of the National Marine Sanctuary Program: Artificial Reef Permitting Guidelines.”

Scuttled Vessels

The scuttling of vessels requires particular attention in this policy because of their size and potential toxicological effects on the environment. As discussed above, sunken ships potentially attract divers away from natural reefs and thus may be beneficial to natural reefs in National Marine Sanctuaries. However, there is a wide array of concerns that must be addressed before intentionally sinking a ship.

The removal of petroleum products, hazardous materials, paint cans, batteries, plastics, oil, and fuel is specified on the U.S. Coast Guard’s Ocean Disposal/Artificial Reef Inspection form. Additionally, under the Toxic Substances Control Act (TSCA), the EPA has the authority to gather information on and regulate chemical substances and mixtures imminently hazardous or presenting unreasonable risk of injury to public health or the environment. Despite these controls, some materials of concern may still remain on items used as artificial reef material. Such materials include: asbestos, polychlorinated biphenols (PCBs), iron, lead paint, and antifouling paint. The National Marine Sanctuaries Program (NMSP) should consider the risks associated with materials remaining on vessels to be used as artificial reefs. The NMSP will consult with appropriate agencies (i.e., U.S. EPA, MARAD) to determine the best management practices to use in evaluating materials for pollution potential).

Asbestos is the name given to six naturally occurring minerals that are used as insulators and fire retardants. Several studies have investigated the effects of asbestos on fish (Batterman and Cook 1981, Belanger *et al.* 1990, Belanger *et al.* 1986, Woodhead *et al.* 1983). The findings indicate that asbestos concentrations on the order of 10^6 to 10^8 fibers/L may cause epidermal lesions, epithelial hypertrophy, kidney damage, decreased orientation and swimming ability, degradation of the lateral line, reduced growth, and increased mortality in fish. Undisturbed, non-friable (not easily crumbled) asbestos has been found to be relatively harmless (Garcia and Salzwedel 1995, Montoya *et al.* 1985).

PCBs may still exist in water-tight gaskets, cable insulation, paint, transformers, capacitors, and other components of decommissioned Navy vessels (Martore *et al.* 1996, Eisler and Belisle 1996). These chemicals have been implicated in: reduced primary productivity in phytoplankton; reduced hatchability of contaminated fish and bird eggs; reproductive failure in seals; altered steroid levels and subsequent reproductive impairment in fish and sea stars; reduced fertilization efficiency in sea urchins; and reduced plasma retinal and thyroid hormone levels potentially leading to increased susceptibility to microbial infections, reproductive disorders and other pathological alternation in seals and other marine mammals (Adams and Slaughter-Williams 1988, Brouwer *et al.* 1989, Clark 1992, den Besten *et al.* 1991).

Antifouling paints typically containing tributyltin (TBT) and copper (Cu) are often used to paint vessel hulls to inhibit the growth of organisms below the water line. An IMO convention to

control the use of harmful anti-fouling systems on ships was adopted on October 5, 2001. The convention will prohibit the use of harmful organotins, including TBT, in anti-fouling paints used on ships and establish a mechanism to prevent the potential future use of other harmful substances in anti-fouling systems. TBT has been found to be toxic to non-target, non-fouling organisms at low levels (approximately 7.5-10.5 ng TBT/L). One of its most marked effects has been the induction of shell thickening and growth anomalies in oysters and imposex in the dogwhelk *Nucella lapillus* potentially leading to sterility (Gibbs *et al.* 1998).⁸ The discovery of the highly toxic nature of TBT-based paints has led many countries to ban the use of these paints for non-aluminum hulled vessels less than 25 meters in length. Copper, though an effective antifoulant, has not been shown to cause extensive effects on non-target organisms at relatively low levels. When present in high concentrations, however, copper can be toxic to aquatic life (Sorrenson 1991). In a study conducted when a cargo ship collided with part of the Great Barrier Reef and remained grounded for 12 days, sediment containing 8.0 mg kg super(-1) TBT, 72 mg kg super(-1) Cu and 92 mg kg super(-1) Zn was found to significantly inhibit larval settlement and metamorphosis (Negri *et al.* 2002). At this level of contamination, larvae survived but contracted to a spherical shape and swimming and searching behavior ceased. At higher contamination levels, 100% mortality was recorded. These results indicate that the contamination of sediment by anti-fouling paint has the potential to significantly reduce coral recruitment in the immediate vicinity of the site and that this contamination may threaten the recovery of the resident coral community unless the paint is removed.

Iron, an essential element like copper, can be contributed to the environment from steel hulls of sunken vessels. As an essential element, iron levels will tend to be closely regulated by organisms, and thus, it is unlikely that any pollution-derived effects will be observed except in severe and localized cases (Thompson 1990). Corals living in seawater with high iron concentrations have been shown to incorporate the iron into their skeletons (Brown *et al.* 1991). Studies on phytoplankton and macroalgae indicate that in areas where plant nutrients such as nitrate and phosphate are abundant the availability of iron is actually a limiting factor in growth and biomass (Coale *et al.* 1996, Frost 1996, Matsunaga *et al.* 1994, Takeda 1998, Wells *et al.* 1995). Hence the concern of unnatural iron inputs from artificial reefs seems to center not on the occurrence of adverse toxicological effects in marine organisms, but rather on the alteration of the composition of natural assemblages of algae and species which compete with algae.

Lead paint has been used on the interiors of some vessels. Lead has no biological function and, therefore, exhibits accumulation trends in organisms (Thompson 1990). Corals have been shown to incorporate lead into their skeletons (Dodge and Gilbert 1984). Unicellular algae and sea urchins appear to be the most sensitive marine organisms (Berhard 1980). Growth inhibition has been observed in the algae species *Thalassiosira pseudonana* and *Porphyridium marinum* exposed to lead as well as in sea urchins.

References

- Adams, J.A. and S. Slaughter-Williams. 1988. The effects of PCBs on fertilization and morphology in *Arbacia punctulata*. *Water Air Soil Pollut.* 38: 299-310.
- Batterman, A.L. and P.M. Cook. 1981. Determination of mineral fiber concentrations in fish tissues. *Can. J. Fish. Aquat. Sci.* 38: 952-959.

⁸ Imposex is defined as the development of specific male sexual organs in females.

- Belanger, S.E., K. Schurr, D.A. Allen, and A.F. Gohara. 1986. Effects of chrysotile asbestos on coho salmon and green sunfish: evidence of pathological stress. *Environ. Res.* 39: 74-85.
- Belanger, S.E., D.S. Cherry, and J. Cairns, Jr. 1990. Functional and pathological impairment of Japanese Medaka (*Oryzias latipes*) by long-term asbestos exposure. *Aquat. Toxicol.* 17: 133-154.
- Brouwer, A., P.J.H. Reijnders, and J.H. Koeman. 1989. Polychlorinated biphenol (PCB)-contaminated fish induces vitamin A and thyroid hormone deficiency in the common seal. *Aquatic Toxicology.* 15: 99-106.
- Brown, B.E., A.W. Tudhope, M.D.A. Le Tissier, and T.P. Scoffin. 1991. A novel mechanism for iron incorporation into coral skeletons. *Coral Reefs* 10: 211-215.
- Clark, R.B. 1992. *Marine Pollution*. Clarendon Press, Oxford, 172.
- Coale, K.H., S.E. Fitzwater, R.M. Gordon, K.S. Johnson, and R.T. Barber. 1996. Control of community growth and export production by upwelled iron in the equatorial Pacific Ocean. *Lett. Nature* 379: 621-624.
- den Beston, P.J., J.M.L. Elenbaas, J.R. Maas, S.J. Dieleman, H.J. Herwig, and P.A. Voogt. 1991. Effects of cadmium and polychlorinated biphenols on steroid metabolism and cytochrome P-450 monooxygenase system in the sea star *Asterias rubens* L. *Aquatic Toxicology.* 20: 95-100.
- Frost, B.W. 1996. Phytoplankton bloom on iron rations. *Nature* 383: 475-476.
- Garcia, C.B. and H. Salzwedel. 1995. Successional patterns on fouling plates in the Bay of Santa Marta, Colombian Caribbean. *An. Inst. Invest. Mar. Punta de Betin.* 24: 95-121.
- Gibbs, P.E., P.L. Pascoe, and G.R. Burt. 1988. Sex change in the female dog-whelck, *Nucella lapillus*, induced by tributyltin from antifouling paints. *J. Mar. Biol. Ass. U.K.* 68:715-731.
- Martore, R.M., T.D. Mathews, and M. Bell. 1998. Levels of PCBs and heavy metals in biota found on ex-military ships used as artificial reefs. South Carolina Department of Natural Resources, Charleston, South Carolina.
- Matsunaga, K., Y. Suzuki, K. Kuma, and I. Kudo. 1994. Diffusion of Fe(II) from an iron propagation cage and its effect on tissue iron and pigments of macroalgae on the cage. *J. Appl. Phycol.* 6: 397-403.
- Montoya, A.J., Q.R. Quesada, Z.E. Madriz, M.E. Castro, and P.O. Urpi. 1985. Comparative analysis of substrates for collection of mangrove oyster spat in Viscaya estuary, Limon, Costa Rica. *Rev. Biol. Trop.* 33: 1-6.
- Negri, A.P., L.D. Smith, N.S. Webster, A.J. Heyward. 2002. Understanding ship-grounding impacts on a coral reef: potential effects of anti-foulant paint contamination on coral recruitment. *Mar. Pollut. Bull.* Vol. 44: 111-117.
- Sorrenson, E.M. 1991. *Metal poisoning in fish*. CRC Press, Inc., Boca Raton, 374 pp.
- Takeda, S. 1998. Influence of iron availability on nutrient consumption ratio of diatoms in oceanic waters. *Nature* 393: 774-777.
- Thompson, D.R. 1990. Metal levels in marine vertebrates. In R.W. Furness and P.S. Rainbow (eds.), *Heavy Metals in the Marine Environment*. CRC Press, Inc., Boca Raton, pp 143-183.
- Wells, M.L. N.M. Price, and K.W. Bruland. 1995. Iron chemistry in seawater and its relationship to phytoplankton: a workshop report. *Mar. Chem.* 48: 157-182.
- Woodhead, A.D., R.B. Setlow, and V. Pond. 1983. The effects of chronic exposure to asbestos fibers in the Amazon molly, *Poecilia Formosa*. *Environ. International.* 9:173-176.