

Fact Sheet Date: March 12, 1998

**NEW YORK STATE  
- HUMAN HEALTH FACT SHEET -**

**Ambient Water Quality Value  
Based on Human Consumption of Fish**

**SUBSTANCE:** Hexachlorobutadiene

**CAS REGISTRY NUMBER:** 87-68-3

**AMBIENT WATER QUALITY VALUE:** 0.01 ug/L

**BASIS:** Bioaccumulation

**INTRODUCTION**

This value applies to the water column and is designed to protect humans from the effects of waterborne contaminants that may bioaccumulate in fish; it is referred to as a Health (Fish Consumption) or H(FC) value. The H(FC) value is based on three components, the toxicity of the substance to humans, the extent to which it bioaccumulates in fish, and the rate of fish consumption.

**SUMMARY OF INFORMATION**

**A. Toxicity**

The toxicity of hexachlorobutadiene (HCBd) relevant to human health is described in a separate fact sheet (NYS, 1997). That fact sheet, which supports an ambient water quality value for protection of sources of potable water, derives an acceptable daily intake, or ADI, of 0.067 ug HCBd/kg/day, for nononcogenic effects.

**B. Bioaccumulation**

A measurement of bioaccumulation is necessary to derive a value to protect human consumers of fish. Bioaccumulation is the process by which a substance becomes concentrated in an organism through the organism's exposure to the contaminant in food and water. Bioaccumulation is represented numerically by a

bioaccumulation factor, or BAF, which is the ratio of the concentration of a substance in the organism to that in the water column.

The term bioconcentration also describes the concentration of a substance in an organism relative to the concentration in the water column. A bioconcentration factor (BCF), however, is measured with exposure to the contaminant by water only. A BCF may be equal to the BAF for many substances, but can substantially underestimate it for others.

U.S. EPA (1995a) has promulgated, as final Federal regulations, procedures for deriving bioaccumulation factors. The procedures are believed appropriate for deriving statewide values and are being used in this fact sheet.

A key aspect of this procedure is that bioaccumulation is believed to be related to the concentration of freely dissolved substance. Hydrophobic organic substances are considered to exist in water in three phases: freely dissolved, sorbed to dissolved organic matter and sorbed to suspended solids (U.S. EPA, 1995b). Because BAF determinations are often based on measurements of total or dissolved substance, a measured BAF must be adjusted based on the estimated fraction of freely dissolved material. In addition, because measured BAFs are determined based on the percent lipid in the species studied, they are adjusted, or normalized, to 100% lipid to allow comparison of BAFs derived from species with different tissue lipid fractions. A BAF adjusted for both fraction freely dissolved and normalized to 100% lipid is referred to as a "baseline BAF."

Although bioaccumulation is related to the freely dissolved substance, water quality criteria are based on total substance. A baseline BAF, therefore, is readjusted to a final BAF by the expected fraction freely dissolved and fish lipid content for the waters for which criteria are established. The relationship of field-measured or final BAF to the baseline BAF is shown in equation 1:

$$(Eq. 1) \quad \text{Baseline BAF} = \left[ \frac{\text{Field or Final BAF}}{f_{fd}} \right] \left[ \frac{1}{f_l} \right]$$

where  $f_l$  = fraction of tissue that is lipid and  $f_{fd}$  = fraction of substance that is freely dissolved.

U.S. EPA (1995b) presented the following equation for estimating  $f_{fd}$ :

$$(Eq. 2) \quad f_{fd} = \frac{1}{1 + \frac{(DOC)(K_{ow})}{10} + (POC)(K_{ow})}$$

where  $K_{ow}$  is the n-octanol-water partition coefficient of the substance in question, and DOC and POC are concentrations of dissolved and particulate organic carbon, respectively, in kg/L. The basis for this equation is described by U.S. EPA (1995b).

When deriving a baseline BAF from a field-measured BAF, DOC and POC levels under which the field BAF was determined are used to calculate a  $f_{fd}$ . When the baseline BAF is readjusted to yield a final BAF, the DOC and POC levels appropriate for the applicability of the criterion are used.

#### Derivation of Baseline BAF for HCBd

U.S. EPA (1995b), as part of the documentation for the Great Lakes Water Quality Initiative, presents baseline BAFs for a number of substances. For HCBd, these values are presented in Table 1.

Table 1	
Baseline BAFs for HCBd (U.S. EPA, 1995b)	
Trophic Level	Baseline BAF (L/kg)
3	354,800
4	43,940

These values have been reviewed and are believed appropriate for both the Great Lakes and the rest of the State. The data U.S. EPA used and calculations needed to derive these values are shown below. (Note: The readjustment of these baseline BAFs to final BAFs is described under Derivation of Water Quality Values).

Table 2						
Data Used to Calculate Baseline BAFs for HCBd						
Fish (Trophic Level)	Fish Tissue Conc. <sup>(1)</sup> (ng/g)	Water Conc. (pg/L)	$f_l$	DOC <sup>(2)</sup> (kg/L)	POC <sup>(2)</sup> (kg/L)	Log $K_{ow}$ <sup>(2)</sup>
Sculpin (3)	0.5 <sup>(1)</sup>	18 <sup>(1)</sup>	0.08 <sup>(1)</sup>	$2 \times 10^{-6}$	0	4.842
R. trout (4)	0.1637 <sup>(4)</sup>	50 <sup>(3)</sup>	0.07592 <sup>(4)</sup>	$2 \times 10^{-6}$	$7.5 \times 10^{-8}$	4.842
<sup>1</sup> Oliver and Niimi (1988) <sup>2</sup> U.S. EPA (1995b) <sup>3</sup> Oliver and Niimi (1983) <sup>4</sup> Geometric mean of 10 values from Oliver and Niimi (1983) (Stephan, 1995)						

Field BAFs are calculated as the ratio of fish tissue to water concentrations of the substance. Field BAFs based on data in Table 2 are presented in Table 3.

Table 3	
Field BAFs for HCBd	
Fish (Trophic Level)	Field BAF (L/kg)
Sculpin (3)	27,780
Rainbow trout (4)	3,274

The fraction freely dissolved is calculated using Equation 2 and data from Table 2. The value for  $f_{fd}$  for the conditions of the sculpin study is 0.986; for the trout study it is 0.9812.

Baseline BAFs are calculated using Equation 1 from the field BAFs in Table 3, the  $f_{fd}$  values above, and  $f_l$  values from Table 2, and are presented below in Table 4.

Table 4	
Baseline BAFs for HCBd	

Fish (Trophic Level)	Baseline BAF (L/kg)
Sculpin (3)	352,200
Rainbow trout (4)	43,940

The baseline BAF for trophic level 3 presented in Table 4 is slightly smaller than the value presented by U.S. EPA (Table 1). The difference is believed due to rounding and considered negligible. Because the two values are sufficiently close, and to be consistent with U.S. EPA guidelines for the Great Lakes, this fact sheet will use the Table 1 value of 354,800 L/kg.

## DERIVATION OF WATER QUALITY VALUE

As required by 6 NYCRR 702.8(a) the water quality value must equal the acceptable daily intake from fish consumption divided by a bioaccumulation factor and by a fish consumption rate of 0.033 kg/day.

### A. Acceptable Daily Intake From Fish Consumption

As required by 6 NYCRR 702.8(b), the most stringent acceptable daily intake from fish consumption is 20% of the ADI for nononcogenic effects, as determined from 6 NYCRR 702.5. That ADI is 0.067 ug HCBd/kg/day from NYS (1997) as described above.

The acceptable daily intake for fish consumption is:

$$0.2 \times 0.067 \text{ ug HCBd/kg/day} = 0.0134 \text{ ug HCBd/kg/day}$$

### B. Final Bioaccumulation Factors (BAF)

As described above, a baseline BAF is adjusted by the fish lipid fraction and the fraction freely dissolved to yield a final BAF for the substance. Equation 1 (above) is rearranged to solve for final BAF:

$$\text{Final BAF} = [(\text{baseline BAF})(f_l) + 1](f_{fd})$$

where values for  $f_l$  and  $f_{fd}$  are appropriate to criteria for New York State. Because, as described below, humans are exposed to fish from two trophic levels, this calculation is performed to generate final BAFs for trophic levels 3 and 4.

A fish lipid content of 3% had previously been used when calculating BAFs for deriving criteria for New York State. U.S. EPA (1995a) apportions daily fish consumption between fish of trophic levels 3 and 4. Specifically, 24% is assigned to trophic level 3 fish, with a standardized lipid fraction of 0.0182 (1.82%), and 76% to trophic level 4 fish, with a standardized lipid fraction of 0.0310 (3.1%). The weighted average lipid fraction of trophic level 3 and 4 fish is thus 0.028 (2.8%), which is very close to the value of 3% that had been used in New York State. U.S. EPA's apportionment approach is believed to be protective of human consumers of fish statewide, and will be used in the derivation of the water quality value in this fact sheet to achieve consistency with requirements for the Great Lakes System.

For deriving  $f_{fd}$  values for the Great Lakes, U.S. EPA (1995a) procedures use DOC and POC values of 2 and 0.04 mg/L respectively. The POC level of 0.04 mg/L is on the low end for the Great Lakes but U.S. EPA selected it to ensure protection throughout the System.

Data on levels of DOC and POC were examined for fresh and marine waters in New York State. Levels of DOC vary somewhat through the State but are fairly close to 2 mg/L. The  $f_{fd}$  is not very sensitive to changes in concentration of DOC. Levels of POC in New York State range from zero to several mg/L, but a sufficient number of near-zero values were found such that the level that EPA uses for the Great Lakes System seems appropriate for statewide standards and at the same time provides consistency with the federal requirements for the Great Lakes System.

Using these values for DOC and POC, equation 2 (above) becomes:

$$f_{fd} = \frac{1}{1 + (0.00000024 \text{ kg/L})(K_{ow})}$$

With a  $K_{ow}$  of 69,500, the fraction freely dissolved is calculated to be 0.984.

As described above, the baseline BAFs for HCBd for trophic levels 3 and 4 are 354,800 and 43,940 L/kg respectively.

The final BAF for trophic level 3 is calculated as:

$$\text{Final BAF}_{TL3} = [(\text{baseline BAF}_{TL3})(f_{lTL3}) + 1](f_{fd}) =$$

$$\text{Final BAF}_{TL3} = [(354,800)(0.0182)+1](0.984) = 6,360 \text{ L/kg}$$

The final BAF for trophic level 4 is calculated as:

$$\text{Final BAF}_{\text{TL4}} = [(\text{baseline BAF}_{\text{TL4}})(f_{\text{I TL4}}) + 1](f_{\text{fd}}) =$$

$$\text{Final BAF}_{\text{TL4}} = [(43,940)(0.0310) + 1](0.984) = 1,340 \text{ L/kg}$$

### C. Human Exposure (Fish Consumption)

6 NYCRR 702.8 requires that H(FC) values be based on a fish consumption rate of 0.033 kg/day.

### D. Calculation of Water Quality Value

The water quality value (WQV) is derived using a human body weight of 70 kg and a daily fish consumption rate of 0.033 kg as shown below. The fish consumption is apportioned as 24% trophic level 3 and 76% trophic level 4.

$$\text{WQV} = \frac{\text{Acceptable Daily Intake from Fish Consumption} \times 70 \text{ kg}}{[(\text{BAF}_{\text{TL3}})(0.24) + (\text{BAF}_{\text{TL4}})(0.76)] \times 0.033 \text{ kg/day}}$$

$$\text{WQV} = \frac{0.0134 \text{ ug HCBD/kg/day} \times 70 \text{ kg}}{[(6,360 \text{ L/kg})(0.24) + (1,340 \text{ L/kg})(0.76)] \times 0.033 \text{ kg/day}}$$

$$= 0.0112 \text{ ug/L, rounded to } 0.01 \text{ ug/L}$$

## REFERENCES

Oliver, B.G. and A.J. Niimi. 1983. Bioconcentration of chlorobenzenes from water by rainbow trout: Correlations with partition coefficients and environmental residues. Environ. Sci. Technol. 17:287-291.

Oliver, B.G. and A.J. Niimi. 1988. Trophodynamic analysis of polychlorinated biphenyl congeners and other chlorinated hydrocarbons in the Lake Ontario ecosystem. Environ. Sci. Technol. 22(4):388-397.

6 NYCRR (New York State Codes, Rules and Regulations). Water Quality Regulations, Surface Water and Groundwater Classifications and Standards: Title 6 NYCRR, Chapter X, Parts 700-705. Albany, NY: Department of Environmental Conservation.

NYS (New York State). 1997. Human Health Fact Sheet. Ambient Water Quality Value for Protection of Sources of Potable Water. Hexachlorobutadiene. Albany, NY: Department of Environmental Conservation.

Stephan, C.E. 1995. Personal communication from Dr. Charles E. Stephan, Environmental Research Lab, U.S. Environmental Protection Agency, Duluth, MN.

U.S. EPA (Environmental Protection Agency). 1995a. Final Water Quality Guidance for the Great Lakes System. 60 Federal Register: 15366-15425. March 23, 1995.

U.S. EPA (Environmental Protection Agency). 1995b. Great Lakes Water Quality Initiative Technical Support Document for the Procedure to Determine Bioaccumulation Factors. Office of Water. EPA-820-B95-005.

New York State Department of Environmental Conservation  
Division of Water  
SJS  
January 27, 1996