

20. References

Chapter 1

- Brown, R.; Barone, S.; Kimmel, C. (2008) Children's health risk assessment: Incorporating a lifestage approach into the risk assessment process. *Birth Defects Res B Dev Reprod Toxicol* (83)6:511–521.
- Georgopoulos, P.G. (2008) A multiscale approach for assessing the interactions of environmental and biological systems in a holistic health risk assessment framework. *Water Air Soil Pollut: Focus* 8:3–21.
- IPCS (International Programme on Chemical Safety). (2001) Glossary of exposure assessment-related terms: A compilation. IPCS Exposure Terminology Subcommittee, International Programme on Chemical Safety, World Health Organization. Available online at http://www.who.int/ipcs/publications/methods/harmonization/en/compilation_nov2001.pdf.
- Price, P.S.; Chaisson, C.F.; Koontz, M.; Ryan, B.; Wilkes, C.; Macintosh, D.; Georgopoulos, P.G. (2003) Construction of a comprehensive chemical exposure framework using person-oriented modeling. The Lifeline Group. Developed for The Exposure Technical Implementation Panel American Chemistry Council; Contract # 1388.
- U.S. EPA (Environmental Protection Agency). (1992) Guidelines for exposure assessment. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/Z-92/001. Available online at <http://www.epa.gov/raf/publications/guidelines-for-exposure-assessment.htm>.
- U.S. EPA (Environmental Protection Agency). (1997) Exposure factors handbook. National Center for Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/P-95/002Fa,b,c. Available from the National Technical Information Service, Springfield, VA, and online at <http://www.epa.gov/ncea/cfh/>.
- U.S. EPA (Environmental Protection Agency). (2005) Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington DC; EPA/630/P-03/003-F. Available at <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.
- U.S. EPA (Environmental Protection Agency). (2006) A framework for assessing health risk of environmental exposures to children. National Center for Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/R-05/093F. Available from the National Technical Information Service, Springfield, VA, and online at <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=158363>.
- U.S. EPA (Environmental Protection Agency). (2011) Exposure factors handbook: 2011 edition. National Center for Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/R-09/052F. Available from the National Technical Information Service, Springfield, VA, and online at <http://www.epa.gov/ncea>.

Chapter 2

- NRC (National Research Council). (1994) Science and judgment in risk assessment. National Academy Press, Washington, DC.
- U.S. EPA (Environmental Protection Agency). (1992) Guidelines for exposure assessment. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/Z-92/001. Available online at <http://www.epa.gov/raf/publications/guidelines-for-exposure-assessment.htm>.

Chapter 3

- Dufour, A.P.; Evans, O.; Behymer, T.D.; Cantu, R. (2006) Water ingestion during swimming activities in a pool: A pilot study. *J Water Health* 4(4):425–430.
- Kahn, H. (2008) Letter from Henry Kahn to Jacqueline Moya, EPA, providing supplemental data to Estimated daily average per capita water ingestion by child and adult age categories based on USDA's 1994–96 and 1998 continuing survey of food intakes by individuals (September 18, 2008).
- Kahn, H.; Stralka, K. (2008a) Estimated daily average per capita water ingestion by child and adult age categories based on USDA's 1994–96 and 1998 continuing survey of food intakes by individuals. *J Exp Sci Environ Epidemiol* 19(4):396–404.
- Kahn, H.; Stralka, K. (2008b) Estimates of water ingestion for women in pregnant, lactating, and non-pregnant and non-lactating child-bearing age groups based on USDA's 1994–96, 1998 continuing survey of food intakes by individuals. *Hum Ecol Risk Assess Int J* 14(6):1273–1290.
- NCHS (National Center for Health Statistics). (1993) Joint policy on variance estimation and statistical reporting standards on NHANES III and CSFII reports: HNIS/NCHS Analytic Working Group recommendations. Human Nutrition Information Service (HNIS)/Analytic Working Group. Agricultural Research Service, Survey Systems/Food Consumption Laboratory, Riverdale, MD.

Chapter 4

- Beamer, P.; Key, M.E.; Ferguson, A.C.; Canales, R.A.; Auyeung, W.; Leckie, J.O. (2008) Quantified activity pattern data from 6 to 27-month-old farmworker children for use in exposure assessment. *Environ Res* 108(2):239–246.
- Greene, M.A. (2002) Mouthing times for children from the observational study. U.S. Consumer Product Safety Commission, Bethesda, MD.
- Groot, M.E.; Lekkerkerk, M.C.; Steenbekkers, L.P.A. (1998) Mouthing behavior of young children: An observational study. Wageningen Agricultural University, Wageningen, The Netherlands.
- Juberg, D.R.; Alfano, K.; Coughlin, R.J.; Thompson, K.M. (2001) An observational study of object mouthing behavior by young children. *Pediatrics* 107(1):135–142.
- Xue, J.; Zartarian, V.; Moya, J.; Freeman, N.; Beamer, P.; Black, K.; Tulve, N.; Shalat, S. (2007) A meta-analysis of children's hand-to-mouth frequency data for estimating nondietary ingestion exposure. *Risk Anal* 27(2):411–420.
- Xue, J.; Zartarian, V.; Tulve, N.; Moya, J.; Freeman, N.; AuYeung, W.; Beamer, P. (2009) A meta-analysis of children's object-to-mouth frequency data for estimating nondietary ingestion exposure. Accepted for publication in the *J Expo Sci Environ Epidemiol*.

Chapter 5

- ATSDR (Agency for Toxic Substances and Disease Registry). (2001) Summary report for the ATSDR soil-pica workshop. ATSDR, Atlanta, GA. March 20, 2001. Available online at: <http://www.atsdr.cdc.gov/NEWS/soilpica.html>.
- Barnes, R.M. (1990) Childhood soil ingestion: How much dirt do kids eat? *Anal Chem* 62:1024–1033.
- Calabrese, E.J.; Stanek, E.J. (1993) Soil pica: Not a rare event. *J Environ Sci Health A28*(2):373–384.
- Calabrese, E.J.; Stanek, E.J. (1995) Resolving intertracer inconsistencies in soil ingestion estimation. *Environ Health Perspect* 103(5):454–456.

- Calabrese, E.J.; Barnes, R.; Stanek, E.J. III; Pastides, H.; Gilbert, C.; Veneman, P.; Wang, X.; Lásztity, A.; Kostecki, P.T. (1989) How much soil do young children ingest: An epidemiologic study. In: Petroleum Contaminated Soils, Lewis Publishers, Chelsea, MI; pp. 363–397 and Regul Toxicol Pharm 10:123–137.
- Calabrese, E.J.; Stanek, E.J.; Gilbert, C.E. (1991) Evidence of soil-pica behavior and quantification of soil ingested. Hum Exp Toxicol 10:245–249.
- Calabrese, E.J.; Stanek, E.J.; Pekow, P.; Barnes, R.M. (1997a) Soil ingestion estimates for children residing on a Superfund site. Ecotoxicol Environ Saf 36:258–268.
- Calabrese, E.J.; Stanek, E.J.; Barnes, R.M. (1997b) Soil ingestion rates in children identified by parental observation as likely high soil ingesters. J Soil Contam 6(3):271–279.
- Davis, S.; Mirick, D. (2006) Soil ingestion in children and adults in the same family. J Exp Anal Environ Epidemiol 16:63–75.
- Davis, S.; Waller, P.; Buschbom, R.; Ballou, J.; White, P. (1990) Quantitative estimates of soil ingestion in normal children between the ages of 2 and 7 years: Population based estimates using aluminum, silicon, and titanium as soil tracer elements. Arch Environ Health 45:112–122.
- Hogan, K.; Marcus, A.; Smith, R.; White, P. (1998) Integrated exposure uptake biokinetic model for lead in children: Empirical comparisons with epidemiologic data. Environ Health Perspect 106(6):1557–1567.
- Özkaynak, H., Xue, J., Zartarian, V.G., Glen, G., Smith, L. (2010) Modeled estimates of soil and dust ingestion rates for children. Risk Anal online prepublication copy.
- Stanek, E.J.; Calabrese, E.J. (1995) Soil ingestion estimates for use in site evaluations based on the best tracer method. Hum Ecol Risk Assess 1:133–156.
- Stanek, E.J.; Calabrese, E.J.; Mundt, K.; Pekow, P.; Yeatts, K.B. (1998) Prevalence of soil mouthing/ingestion among healthy children aged 1 to 6. J Soil Contam 7(2):227–242.
- Van Wijnen, J.H.; Clausing, P.; Brunekreff, B. (1990) Estimated soil ingestion by children. Environ Res 51:147–162.
- Vermeer, D.E.; Frate, D.A. (1979) Geophagia in rural Mississippi: Environmental and cultural contexts and nutritional implications. Am J Clin Nutr 32:2129–2135.
- Wong, M.S. (1988) The role of environmental and host behavioural factors in determining exposure to infection with *Ascaris lumbricoides* and *Trichuris trichiura*. Ph.D. Thesis, Faculty of Natural Sciences, University of the West Indies.

Chapter 6

- Arcus-Arth, A; Blaisdell, R.J. (2007) Statistical distributions of daily breathing rates for narrow age groups of infants and children. Risk Anal 27(1):97–110.
- Brochu, P.; Ducré-Robitaille, J.; Brodeur, J. (2006) Physiological daily inhalation rates for free-living individuals aged 1 month to 96 years, using data from doubly labeled water measurements: A proposal for air quality criteria, standard calculations and health risk assessment. Hum Ecol Risk Assess 12:675–701.
- Stifelman, M. (2007) Using doubly-labeled water measurements of human energy expenditure to estimate inhalation rates. Sci Total Environ 373:585–590.

U.S. EPA (Environmental Protection Agency). (1992) Guidelines for exposure assessment. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/Z-92/001. Available online at <http://www.epa.gov/raf/publications/guidelines-for-exposure-assessment.htm>.

U.S. EPA (Environmental Protection Agency). (2009) Metabolically-derived human ventilation rates: A revised approach based upon oxygen consumption rates. National Center for Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/R-06/129F. Available from the National Technical Information Service, Springfield, VA.

Chapter 7

Boniol, M.; Verriest, J.P.; Perdeux, R.; Dore, J-F. (2007) Proportion of skin surface area of children and young adults from 2 to 18 years old. *J Investig Dermatol* 128:461–464.

Holmes, Jr., K.K.; Shirai, J.H.; Richter, K.Y.; Kissel, J.C. (1999) Field measurement of dermal loadings in occupational and recreational activities. *Environ Res* 80:148–157, Section A.

Kissel, J.C.; Richter, K.; Fenske, R. (1996) Field measurements of dermal soil loading attributable to various activities: Implications for exposure assessment. *Risk Anal* 16(1):116–125.

Shoaf, M.B.; Shirai, J.H.; Kedan, G.; Schaum, J.; Kissel, J.C. (2005) Child dermal sediment loads following play in a tide flat. *J Expo Anal Environ Epidemiol* 15:407–412.

U.S. EPA (Environmental Protection Agency). (1985) Development of statistical distributions or ranges of standard factors used in exposure assessments. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/8-85/010. Available from the National Technical Information Service, Springfield, VA, PB85-242667.

U.S. EPA (Environmental Protection Agency). (1992a) Guidelines for exposure assessment. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/Z-92/001. Available online at <http://www.epa.gov/raf/publications/guidelines-for-exposure-assessment.htm>.

U.S. EPA (Environmental Protection Agency). (1992b) Dermal exposure assessment: Principles and applications. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; U.S. EPA/600/8-91/91. Available from the National Technical Information Service, Springfield, VA, and online at <http://www.epa.gov/ncea>.

U.S. EPA (Environmental Protection Agency). (2004) Risk assessment guidance for Superfund (RAGS), Volume I: Human health evaluation manual, Part E. Washington, DC; EPA/540/R-99/005.

U.S. EPA (Environmental Protection Agency). (2005) Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington DC; EPA/630/P-03/003-F. Available at <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.

Chapter 8

U.S. EPA (Environmental Protection Agency). (1989) Risk assessment guidance for Superfund, Volume I: Human health evaluation manual. Office of Emergency and Remedial Response, Washington, DC; EPA/540/1-89/002.

U.S. EPA (Environmental Protection Agency). (2005) Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington DC; EPA/630/P-03/003-F. Available at <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.

Chapter 9

NCHS (National Center for Health Statistics). (1993) Joint policy on variance estimation and statistical reporting standards on NHANES III and CSFII reports: HNIS/NCHS Analytic Working Group recommendations. Human Nutrition Information Service (HNIS)/Analytic Working Group. Agricultural Research Service, Survey Systems/Food Consumption Laboratory, Riverdale, MD.

U.S. EPA (Environmental Protection Agency). (2005) Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington DC; EPA/630/P-03/003-F. Available at <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.

Chapter 10

NCHS (National Center for Health Statistics). (1993) Joint policy on variance estimation and statistical reporting standards on NHANES III and CSFII reports: HNIS/NCHS Analytic Working Group recommendations. Human Nutrition Information Service (HNIS)/Analytic Working Group. Agricultural Research Service, Survey Systems/Food Consumption Laboratory, Riverdale, MD.

NMFS (National Marine Fisheries Service). (1993) Data tapes for the 1993 NMFS provided to U.S. EPA. National Center for Environmental Assessment.

U.S. EPA (Environmental Protection Agency). (2005) Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington DC; EPA/630/P-03/003-F. Available at <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.

Chapter 11

NCHS (National Center for Health Statistics). (1993) Joint policy on variance estimation and statistical reporting standards on NHANES III and CSFII reports: HNIS/NCHS Analytic Working Group recommendations. Human Nutrition Information Service (HNIS)/Analytic Working Group. Agricultural Research Service, Survey Systems/Food Consumption Laboratory, Riverdale, MD.

U.S. EPA (Environmental Protection Agency). (2005) Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington DC; EPA/630/P-03/003-F. Available at <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.

U.S. EPA (Environmental Protection Agency). (2007) Analysis of fat intake based on the U.S. Department of Agriculture's 1994–96, 1998 continuing survey of food intakes by individuals (CSFII). National Center for Environmental Assessment, Washington, DC; EPA/600/R-05/021F. Available from the National Technical Information Service, Springfield, VA, and online at <http://www.epa.gov/ncea>.

Chapter 12

NCHS (National Center for Health Statistics). (1993) Joint policy on variance estimation and statistical reporting standards on NHANES III and CSFII reports: HNIS/NCHS Analytic Working Group recommendations. Human Nutrition Information Service (HNIS)/Analytic Working Group. Agricultural Research Service, Survey Systems/Food Consumption Laboratory, Riverdale, MD.

U.S. EPA (Environmental Protection Agency). (2005) Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington DC; EPA/630/P-03/003-F. Available at <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.

Chapter 13

- Phillips, L.; Moya, J. (2011) Estimation of age-specific per capita home-produced food intake among populations that garden, farm, or raise animals. *J Expo Anal Environ Epidemiol* advance online publication: 27 April 2011:1–8.
- USDA (U.S. Department of Agriculture). (1987–1988) Dataset: Nationwide food consumption survey 1987/88 household food use. Washington, DC. 1987/88 NFCS Database.
- U.S. EPA (Environmental Protection Agency). (1989) Risk assessment guidance for Superfund (RAGS), Volume I: Human health evaluation manual, Part A. Office of Solid Waste and Emergency Response, Washington, DC; EPA/540/1-89/002. Available online at <http://www.epa.gov/oswer/riskassessment/ragsa/index.htm>.
- U.S. EPA (Environmental Protection Agency). (1996) Soil screening fact sheet guidance. Office of Solid Waste and Emergency Response, Washington, DC; EPA/540/F-95/041. Available online at <http://www.epa.gov/superfund/health/conmedia/soil/index.htm>.
- U.S. EPA (Environmental Protection Agency). (2005) Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. Risk Assessment Forum, Office of Research and Development, Washington DC; EPA/630/P-03/003-F. Available at <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.

Chapter 14

- NCHS (National Center for Health Statistics). (1993) Joint policy on variance estimation and statistical reporting standards on NHANES III and CSFII reports: HNIS/NCHS Analytic Working Group recommendations. Human Nutrition Information Service (HNIS)/Analytic Working Group. Agricultural Research Service, Survey Systems/Food Consumption Laboratory, Riverdale, MD.
- USDA (U.S. Department of Agriculture). (2000) 1994–96, 1998 continuing survey of food intakes by individuals (CSFII). CD-ROM. Agricultural Research Service, Beltsville Human Nutrition Research Center, Beltsville, MD. Available from the National Technical Information Service, Springfield, VA; PB-2000-500027.
- U.S. EPA (Environmental Protection Agency). (2007) Analysis of total food intake and composition of individual's diet based on USDA's 1994–96, 1998 continuing survey of food intakes by individuals (CSFII). National Center for Environmental Assessment, Washington, DC; EPA/600/R-05/062F. Available from the National Technical Information Service, Springfield, VA, and online at <http://www.epa.gov/ncea>.

Chapter 15

- AAP (American Academy of Pediatrics). (2005) Breast feeding and the use of human milk. Policy statement. *Pediatrics*. 115(2):496–506. Available online at <http://aappolicy.aappublications.org/cgi/content/full/pediatrics;115/2/496>.
- Arcus-Arth, A.; Krowech, G.; Zeise, L. (2005) Human milk and lipid intake distributions for assessing cumulative exposure and risk. *J Expos Anal Environ Epidemiol* 15:357–365.
- Butte, N.F.; Garza, C.; Smith, E.O.; Nichols, B.L. (1984) Human milk intake and growth in exclusively breast-fed infants. *J Pediatr* 104:187–195.
- Butte, N.; Wong, W.; Hopkinson, J.; Smith E.; Ellis, J. (2000) Infant feeding mode affects early growth and body composition. *Pediatrics* 106:1355–1366.
- Chen, A.; Rogan, W.J. (2004) Breastfeeding and the risk of postneonatal death in the United States. *Pediatrics* 113:435–439.
- Dewey, K.G.; Lönnerdal, B. (1983) Milk and nutrient intake of breast-fed infants from 1 to 6 months: Relation to growth and fatness. *J Pediatr Gastroenterol Nutr* 2:497–506.

- Dewey, K.G.; Heinig, J.; Nommsen, L.A.; Lönnerdal, B. (1991a) Maternal versus infant factors related to human milk intake and residual volume: The DARLING study. *Pediatrics* 87:829–837.
- Dewey, K.G.; Heinig, J.; Nommsen, L.; Lönnerdal, B. (1991b) Adequacy of energy intake among breast-fed infants in the DARLING study: Relationships to growth, velocity, morbidity, and activity levels. *J Pediatr* 119:538–547.
- Neville, M.C.; Keller, R.; Seacat, J.; Lutes, V.; Neifert, M.; et al. (1988) Studies in human lactation: Milk volumes in lactating women during the onset of lactation and full lactation. *Am J Clin Nutr* 48:1375–1386.
- Pao, E.M.; Hines, J.M.; Roche, A.F. (1980) Milk intakes and feeding patterns of breast-fed infants. *J Am Diet Assoc* 77:540–545.

Chapter 16

- Carey, M. (1988) Occupational tenure in 1987: Many workers have remained in their fields. *Mon Labor Rev* October 1988:3–12.
- Carey, M. (1990) Occupational tenure, employer tenure, and occupational mobility. *Occup Outlook Quart* Summer 1990:55–60.
- Johnson, T.; Capel, J. (1992) A monte carlo approach to simulating residential occupancy periods and its application to the general U.S. population. U.S. EPA, Office of Air Quality and Standards, Research Triangle Park, NC.
- U.S. Bureau of the Census. (2008) American housing survey for the United States in 2007. U.S. Government Printing Office, Washington, DC.
- U.S. EPA (Environmental Protection Agency). (1996) Descriptive statistics tables from a detailed analysis of the National Human Activity Pattern Survey (NHAPS) data. Office of Research and Development, Washington, DC; EPA/600/R-96/148.
- Wiley, J.A.; Robinson, J.P.; Cheng, Y.; Piazza, T.; Stork, L.; Plasden, K. (1991) Study of children's activity patterns. California Environmental Protection Agency, Air Resources Board Research Division, Sacramento, CA.

Chapter 17

- Abt Associates. (1992) Methylene chloride consumer products use survey findings. Prepared by Abt Associates, Inc. for the U.S. Consumer Product Safety Commission, Bethesda, MD.
- Bass, J.; Ortega, L.; Rosales, C.; Petersen, N.; Philen, R. (2001) What's being used at home: A household pesticide survey. *Pub Health* 9(3):138–144.
- CTFA (Cosmetic, Toiletry and Fragrance Association). (1983) Summary of the results of surveys of the amount and frequency of use of cosmetic products by women. Prepared by Environ Corporation, Washington, DC, for CTFA Inc., Washington, DC.
- Franklin, P. (2008) Household chemicals: Good housekeeping or occupational hazard. *Eur Respir J* 31:489–491.
- Hall, B.; Tozer, S; Safford, B; Coroama, M; Steiling, W; Leneveu-Duchemin, MC; McNamara C.; Gibney, M. (2007) European consumer exposure to cosmetic products, a framework for conducting population exposure assessments. *Food Chem Toxicol* 45(11):2097–2108.
- Loretz, L.; Api, A.; Barraji, L.; Burdick, J.; Dressler, W.; Gettings, S.; Hsu, H.; Pan, Y.; Re, T.; Renskers, K.; Rothenstein, A.; Scrafford, C.; Sewall, C. (2005) Exposure data for cosmetic products: Lipstick, body lotion, and face cream. *Food Chem Toxicol* 43:279–291.

- Loretz, L.; Api, A.; Barraji, L.; Burdick, J.; Davis, D.; Dressler, W.; Gilberti, E.; Jarrett, G.; Mann, S.; Pan, Y.; Re, T.; Renskers, K.; Scrafford, C.; Vater, S. (2006) Exposure data for personal care products: Hairspray, spray perfume, liquid foundation, shampoo, body wash, and solid antiperspirant. *Food Chem Toxicol* 44:2008–2018.
- Loretz, L.; Api, A.; Babcock, L.; Barraji, L.; Burdick, J.; Cater, K.; Jarrett, G.; Mann, S.; Pan, Y.; Re, T.; Renskers, K.; Scrafford, C. (2008) Exposure data for cosmetic products: Facial cleanser, hair conditioner, and eye shadow. *Food Chem Toxicol* 46:1516–1524.
- Sathyanarayana, S.; Karr, C.; Lozano, P.; Brown, E.; Calafat, M. (2008) Baby care products: Possible sources of infant phthalate exposure. *Pediatrics* 121:260–268.
- Steenbekkers, L.P. (2001) Methods to study everyday use of products in households: The Wageningen mouthing study. *Am Occup Hyg* 45(1001):125–129.
- U.S. EPA (Environmental Protection Agency). (1996) Descriptive statistics tables from a detailed analysis of the National Human Activity Pattern Survey (NHAPS) data. Office of Research and Development, Washington, DC; EPA/600/R-96/148.
- Weegels, M.F.; van Veen, M.P. (2001) Variation of consumer contact with household products: A preliminary investigation. *Risk Anal* 21(3):499–511.
- Westat. (1987a) Household solvent products: A national usage survey. Under Subcontract to Battelle Columbus Div., Washington DC. Prepared for U.S. Environmental Protection Agency, Washington, DC. Available from the National Technical Information Service, Springfield, VA; PB-88-132881.
- Westat. (1987b) National usage survey of household cleaning products. Prepared for U.S. Environmental Protection Agency, Office of Toxic Substances and Office of Pesticides and Toxic Substances, Washington, DC. Available from the National Technical Information Service, Springfield, VA.
- Westat. (1987c) National household survey of interior painters. Prepared for U.S. Environmental Protection Agency, Office of Toxic Substances and Office of Pesticides and Toxic Substances, Washington DC.

Chapter 18

- Xu, J., Kochanek, K., Murphy, S., and Tejada-Vera, B. (2010, May 20) Deaths: Final data for 2007. *National Vital Statistics Reports*, 58(19). Available online at http://www.cdc.gov/NCHS/data/nvsr/nvsr58/nvsr58_19.pdf.

Chapter 19

- Koontz, M.D.; Rector, H.E. (1995) Estimation of distributions for residential air exchange rates. EPA Contract No. 68-D9-0166, Work Assignment No. 3-19, U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics, Washington, DC.
- Turk, B.H., Brown, J.T., Geisling-Sobotka, K., Froehlich, D.A., Grimsrun, D.T., Harrison, J., Koonce, J.F., Prill, R.J., and Revzan, K.L. (1987) Indoor Air Quality and Ventilation Measurements in 38 Pacific Northwest commercial Buildings. Volume 1: Measurement Results and Interpretation. Final Report. Prepared for U.S. DOE. DE-AC03-76SF00098.
- U.S. DOE (Department of Energy). (2008a) Residential Energy Consumption Survey (RECS). Report No. DOE/EIA-0314 (93), U.S. Department of Energy, Energy Information Administration, Washington, DC.
- U.S. DOE (Department of Energy). (2008b) Commercial Buildings Energy Consumption Survey (CBECS). Form EIA-871A. U.S. Department of Energy, Energy Information Administration, Washington, DC. Available at <http://www.eia.doe.gov/emeu/cbecs>.