

## ASTHMA FACTS

### **Asthma continues to be a serious public health problem in the United States.**

According to the U.S. Centers for Disease Control and Prevention (CDC)'s 2015 and 2016 National Health Interview Surveys (NHIS):

- An estimated 24.6 million people,<sup>1</sup> including 6.1 million children,<sup>2</sup> have asthma.
- More than 11.5 million people with asthma, including nearly 3 million children, report having had one or more asthma attacks in 2015.<sup>3</sup>

Uncontrolled asthma is a common reason people seek medical attention.

- The 2010 National Hospital Ambulatory Medical Care Survey reported 1.3 million outpatient department visits with asthma as the primary diagnosis.<sup>4</sup>
- The 2012 National Ambulatory Medical Care Survey reported 10.5 million physician office visits with asthma as the primary diagnosis.<sup>5</sup>
- The National Hospital Ambulatory Medical Care Survey: 2014 Emergency Department Summary Tables showed that asthma was the primary diagnosis for more than 2.0 million emergency department visits.<sup>6</sup>
- The 2011-2012 National Survey for Children's Health reported that children who have asthma (3.4%) are more likely to use a hospital emergency department as their usual place for medical care than children without asthma (2.1%).<sup>7</sup>

Asthma is a common chronic disease in children.

- In 2016, the prevalence of asthma in children was 8.3 percent, meaning about 1 in 12 children had asthma.<sup>2</sup>
- In 2013, approximately 13.8 million missed school days were reported due to asthma.<sup>8</sup>

Non-Hispanic Blacks have a higher asthma mortality rate than people of other races or ethnicities.

- According to the CDC's 2015 summary of asthma mortality data, Black Americans have a higher asthma death rate - at 23.9 deaths per million persons - than non-Hispanic whites (8.4 deaths per million persons), Hispanics (7.3 deaths per million persons), and other non-Hispanics (10.0 deaths per million persons).<sup>9</sup>

### **The economic costs of asthma are high.**

- The annual economic cost of asthma in 2007, including medical costs and lost school and work days, amounted to more than \$56 billion.<sup>10</sup>
- In 2012, the median annual medical cost of asthma was \$983 per child, with a range of \$833 in Arizona to \$1,121 in Michigan.<sup>11</sup>

### **Reducing exposure to environmental factors, such as indoor asthma triggers, is important for asthma management.**

- On average, Americans spend about 90 percent of their time indoors.<sup>12</sup>
- Indoor environmental factors called asthma triggers - such as dust mites, mold, cockroaches, pet dander and secondhand smoke - can exacerbate asthma symptoms.<sup>13</sup>
- With an asthma action plan that includes medical treatment and control of environmental triggers, people with asthma can lead healthy, active lives.<sup>14</sup>

## References

1. CDC. 2015. *NHIS Data; Table 3-1*. [www.cdc.gov/asthma/nhis/2015/table3-1.htm](http://www.cdc.gov/asthma/nhis/2015/table3-1.htm).
2. CDC. 2018. *Morbidity and Mortality Weekly Report: Vital Signs: Asthma in Children — United States, 2001–2016*. <https://www.cdc.gov/mmwr/volumes/67/wr/mm6705e1.htm>.
3. CDC. 2015. *NHIS Data; Table 5-1*. [www.cdc.gov/asthma/nhis/2015/table5-1.htm](http://www.cdc.gov/asthma/nhis/2015/table5-1.htm).
4. CDC. 2010. *National Hospital Ambulatory Medical Care Survey: 2010 Outpatient Department Summary Tables; Table 11*. [www.cdc.gov/nchs/data/ahcd/nhamcs\\_outpatient/2010\\_opd\\_web\\_tables.pdf](http://www.cdc.gov/nchs/data/ahcd/nhamcs_outpatient/2010_opd_web_tables.pdf).
5. CDC. 2012. *National Ambulatory Medical Care Survey: 2012 State and National Summary Tables; Table 16*. [www.cdc.gov/nchs/data/ahcd/namcs\\_summary/2012\\_namcs\\_web\\_tables.pdf](http://www.cdc.gov/nchs/data/ahcd/namcs_summary/2012_namcs_web_tables.pdf).
6. CDC. 2015. *National Hospital Ambulatory Medical Care Survey: 2014 Emergency Department Summary Tables; Table 12*. [https://www.cdc.gov/nchs/data/nhamcs/web\\_tables/2014\\_ed\\_web\\_tables.pdf](https://www.cdc.gov/nchs/data/nhamcs/web_tables/2014_ed_web_tables.pdf).
7. CDC. November 2016. *Usual Place for Medical Care Among Children*. Accessed April 24, 2018. [www.cdc.gov/asthma/asthma\\_stats/usualplaceforcare.htm](http://www.cdc.gov/asthma/asthma_stats/usualplaceforcare.htm).
8. CDC. October 2015. “Asthma-Related Missed School Days Among Children Aged 5–17 Years.” *AsthmaStats Factsheet*. Accessed April 7, 2017. [www.cdc.gov/asthma/asthma\\_stats/missing\\_days.htm](http://www.cdc.gov/asthma/asthma_stats/missing_days.htm).
9. CDC. 2015. *Most Recent Asthma Data; Mortality Table*. Accessed April 24, 2018. [www.cdc.gov/asthma/most\\_recent\\_data.htm](http://www.cdc.gov/asthma/most_recent_data.htm).
10. CDC. 2011. “Asthma in the U.S.” *CDC Vital Signs*. [www.cdc.gov/vitalsigns/asthma](http://www.cdc.gov/vitalsigns/asthma).
11. Nurmagambetov T., Khavjou O., Murphy L., Orenstein D. “State-level medical and absenteeism cost of asthma in the United States.” 2017. *Journal of Asthma*; 54:357–70. <https://doi.org/10.1080/02770903.2016.1218013>.
12. Klepeis, N.E., Nelson, W.C., Ott, W.R., Robinson, J.P., Tsang, A.M., Switzer, P., Behar, J.V., Hern, S.C., Engelmann, W.H. 2001. “The National Activity Pattern Survey: A Resource for Assessing Exposure to Environmental Pollutants.” *Journal of Exposure Analysis and Environmental Epidemiology*; 11(3):231-52.
13. Kanchongkittiphon, W., Mendell, M.J., Gaffin, J.M., Wang, G., Phipatanakul, W. January 2015. “Indoor Environmental Exposures and Exacerbation of Asthma: An Update to the 2000 Review by the Institute of Medicine.” *Environmental Health Perspectives*; 123:6–20. <https://ehp.niehs.nih.gov/1307922>.
14. Matsui, E.C., Abramson, S.L., Sandel, M.T., Section on Allergy and Immunology, Council on Environmental Health. November 2016. “Indoor Environmental Control Practices and Asthma Management.” *Pediatrics*; 135(5): e1-e11. <http://pediatrics.aappublications.org/content/138/5/e20162589>.