

Risk Evaluation for Trichloroethylene

Systematic Review Supplemental File:

Data Quality Evaluation of Environmental Releases and Occupational Exposure Common Sources

CASRN: 79-01-6

February 2020

This document is a compilation of tables for the data extraction and evaluation of common sources for environmental releases and occupational exposure of the first 10 chemicals. This document may contain sources that were not used for the risk evaluation of Trichloroethylene. Each table shows the data point or set or information element that was extracted evaluated from data source in accordance with and a Appendix D of the Application of Systematic Review in TSCA Risk Evaluations. If the source contains more than one data set or information element, the review provides an overall confidence score for each data set or information element that is found in the source. Therefore, it is possible that a source may have more than one overall quality/confidence score.

Table of Contents

	Page
Releases to the Environment	3
Occupational Exposure	11
Facility	51

Explanatory Notes

These explanatory notes provide context to understand the short comments in the data evaluation tables.

Domain	Metric	Description of Comments Field
Reliability	Methodology	Indicates the sampling/analytical methodology, estimation method, or type of publication
Representativeness	Geographic Scope	Indicates the country of the study, publication, or underlying data
	Applicability	Indicates whether the data are for a condition of use within scope of the Risk Evaluation
	Temporal Representativeness	Provides the year of study, publication, or underlying data
	Sample Size	Describes the distribution of the sample or underlying data
Accessibility / Clarity	Metadata Completeness	Describes the completeness of the metadata
Variability and Uncertainty	Metadata Completeness	Indicates if study or publication addresses variability and uncertainty of the data or information

Releases to the Environment

Source Citation: Type of Data Source Hero ID		2017. Toxics Release Inventory o the Environment; Environmen			ar 2016.	
EXTRACTION Parameter			Data			
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Environmental Media: Release or Emission Factor:			All All Provides Provides			
EVALUATION						
Domain		Metric	Rating	MWF^{\star}	Score	Comments
Domain 1: Reliab	oility Metric 1:	Methodology	Low	× 1	3	Methodology used by submitters to estimate release data is not known.
Domain 2: Repres	sentative					
Domain 2. Ropros	Metric 2:	Geographic Scope	High	\times 1	1	TRI is U.S. based data
	Metric 3:	Applicability	High	\times 2	2	\ensuremath{TRI} includes industries included in the scopes of multiple chemicals
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	TRI data are from 2016
	Metric 5:	Sample Size	Medium	× 1	2	Due to reporting requirements, statistical representativeness is unclear. $$
Domain 3: Access	sibility/Clar	ity				
-	Metric 6:	Metadata Completeness	Low	× 1	3	TRI only includes release media but no other metadata.
Domain 4: Variab						
	Metric 7:	Metadata Completeness	Low	× 1	3	\ensuremath{TRI} does not address variability or uncertainty in submitter provided data.
Overall Quality D	eterminatio	$ m n^{\dagger}$	Medium		1.8	

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

			()			(818 1 1 1 1 1 1 1 1 1
Source Citation:	U.S. EPA. year 2017.	2017. Toxics Release Inventory	(TRI) basic	plus data	a file, He	exabromocyclododecane (CAS $\#$ 25637-99-4), reporting
Type of Data Source		the Environment; Environmen	tal Release	Data:		
Hero ID	5079078	, , , , , , , , , , , , , , , , , , ,	our recrease	2 ava,		
EXTRACTION						
Parameter			Data			
Life Cycle Stage:			All			
Life Cycle Descrip	otion (Subca	ategory of Use):	All			
Environmental M		, , , , , , , , , , , , , , , , , , ,	Provides	media of	release	
Release or Emissi	on Factor:		Provides	release da	ata	
EVALUATION						
Domain		Metric	Rating	MWF*	Score	Comments
Domain 1: Reliab	ility					
	Metric 1:	Methodology	Low	× 1	3	Methodology used by submitters to estimate release data is not known. $$
Domain 2: Repres	sentative					
Bolliam 2. Tepro.	Metric 2:	Geographic Scope	High	$\times 1$	1	TRI is U.S. based data
	Metric 3:	Applicability	High	\times 2	2	\ensuremath{TRI} includes industries included in the scopes of multiple chemicals
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	TRI data are from 2017
	Metric 5:	Sample Size	Medium	× 1	2	Due to reporting requirements, statistical representativeness is unclear. $$
Domain 3: Access	sibility/Clar	itv				
	Metric 6:	Metadata Completeness	Low	× 1	3	TRI only includes release media but no other metadata.
Domain 4: Variab	oility and Ur	ncertainty				
	Metric 7:	Metadata Completeness	Low	× 1	3	\ensuremath{TRI} does not address variability or uncertainty in submitter provided data.
Overall Quality D	eterminatio	\mathbf{n}^{\dagger}	Medium		1.8	

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID		2016. EPA Discharge Monitoring the Environment; Environment				
EXTRACTION Parameter			Data			
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Environmental Media: Release or Emission Factor:		All All Provides Provides				
EVALUATION						
Domain		Metric	Rating	MWF^{\star}	Score	Comments
Domain 1: Reliab	oility Metric 1:	Methodology	Low	× 1	3	Methodology used by submitters to estimate release data is not known. $\label{eq:methodology}$
Domain 2: Repres	sentative					
Domain 2. Repres	Metric 2:	Geographic Scope	High	\times 1	1	DMR is U.S. based data
	Metric 3:	Applicability	High	\times 2	2	DMR includes industries included in the scopes of multiple chemicals
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	DMR data are from 2016
	Metric 5:	Sample Size	Medium	× 1	2	Universe is limited to NPDES permit holders; statistical representativeness is unclear. $$
Domain 3: Access	sibility/Clar	ity				
	Metric 6:	Metadata Completeness	Low	× 1	3	DMR only includes release media but no other metadata.
Domain 4: Variab	oility and U	ncertainty				
2011011 1. (0110).	Metric 7:	Metadata Completeness	Low	× 1	3	DMR does not address variability or uncertainty in submitter provided data.
Overall Quality D	Determinatio	n^\dagger	Medium		1.8	

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

	2018. 2014 National Emissions of the Environment; Environment				
EXTRACTION					
Parameter		Data			
Life Cycle Stage: Life Cycle Description (Subca Release Source: Environmental Media: Release or Emission Factor: Release Days per Year: P2 Control & percent Efficier	All Provides unit/process of release. Provides media of release Provides release data Provides annual operating time. Provides controls information.				
EVALUATION Domain	Metric	Rating	MWF*	Score	Comments
Domain 1: Reliability Metric 1:	Methodology	Medium	× 1	2	Submitters provide general method used to calculate emissions, but details not provided.
Domain 2: Representative					
Metric 2:	Geographic Scope	High	$\times 1$	1	NEI is U.S. based data
Metric 3:	Applicability	High	× 2	2	NEI includes industries included in the scopes of multiple chemicals.
Metric 4:	Temporal Representativeness	High	$\times 2$	2	NEI data are from 2014
Metric 5:	Sample Size	Medium	× 1	2	Universe is limited to units subject to NESHAP with threshold potential to emit, although states may have different requirements; statistical representativeness is unclear.
Domain 3: Accessibility/Clar	ity.				
Metric 6:	Metadata Completeness	High	× 1	1	NEI includes release media and generally also includes daily and annual operating time, specific unit/process that is the source of release, and presence of engineering controls.
Domain 4: Variability and U	ncortainty				
	Metadata Completeness	Low	× 1	3	NEI does not address variability or uncertainty in submitter provided data.
Overall Quality Determination	${f n}^{\dagger}$	High		1.4	
	Cor	ntinued on 1	next page	;	

- continued from previous page

	· ·				
Metric	Rating	\mathbf{MWF}^{\star}	Score	Comments	
	o the Environment; Environ	o the Environment; Environmental Release	2018. 2014 National Emissions Inventory Report. o the Environment; Environmental Release Data; Metric Rating MWF*	o the Environment; Environmental Release Data;	o the Environment; Environmental Release Data;

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID		1995. Protocol for Equipment I o the Environment; Environmen			nates. E	EPA-453/R-95-017.	
EXTRACTION Parameter			Data				
Life Cycle Stage: Release or Emission Factor:			Tank Truck and Railcar Loading Model Cited for emission factors used for the inhalation exposure and release model.				
EVALUATION							
Domain		Metric	Rating	MWF*	Score	Comments	
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	EPA coordinated the data gathering activities; methodology expected to be accurate and comprehensive of all leak release sources.	
Domain 2: Repres	sentative						
	Metric 2:	Geographic Scope	High	× 1	1	Data are U.S. based.	
	Metric 3:	Applicability	Medium	\times 2	4	EPA-coordinated studies were of synthetic organic chemical manufacturing industry (SOCMI) type facilities, which may include industries within the scopes of the chemicals, but may also include industries outside of the scopes.	
	Metric 4:	Temporal Representativeness	Low	\times 2	6	Underlying data collected through studies from 1980 to 1990. Data more than 20 years old.	
	Metric 5:	Sample Size	Low	× 1	3	Emission factors are presented only as averages; underlying distribution is not characterized.	
Domain 3: Access	sibility/Clar Metric 6:	ity Metadata Completeness	Medium	× 1	2	Metadata includes release media and equipment type that is the source of the release. Does not include the duration over which the emission factors were derived.	
Domain 4: Variab	oility and Ur Metric 7:	ncertainty Metadata Completeness	Low	× 1	3	Does not address variability or uncertainty in average emission factors.	
Overall Quality D	eterminatio	${ m n}^{\dagger}$	Medium		2.2		

^{*} MWF = Metric Weighting Factor † If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	OPW Engineered Systems. 2014. Loading Systems Catalog. OPW Engineered Systems: A Dover Company. ES-LS-6/15-2N Uploaded November 18, 2014.									
Type of Data Source Hero ID	Releases to 5097888	Releases to the Environment; Reports for Data or Information Other than Exposure or Release Data; 5097888								
EXTRACTION										
Parameter			Data							
Life Cycle Stage: Release or Emission					Loading Model umes used to calculate air emissions in model.					
EVALUATION										
Domain		Metric	Rating	MWF*	Score	Comments				
Domain 1: Reliabi	ility									
	Metric 1:	Methodology	High	× 1	1	Data are provided by loading systems vendor; it is expected vendor would provide accurate data on their own loading systems.				
Domain 2: Repres	sentative									
•	Metric 2:	Geographic Scope	High	\times 1	1	Vendor is U.S. based.				
	Metric 3:	Applicability	High	\times 2	2	The loading systems offered in vendor's catalog are applicable for the container types within scope of the model.				
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	Catalog is indicated as copyrighted as of 2015.				
	Metric 5:	Sample Size	High	× 1	1	Vendor's catalog offers loading systems of a variety of sizes, and dimensions are provided for each offered size.				
Domain 3: Access	ibility/Clar	ity								
	Metric 6:	Metadata Completeness	High	× 1	1	All needed metadata for loading system dimensions are provided.				
Domain 4: Variab	ility and U	ncertainty								
	Metric 7:	Metadata Completeness	Medium	× 1	2	Variability among loading systems across vendors is unknown; it is uncertain if the distribution of this vendor's products are capture the distribution across all vendors. However, it is expected that these systems are a reasonable representation of the systems offered in the U.S.				
Overall Quality D	eterminatio	$^{-}$	High		1.1					

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Occupational Exposure

Source Citation: Type of Data Source Hero ID		2016. May 2016 Occupational Enal Exposure; Reports for Data				mates: National Industry-Specific Estimates. Exposure or Release Data;
EXTRACTION Parameter			Data			
Life Cycle Stage: Life Cycle Descrip Number of Sites: Number of Worke	` `	ategory of Use):		_		l to estimate number of sites and workers. l to estimate number of sites and workers.
EVALUATION						
Domain		Metric	Rating	MWF^{\star}	Score	Comments
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	BLS is expected to use reliable survey methods.
Domain 2: Repre	sentative					
Domain 2. Repres	Metric 2:	Geographic Scope	High	\times 1	1	U.S. based economic data
	Metric 3:	Applicability	High	\times 2	2	These economic data cover all industry and occupation types in scope for all chemicals.
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	The BLS OES data are from 2016
	Metric 5:	Sample Size	High	× 1	1	The BLS OES program provides detailed statistics and estimated relative standard error for each state, industry, and occupation survey conducted (https://www.bls.gov/oes/current/oes_research_estimates.htm).
Domain 3: Access	sibility/Clar	itar				
Domain 5. Access	Metric 6:	Metadata Completeness	Medium	\times 1	2	BLS documents results and methods, but underlying survey results not accessible.
Domein 4. Variat	vility and II-	acortainty				
Domain 4: Variab	Metric 7:	Metadata Completeness	Medium	× 1	2	Limited discussion of variability and uncertainty in results.
Overall Quality I	Determinatio	${f n}^\dagger$	High		1.2	

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Type of Data Source		ns Bureau. 2015. Statistics of U nal Exposure; Reports for Data				Exposure or Release Data;		
EXTRACTION Parameter			Data					
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Number of Sites: Number of Workers:				All Used to develop a method to estimate number of sites and workers. Used to develop a method to estimate number of sites and workers.				
EVALUATION Domain		Metric	Rating	MWF*	Score	Comments		
Domain 1: Reliabil	lity Metric 1:	Methodology	High	× 1	1	U.S. Census Bureau is expected to use reliable survey and census methods.		
Domain 2: Represe	entative Metric 2:	Geographic Scope	High	× 1	1	U.S. based economic data		
	Metric 3:	Applicability	High	\times 2	2	These economic data cover all industry and occupation types in scope for all chemicals.		
	Metric 4:	Temporal Representativeness	High	\times 2	2	The Census Bureau SUSB data are from 2015		
	Metric 5:	Sample Size	High	× 1	1	The SUSB is a compilation of data extracted from the Business Register, U.S. Census Bureau's "most complete, current, and consistent data for U.S. business establishments." Incorporates data from economic censuses and current business surveys, quarterly and annual Federal tax records, and other departmental and federal statistics. Expected to be sufficiently representative. (https://www.census.gov/programs-surveys/susb/about.html)		
Domain 3: Accessi	bility/Clari Metric 6:	ity Metadata Completeness	Medium	× 1	2	U.S. Census Bureau documents results and methods, but un-		
						derlying survey results not accessible.		
Domain 4: Variabi	lity and Ur Metric 7:	ncertainty Metadata Completeness	Medium	× 1	2	Limited discussion of variability and uncertainty in results.		
Overall Quality De	eterminatio	${f n}^{\dagger}$	High		1.2			
		Cor	ntinued on 1	next nage	<u> </u>			

- continued from previous page

Source Citation: Type of Data Source Hero ID	U.S. Census Bureau. 2015. Statistics Occupational Exposure; Reports for D 5097881		,	,	or Release Data;	
EVALUATION						
Domain	Metric	Rating	\mathbf{MWF}^{\star}	Score	Comments	

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID		U.S. BLS. 2014. Employee Tenure News Release. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5080421									
EXTRACTION Parameter			Data								
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Exposure Frequency:			All Used to develop estimates of exposure frequency (working days per year and working years)								
EVALUATION Domain		Metric	Rating	MWF*	Score	Comments					
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	BLS is expected to use reliable survey methods.					
Domain 2: Repre											
	Metric 2: Metric 3:	Geographic Scope Applicability	High High	$\begin{array}{c} \times \ 1 \\ \times \ 2 \end{array}$	$\frac{1}{2}$	U.S. based economic data These economic data cover all industry types in scope for all chemicals.					
	Metric 4:	Temporal Representativeness	High	\times 2	2	Median employee tenure with current employer was obtained from the BLS Current Population Survey for January 2014.					
	Metric 5:	Sample Size	High	× 1	1	The Current Population Survey (CPS) is a monthly survey of about 60,000 households. BLS provides detailed statistical treatment of surveys. Expected to be sufficiently representative. (https://www.bls.gov/cps/documentation.htm#reliability)					
Domain 3: Access	sibility/Clar Metric 6:	ity Metadata Completeness	Medium	× 1	2	DIC decomposite and methods but underlying assured					
	MICHIE O.	Metadata Completeness	Mediuii	^ 1		BLS documents results and methods, but underlying survey results not accessible.					
Domain 4: Variab	oility and Un Metric 7:	ncertainty Metadata Completeness	Medium	× 1	2	Limited discussion of variability and uncertainty in results.					
Overall Quality I	Oeterminatio	${f n}^{\dagger}$	High		1.2						

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID		U.S. BLS. 2015. Hours and Employment by Industry Tables - August 6, 2015. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5079873								
EXTRACTION Parameter			Data							
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Exposure Frequency:										
EVALUATION										
Domain		Metric	Rating	MWF*	Score	Comments				
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	BLS is expected to use reliable survey methods.				
Domain 2: Repres	sontativo									
Domain 2. Repres	Metric 2:	Geographic Scope	High	\times 1	1	U.S. based economic data				
	Metric 3:	Applicability	High	\times 2	2	These economic data cover all industry types in scope for all chemicals.				
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	Hours and employment data are from 2016.				
	Metric 5:	Sample Size	High	× 1	1	BLS Labor Productivity and Costs data are used to aid economic policymaking, among other uses, and are expected to be sufficiently representative.				
Domain 3: Access	sibility/Clar	itx								
Domain 9. Access	Metric 6:	Metadata Completeness	Medium	× 1	2	BLS documents results and methods, but underlying survey results not accessible.				
Domain 4: Variab	oility and U	ncertainty								
	Metric 7:	Metadata Completeness	Medium	× 1	2	Limited discussion of variability and uncertainty in results.				
Overall Quality D	Determination	$^{-}$	High	_	1.2					

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID		U.S. Census Bureau. 2019. Survey of Income and Program Participation data. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5080429									
EXTRACTION Parameter			Data								
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Exposure Frequency:			All Used to develop estimates of exposure frequency (working days per year and working years)								
EVALUATION		N. i	D 4:	MIND+	ď						
Domain		Metric	Rating	MWF*	Score	Comments					
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	U.S. Census Bureau is expected to use reliable survey and census methods.					
Domain 2: Repre	contativo										
Domain 2. Repres	Metric 2:	Geographic Scope	High	× 1	1	U.S. based economic data					
	Metric 3:	Applicability	High	\times 2	2	These economic data cover all industry types in scope for all chemicals.					
	Metric 4:	Temporal Representativeness	High	\times 2	2	EPA used the 2008 SIPP Panel Wave 1 (interview months of September through December 2008).					
	Metric 5:	Sample Size	High	× 1	1	The SIPP survey is a continuous series of national panels, with sample size ranging from 14,000 to 52,000 interviewed households. Panels range from 2.5 to 4 years. Expected to be sufficiently representative. (https://www.census.gov/programs-surveys/sipp/about/sipp-introduction-history.html)					
Domain 3: Access	sibility/Clar	ity									
Domain 6. 11000s.	Metric 6:	Metadata Completeness	Medium	× 1	2	BLS documents results and methods, but underlying survey results not accessible.					
Domain 4: Varial	ality and II	ocertainty									
Domain 4. Variat	Metric 7:	Metadata Completeness	Medium	× 1	2	Limited discussion of variability and uncertainty in results.					
Overall Quality I	Determination	$^{-}$	High		1.2						

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Cherrie, JW; Semple, S; Brouwer, D. 2004. Gloves and Dermal Exposure to Chemicals: Proposals for Evaluating Workplace Effectiveness. Annals of Occupational Hygiene. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5080435								
EXTRACTION Parameter			Data						
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure: PPE:				All Used to develop a dermal exposure assessment method for volatile liquids. Provides concepts of glove effectiveness.					
EVALUATION Domain		Metric	Rating	MWF*	Score	Comments			
Domain		WEGITC	Training	IVI VV I	Score	Comments			
Domain 1: Reliab	Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.			
Domain 2: Repres	sentative Metric 2:	Geographic Scope	N/A		N/A	N/A. Geographic scope is not applicable to scientific research of dermal exposures.			
	Metric 3:	Applicability	High	\times 2	2	Article studies effectiveness of gloves in the workplace, which is applicable to the scopes of multiple chemicals.			
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Article was published in 2004; more than 10 but less than 20 years old.			
	Metric 5:	Sample Size	N/A		N/A	N/A. Article presents concepts of dermal exposure and glove effectiveness. Sample size is not applicable.			
Domain 3: Accessibility/Clarity Metric 6: Metadata Completeness		High	× 1	1	Article is well documented with methods, assumptions, and sources.				
Domain 4: Variab	oility and Ur Metric 7:	ncertainty Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.			
Overall Quality I	Determination	<u> </u>	High		1.3				

 $[\]star$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Dancik, Y; Bigliardi, PL; Bigliardi-Qi, Mei. 2015. What happens in the skin? Integrating skin permeation kinetics into studies of developmental and reproductive toxicity following topical exposure. Reproductive Toxicology. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 3223617									
EXTRACTION										
Parameter			Data							
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			All Used to develop a dermal exposure assessment method for volatile liquids.							
EVALUATION										
Domain		Metric	Rating	MWF^{\star}	Score	Comments				
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.				
Domain 2: Repres	contativo									
Bolliam 2. Repres	Metric 2:	Geographic Scope	N/A		N/A	N/A. Geographic scope is not applicable to scientific research of dermal exposures.				
	Metric 3:	Applicability	High	\times 2	2	Article studies skin permeation kinetics, which is applicable to the scopes of multiple chemicals.				
	Metric 4:	Temporal Representativeness	High	\times 2	2	Article was published in 2015; less than 10 years old.				
	Metric 5:	Sample Size	N/A		N/A	$\rm N/A.$ Article studies science of skin permeation and toxicity. Sample size is not applicable.				
Domain 3: Access	sibility/Clar	itv								
Domain of Troops	Metric 6:	Metadata Completeness	High	× 1	1	Article is well documented with methods, assumptions, and sources.				
Domain 4: Variab	oility and U	ncertainty								
Domain 1. Vancon	Metric 7:	Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.				
Overall Quality D	eterminatio	${ m n}^{\dagger}$	High		1.0					

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Frasch, HF; Bunge, AL. 2015. The transient dermal exposure II: post-exposure absorption and evaporation of volatile compounds. Journal of Pharmaceutical Sciences. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 3230538								
EXTRACTION Parameter			Data						
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			All Used to uids.	All Used to develop a dermal exposure assessment method for volatile liq-					
EVALUATION		26	.	2000		· ·			
Domain		Metric	Rating	MWF*	Score	Comments			
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.			
D : 0 D									
Domain 2: Repres	Metric 2:	Geographic Scope	N/A		N/A	N/A. Geographic scope is not applicable to scientific research of dermal exposures.			
	Metric 3:	Applicability	High	\times 2	2	Article studies transient dermal exposure of volatile chemicals that evaporate and absorb into skin simultaneously, which is applicable to the scopes of multiple chemicals.			
	Metric 4:	Temporal Representativeness	High	\times 2	2	Article was published in 2015; less than 10 years old.			
	Metric 5:	Sample Size	N/A		N/A	N/A. Article studies science of skin permeation and evaporation. Sample size is not applicable.			
Domain 3: Access	vibility/Clari	itv							
Domain 6. Meess	Metric 6:	Metadata Completeness	High	× 1	1	Article is well documented with methods, assumptions, and sources.			
Domain 4: Variab	ility and Ur	acortainty							
Domain 4. variab	Metric 7:	Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.			
Overall Quality D	eterminatio	\mathbf{n}^{\dagger}	High		1.0				

 $^{^\}star$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Frasch FH. 2012. Dermal Absorption of Finite doses of Volatile Compounds. Journal of Pharmaceutical Sciences. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5097903								
EXTRACTION Parameter			Data						
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			All Used to uids.	All Used to develop a dermal exposure assessment method for volatile liq-					
EVALUATION									
Domain		Metric	Rating	MWF^{\star}	Score	Comments			
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.			
Domain 2: Repres	sentative								
1	Metric 2:	Geographic Scope	N/A		N/A	N/A. Geographic scope is not applicable to scientific research of dermal exposures.			
	Metric 3:	Applicability	High	\times 2	2	Article studies transient dermal exposure of volatile chemicals that evaporate and absorb into skin simultaneously, which is applicable to the scopes of multiple chemicals.			
	Metric 4:	Temporal Representativeness	High	\times 2	2	Article was published in 2012; less than 10 years old.			
	Metric 5:	Sample Size	N/A		N/A	N/A. Article studies science of skin permeation and evaporation. Sample size is not applicable.			
Domain 3: Access	sibility/Clar	itv							
	Metric 6:	Metadata Completeness	High	× 1	1	Article is well documented with methods, assumptions, and sources.			
Domain 4: Variab	oility and Ur	ncertainty							
	Metric 7:	Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.			
Overall Quality D	eterminatio	${ m n}^{\dagger}$	High		1.0				

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source	and Enviro	Frasch, HF; Dotson, GS; Barbero, AM. 2011. In vitro human epidermal penetration of 1-bromopropane. Journal of Toxicology and Environmental Health, Part A: Current Issues. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data;							
Hero ID	1247930								
EXTRACTION Parameter			Data						
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			All Used to uids.	All Used to develop a dermal exposure assessment method for volatile liq-					
EVALUATION									
Domain		Metric	Rating	MWF^{\star}	Score	Comments			
Domain 1: Reliab	· ·			_	_				
	Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.			
Domain 2: Repres	sentative								
2. 100p10.	Metric 2:	Geographic Scope	N/A		N/A	N/A. Geographic scope is not applicable to scientific research of dermal exposures.			
	Metric 3:	Applicability	High	\times 2	2	Article studies human epidermal penetration of 1-BP, which is applicable to the scope of 1-BP.			
	Metric 4:	Temporal Representativeness	High	\times 2	2	Article was published in 2011; less than 10 years old.			
	Metric 5:	Sample Size	N/A		N/A	$\rm N/A.$ Article studies science of skin permeation and evaporation. Sample size is not applicable.			
Domain 3: Access	sibility/Clar	itv							
	Metric 6:	Metadata Completeness	High	× 1	1	Article is well documented with methods, assumptions, and sources.			
Domain 4: Variab	sility and III	acertainty							
Domain 4. Variab	Metric 7:	Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.			
Overall Quality D	eterminatio	${f n}^{\dagger}$	High		1.0				

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Garrod, AN; Phillips, AM; Pemberton, JA. 2001. Potential exposure of hands inside protective gloves" a summary of data from non-agricultural pesticide surveys. Annals of Occupational Hygiene. Occupational Exposure; Monitoring Data; 5080256								
EXTRACTION Parameter			Data						
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:				All All Used to develop a dermal exposure assessment method for volatile liquids.					
EVALUATION Domain		Metric	Rating	MWF*	Score	Comments			
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.			
Domain 2: Repres	sentative								
_	Metric 2:	Geographic Scope	Medium	\times 1	2	Study measured dermal exposures during activities in the UK. $$			
	Metric 3:	Applicability	High	\times 2	2	Study measured dermal exposures during occupational activities, which is generally relevant to the scopes of multiple chemicals.			
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Article was published in 2001; more than 10 years but less than 20 years old.			
	Metric 5:	Sample Size	High	× 1	1	Statistics of the inside-glove exposures measured are well characterized.			
Domain 3: Access	sibility/Clar	ity							
	Metric 6:	Metadata Completeness	High	× 1	1	Metadata of the measured exposures are well documented.			
Domain 4: Variab	oility and Ui	ncertainty							
	Metric 7:	Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.			
Overall Quality D	eterminatio	${f n}^\dagger$	High		1.3				

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Kasting, BG; Miller, MA. 2006. Kinetics of finite dose absorption through skin 2: Volatile compounds. Journal of Pharmaceutical Sciences. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5018573							
EXTRACTION	5018573							
Parameter			Data					
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			All All Used to ouids.	All Used to develop a dermal exposure assessment method for volatile liq-				
EVALUATION								
Domain		Metric	Rating	MWF*	Score	Comments		
Domain 1: Reliab	oility							
	Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.		
Domain 2: Repres	sentative							
_ 3333332 _ 333F 33	Metric 2:	Geographic Scope	N/A		N/A	N/A. Geographic scope is not applicable to scientific research of dermal exposures.		
	Metric 3:	Applicability	High	\times 2	2	Article studies transient dermal exposure of volatile chemicals that evaporate and absorb into skin simultaneously, which is applicable to the scopes of multiple chemicals.		
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Article was published in 2006; more than 10 years but less than 20 years old.		
	Metric 5:	Sample Size	N/A		N/A	$\rm N/A.$ Article studies science of skin permeation and evaporation. Sample size is not applicable.		
Domain 3: Access	sibility/Clar	itz						
Domain 5. Access	Metric 6:	Metadata Completeness	High	× 1	1	Article is well documented with methods, assumptions, and sources.		
Domain 4: Variab	oility and U	ncertainty						
	Metric 7:	Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.		
Overall Quality D	Determinatio	\mathbf{n}^{\dagger}	High		1.3			

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Marquart, H; Franken, R; Goede, H; Fransman, W; Schinkel, J. 2017. Validation of the dermal exposure model in ECETOC TRA. Annals of Work Exposures and Health. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5080455								
EXTRACTION Parameter			Data						
Life Cycle Stage: Life Cycle Descrip Route of Exposure		ategory of Use):	All All Used to cuids.	levelop a	dermal	exposure assessment method for volatile liq-			
EVALUATION									
Domain		Metric	Rating	MWF*	Score	Comments			
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.			
Domain 2: Repres	sentative Metric 2:	Geographic Scope	N/A		N/A	N/A. Geographic scope not applicable to the validation of the ECETOC TRA model.			
	Metric 3:	Applicability	High	\times 2	2	ECETOC TRA model and exposure studies used for validation cover a variety of occupational scenarios, which are applicable to the scopes of multiple chemicals.			
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	Article was published in 2017; less than 10 years old.			
	Metric 5:	Sample Size	Medium	× 1	2	Statistics of dermal exposure observations obtained from the literature are not fully characterized.			
Domain 3: Access	ibility/Clar Metric 6:	ity Metadata Completeness	Medium	× 1	2	Article is well documented with methods, assumptions, and results; however, sources used from literature search are not fully described and the metadata associated with the literature review exposure studies are not provided.			
Domain 4: Variab	ility and Ur Metric 7:	ncertainty Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.			
Overall Quality D	eterminatio	\mathbf{n}^{\dagger}	High		1.3				

 $[\]label{eq:mwf} ^\star \, \mathrm{MWF} = \mathrm{Metric} \, \, \mathrm{Weighting} \, \, \mathrm{Factor} \\ ^\dagger \, \mathrm{If} \, \, \mathrm{any} \, \, \mathrm{individual} \, \, \mathrm{metrics} \, \, \mathrm{are} \, \, \mathrm{deemed} \, \, \mathrm{Unacceptable}, \, \mathrm{then} \, \, \mathrm{the} \, \, \mathrm{overall} \, \, \mathrm{rating} \, \, \mathrm{is} \, \, \mathrm{also} \, \, \mathrm{unacceptable}. \, \, \mathrm{Otherwise}, \, \mathrm{the} \, \, \mathrm{overall} \, \, \mathrm{rating} \, \, \mathrm{is} \, \, \mathrm{based} \, \, \mathrm{on} \, \, \mathrm{the} \, \, \mathrm{following} \, \, \mathrm{scale} :$ High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID		Baldwin, PE; Maynard, AD. 1998. A Survey of Wind Speed in Indoor Workplaces. Annals of Occupational Hygiene. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 3045135									
EXTRACTION Parameter			Data								
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			All Used to ouids.	All Used to develop a dermal exposure assessment method for volatile liq-							
EVALUATION											
Domain		Metric	Rating	MWF^{\star}	Score	Comments					
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.					
Domain 2: Repres		VI VIII VOI									
Domain 2. Repres	Metric 2:	Geographic Scope	Medium	× 1	2	Article studied wind speeds in indoor workplaces in the UK.					
	Metric 3:	Applicability	High	\times 2	2	The types of workplaces studied include workplaces applicable to the scopes of multiple chemicals.					
	Metric 4:	Temporal Representativeness	Medium	$\times 2$	4	Article was published in 1998; more than 10 years but less than 20 years old.					
	Metric 5:	Sample Size	High	\times 1	1	Statistics of wind speed surveys are well characterized.					
Domain 3: Access	sibility/Clar	ity									
	Metric 6:	Metadata Completeness	High	× 1	1	Article is well documented with methods, assumptions, results, and sources.					
Domain 4: Variab	oility and U	ncertainty									
	Metric 7:	Metadata Completeness	High	× 1	1	Detailed discussion on variability/uncertainty.					
Overall Quality Determination [†]			High		1.3						

^{*} MWF = Metric Weighting Factor † If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:		17. Chemical Exposure Health I	Data (CEH)	D) provid	led by (OSHA to EPA.
Type of Data Source		nal Exposure; Monitoring Data;				
Hero ID	3827305					
EXTRACTION						
Parameter			Data			
Life Cycle Stage:			All			
Life Cycle Descri		_ ,	All			
Exposure Concen	tration (Uni	it):	Provides	personal	breathi:	ng zone and area monitoring data.
EVALUATION						
Domain		Metric	Rating	\mathbf{MWF}^{\star}	Score	Comments
D D !! 1	•1• .					
Domain 1: Reliab		26.1.1.1	TT: 1			
	Metric 1:	Methodology	High	$\times 1$	1	OSHA and state inspectors are expected to use OSHA or NIOSH sampling methods. Samples sent to the OSHA SLTC
						are expected to be analyzed using OSHA or NIOSH analytical
						methods.
Domain 2: Repre	contativo					
Domain 2. Repre	Metric 2:	Geographic Scope	High	× 1	1	U.S. based exposure data
	Metric 3:	Applicability	Medium	$\stackrel{\wedge}{\times} \stackrel{1}{2}$	4	The OSHA data include occupational scenarios within the
	wienie o.	ripplicability	Wediam	A 2	1	scopes of the chemicals as identified by NAICS code and facility
						name. However, some occupational scenarios are not clear and
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	cannot be clearly mapped to conditions of use within scope. Data provided by OSHA are not more than 10 years old.
	Metric 5:	Sample Size	High	$\times 2 \times 1$	1	Individual measurements are provided so the sample sets can
	Metric 5.	Sample Size	IIIgii	^ 1	1	be fully statistically characterized.
Domain 3: Access	٠,	•	3.6 1:	1	0	
	Metric 6:	Metadata Completeness	Medium	$\times 1$	2	OSHA data include sample type and exposure type. Sample times also provided. Exposure frequency is inconsistently pro-
						vided. Worker job descriptions provided, but often lacks suffi-
						cient clarity.
Domain 4: Variability and Uncertainty						
Domain 4. variai	Metric 7:	•	Low	× 1	3	OCIIA data da not discuss reviability en un containty
	Menic 7.	Metadata Completeness	ьом	^ I	ა 	OSHA data do not discuss variability or uncertainty.
Organall Organitas I) at amma im a t ! -	m†	II: odo		1 6	
Overall Quality I	peterminatio	on ·	High		1.6	

^{*} MWF = Metric Weighting Factor † If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

	Defense Occupational and Environmental Health Readiness System - Industrial Hygiene (DOEHRS-IH). 2018. Email between DOD and EPA: RE: [Non-DoD Source] Update: DoD exposure data for EPA risk evaluation - EPA request for additional information. Occupational Exposure; Monitoring Data; 5178607							
EXTRACTION Parameter			Data					
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Exposure Concentration (Unit):			All All Provides					
EVALUATION Domain		Matria	Datina	MWF*	Caona	Comments		
Domain		Metric	Rating	M W F ^	Score	Comments		
Domain 1: Reliabil	ity Metric 1:	Methodology	High	× 1	1	DOD service branches use OSHA and NIOSH methods and DOD methods, which are expected to be equivalent to OSHA or NIOSH methods.		
Domain 2: Represe	entative							
	Metric 2:	Geographic Scope	High	\times 1	1	U.S. based exposure data		
	Metric 3:	Applicability	Medium	× 2	4	The DOD data include occupational conditions of use within the scopes of the chemicals, although additional uses poten- tially outside of scope may also be included. However, some occupational scenarios are not clear and cannot be clearly mapped to conditions of use within scope.		
	Metric 4:	Temporal Representativeness	High	\times 2	2	Approximately 82 percent of the samples provided by DOD are not more than 10 years old.		
	Metric 5:	Sample Size	High	× 1	1	Individual measurements are provided so the sample sets car be fully statistically characterized.		
Domain 3: Accessib	bility/Clari	itv						
		Metadata Completeness	Medium	× 1	2	DOD data include sample type (PBZ), sample time, process duration and frequency, and workshift duration. Process and worker job descriptions are provided, but inconsistent in detail and often lack sufficient clarity.		
Domain 4: Variabil	lity and Ur	ncertainty						
	Metric 7:	Metadata Completeness	Low	× 1	3	DOD data do not discuss variability or uncertainty.		
		Con	tinued on 1	next page	9			

- continued from previous page

	Contr	naca nom j	or c v rous	page			
Source Citation:	Defense Occupational and Environmental Health Readiness System - Industrial Hygiene (DOEHRS-IH). 2018. Email between DOD and EPA: RE: [Non-DoD Source] Update: DoD exposure data for EPA risk evaluation - EPA request for additional information.						
Type of Data Source	Occupational Exposure; Monitoring Dat	a;					
Hero ID	5178607	,					
EVALUATION							
Domain	Metric	Rating	MWF^{\star}	Score	Comments		
Overall Quality I	Oetermination [†]	High		1.6			

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

of chlorin Type of Data Source Occupati	California Air Resources Board. 2000. Initial statement of reasons for the proposed airborne toxic control measure for emissions of chlorinated toxic air contaminants from automotive maintenance and repair activities. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data;							
Hero ID 5071458								
EXTRACTION Parameter		Data						
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			Brake Servicing Model Brake Servicing Model Used to develop an inhalation exposure model.					
EVALUATION								
Domain	Metric	Rating	\mathbf{MWF}^{\star}	Score	Comments			
Domain 1: Reliability Metric 1	Methodology	High	× 1	1	CARB is expected to use reliable data collection and survey methods.			
Domain 2: Representative								
Metric 2	Geographic Scope	High	$\times 1$	1	Data surveyed and collected from U.S. (California) facilities			
Metric 3	Applicability	High	\times 2	2	The CARB data are specific to brake servicing and include halogenated solvent aerosol brake cleaners, which is applicable to the scope of the model.			
Metric 4	Temporal Representativeness	Medium	\times 2	4	The report was published in 2000, the manufacturer and facility surveys were conducted in 1997 and 1998, and site visits were conducted circa 1998. All less than 20 years old (from 2016).			
Metric 5	Sample Size	Medium	× 1	2	Some data elements from site visits include all individual data points; some surveyed data elements include some statistics (more than range but not full distribution), and some data elements have limited distribution information.			
Domain 3: Accessibility/Cla	oritz.							
Metric 6		High	× 1	1	Report fully documents its data sources, assessment methods, results, and assumptions.			
Domain 4: Variability and V	Incortainty							
Metric 7	· ·	High	× 1	1	Report discusses and addresses variability and uncertainty.			
Overall Quality Determination [†]		High		1.3				
		ntinued on 1						

- continued from previous page

		maea nom previous pag	·6°
Source Citation:	California Air Resources Board. 2000. of chlorinated toxic air contaminants		or the proposed airborne toxic control measure for emission e and repair activities.
Type of Data Source Hero ID	Occupational Exposure; Reports for I 5071458	a or Information Other tha	an Exposure or Release Data;
EVALUATION			
Domain	Metric	Rating MWF* Scor	ore Comments

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source	Demou, E.,Hellweg, S.,Wilson, M. P.,Hammond, S. K.,McKone, T. E 2009. Evaluating indoor exposure modeling alternatives for LCA: A case study in the vehicle repair industry. Environmental Science and Technology. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data;							
Hero ID	2591566							
EXTRACTION								
Parameter			Data					
Life Cycle Stage:			Brake Sei	rvicing M	odel			
Life Cycle Descrip	tion (Subca	ategory of Use):	Brake Sei					
Route of Exposure	e:		Used to d	levelop a	n inhala	tion exposure model.		
EVALUATION								
Domain		Metric	Rating	MWF^{\star}	Score	Comments		
Domain 1: Reliab	ility							
	Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.		
Domain 2: Repres	sentative							
•	Metric 2:	Geographic Scope	Medium	× 1	2	Air ventilation rate data are at least in part based on European data (but may also include U.S. data).		
	Metric 3:	Applicability	High	$\times 2$	2	Ventilation rate data are applicable to the scope of the model.		
	Metric 4:	Temporal Representativeness	Low	\times 2	6	Paper published in 2009; data are based on 2006 and 1991 data. Data are in part more than than 20 years old (as measured from 2016).		
	Metric 5:	Sample Size	Medium	× 1	2	Ventilation rate provided as range with uncertain distribution.		
Domain 3: Access	ibility/Clar	itx						
Domain 5. Access	Metric 6:	Metadata Completeness	Medium	\times 1	2	Sources are cited, but does not provide details on how reported values were derived from cited sources.		
Domain 4: Variab	ility and II.	ncartainty						
Domain 4. variab	Metric 7:	Metadata Completeness	Medium	× 1	2	Variability of ventilation rates provided, but uncertainty not discussed.		
Overall Quality Determination [†]			Medium		1.9			

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Golsteijn, L., Huizer, D., Hauck, M., van Zelm, R., Huijbregts, M. A 2014. Including exposure variability in the life cycle impact assessment of indoor chemical emissions: the case of metal degreasing. Environment International. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 2537636								
EXTRACTION									
Parameter			Data						
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			Brake Servicing Model Brake Servicing Model Used to develop an inhalation exposure model.						
EVALUATION									
Domain		Metric	Rating	MWF^{\star}	Score	Comments			
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.			
Domain 2: Repres	contativo								
Domain 2. Repres	Metric 2:	Geographic Scope	Medium	× 1	2	Air ventilation rate data based on European data.			
	Metric 3:	Applicability	High	$\times 2$	$\stackrel{-}{2}$	Ventilation rate data are applicable to the scope of the model.			
	Metric 4:	Temporal Representativeness	High	\times 2	2	Ventilation rate data based on 2012 and 2003 sources. Article published in 2014.			
	Metric 5:	Sample Size	Medium	\times 1	2	Ventilation rate provided as range with uncertain distribution.			
Domain 3: Access	sibility/Clar Metric 6:	rity Metadata Completeness	Medium	× 1	2	Sources are cited, but does not provide details on how reported values were derived from cited sources.			
Domain 4: Variab	oility and Un Metric 7:	ncertainty Metadata Completeness	Medium	× 1	2	Variability of ventilation rates provided, but uncertainty not discussed.			
Overall Quality Determination [†]			High		1.4				

^{*} MWF = Metric Weighting Factor † If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Hellweg, S; Demou, E; Bruzzi, R; Meijer, A; Rosenbaum, RK; Huijbregts, MA; Mckone, TE. 2009. Integrating human indoor air pollutant exposure within Life Cycle Impact Assessment. Environmental Science and Technology. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 634560							
EXTRACTION Parameter			Data					
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			Brake Servicing Model Brake Servicing Model Used to develop an inhalation exposure model.					
EVALUATION								
Domain		Metric	Rating	\mathbf{MWF}^{\star}	Score	Comments		
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.		
Di. 9. D	4-4:							
Domain 2: Repres	Metric 2:	Geographic Scope	Medium	× 1	2	Air ventilation rate data are at least in part based on European data (but may also include U.S. data).		
	Metric 3:	Applicability	High	$\times 2$	2	Ventilation rate data are applicable to the scope of the model.		
	Metric 4:	Temporal Representativeness	Low	\times 2	6	Paper published in 2009; data appear to be from sources dating from 1989 to 1993.		
	Metric 5:	Sample Size	Medium	\times 1	2	Ventilation rate provided as range with uncertain distribution.		
Domain 3: Access	sibility/Clar Metric 6:	ity Metadata Completeness	Medium	× 1	2	Sources are cited, but does not provide details on how reported values were derived from cited sources.		
Domain 4: Variab	oility and Ur Metric 7:	ncertainty Metadata Completeness	Medium	× 1	2	Variability of ventilation rates provided, but uncertainty not		
Overall Quality Determination [†]			Medium		1.9	discussed.		

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	Scientific Consulting Group, Inc 2013. Final peer review comments for the OPPT trichloroethylene (TCE) draft risk assessment.									
Type of Data Source Hero ID	Occupation 3044932	Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 3044932								
EXTRACTION										
Parameter			Data							
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Route of Exposure:			Brake Ser	Brake Servicing Model Brake Servicing Model Used to develop an inhalation exposure model.						
EVALUATION										
Domain		Metric	Rating	\mathbf{MWF}^{\star}	Score	Comments				
Domain 1: Reliab	ility Metric 1:	Methodology	Low	× 1	3	Peer reviewer does not provide data sources or techniques used to arrive at ventilation rate estimates.				
Domain 2: Repres				_						
	Metric 2:	Geographic Scope	High	$\times 1$	1	Peer reviewer's experience appears to be U.S. based.				
	Metric 3:	Applicability	High	$\times 2$	2	Ventilation rate data are applicable to the scope of the model.				
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Time period of the peer reviewer's observations not provided, but not expected to be outdated.				
	Metric 5:	Sample Size	Medium	× 1	2	Ventilation rate provided as range with uncertain distribution.				
Domain 3: Access	ibility /Clan	;+								
Domain 5. Access	Metric 6:	Metadata Completeness	Low	\times 1	3	Underlying data sources not transparent.				
Domain 4: Variab	Metric 7:	ncertainty Metadata Completeness	Medium	× 1	2	Variability of ventilation rates provided, but uncertainty not discussed.				
Overall Quality Determination [†]			Medium		1.9					

^{*} MWF = Metric Weighting Factor † If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	Blando, J. D., Schill, D. P., De La Cruz, M. P., Zhang, L., Zhang, J 2010. Preliminary study of propyl bromide exposure among New Jersey dry cleaners as a result of a pending ban on perchloroethylene. Journal of the Air and Waste Management Association.									
Type of Data Source Hero ID	Occupation 1619253	Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 1619253								
EXTRACTION Parameter			Data							
Life Cycle Stage: Route of Exposure:			1) Dry Cleaning Release Model2) Dry Cleaning Exposure Model Provides data used in inhalation exposure and release models (number of loads per day).							
EVALUATION										
Domain		Metric	Rating	MWF^{\star}	Score	Comments				
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1					
	Metric 1:	wiethodology	Trigii	X 1	1	Article is published in peer-reviewed scientific journal.				
Domain 2: Repres			TT: 1							
	Metric 2: Metric 3:	Geographic Scope Applicability	High High	$\begin{array}{c} \times \ 1 \\ \times \ 2 \end{array}$	$\frac{1}{2}$	Studies New Jersey (U.S.) based dry cleaners. Observed dry cleaning data are applicable to the scope of the				
		· ·	IIIgii		2	model.				
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	Paper published in 2010, site visits conducted circa 2009; less than 10 years old.				
	Metric 5:	Sample Size	High	× 1	1	Limited number of samples, but individual data points allow characterization of distribution.				
Domain 3: Access	sibility/Clar	itv								
	Metric 6:	Metadata Completeness	High	× 1	1	Article fully documents its data sources, assessment methods, results, and assumptions.				
Domain 4: Variab	sility and IIs	acortainty								
Domain 4. Variat	Metric 7:	Metadata Completeness	High	× 1	1	Discusses uncertainty and variability in observed data.				
Overall Quality Determination [†]			High		1.0					

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Division, I	Air Resources Board. 2006. C Emissions Assessment Branch. nal Exposure; Reports for Data			J	istry Technical Assessment Report. Stationary Source Exposure or Release Data;		
EXTRACTION								
Parameter			Data					
Life Cycle Stage: Route of Exposure:				1) Dry Cleaning Release Model2) Dry Cleaning Exposure Model3) Spot Cleaning Exposure Model Provides data used in inhalation exposure and release models.				
EVALUATION								
Domain		Metric	Rating	MWF^{\star}	Score	Comments		
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	CARB is expected to use reliable data collection and survey methods.		
Domain 2: Repres	sontativo							
Domain 2. Repres	Metric 2:	Geographic Scope	High	$\times 1$	1	Data surveyed and collected from U.S. (California) facilities		
	Metric 3:	Applicability	High	\times 2	2	Observed dry cleaning data are applicable to the scope of the model.		
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Report published in 2006, data surveyed and collected circa 2003; more than 10 years old but less than 20 years.		
	Metric 5:	Sample Size	High	\times 1	1	Collected data are generally provided with robust statistics.		
Domain 3: Access	sibility/Clar	itv						
Domain 9. Necesi	Metric 6:	Metadata Completeness	High	× 1	1	Article fully documents its data sources, assessment methods, results, and assumptions.		
Domain 4: Variab	oility and U	ncertainty						
Domain 1. Variat	Metric 7:	Metadata Completeness	High	× 1	1	Discusses uncertainty and variability in observed data.		
Overall Quality D)eterminatio	${ m n}^{\dagger}$	High		1.2			

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID		97. Hazard control: Control of e nal Exposure; Reports for Data				e in commercial drycleaning (machine design) (HC 18). Exposure or Release Data;
EXTRACTION Parameter			Data			
Life Cycle Stage: Route of Exposure:				_		Model2) Dry Cleaning Exposure Model alation exposure and release models.
EVALUATION						
Domain		Metric	Rating	MWF*	Score	Comments
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	NIOSH is expected to use reliable data collection and survey methods.
Domain 2: Repres	sentative					
	Metric 2:	Geographic Scope	High	$\times 1$	1	Data are based on U.S. dry cleaning machines.
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Document published in 1997; more than 10 years old but generally less than 20 years.
	Metric 5:	Sample Size	Low	× 1	3	Data characterized with no statistics.
Domain 3: Access	ibility/Clar Metric 6:	ity Metadata Completeness	Low	× 1	3	Provides results but generally does not provide data sources, methods, or assumptions.
Domain 4: Variab	ility and Uı Metric 7:	ncertainty Metadata Completeness	Low	× 1	3	Generally does not discuss variability or uncertainty.
Overall Quality D	Overall Quality Determination [†]		Medium		1.9	

 $^{^\}star$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	risk factor	s.			v	eaners: chemical exposures, fire hazards, and ergonomic
Type of Data Source Hero ID	Occupation 3044963	nal Exposure; Reports for Data	or Informa	tion Othe	er than	Exposure or Release Data;
EXTRACTION						
Parameter			Data			
Life Cycle Stage: Route of Exposur				Model2) Dry Cleaning Exposure Model alation exposure and release models.		
EVALUATION						
Domain		Metric	Rating	MWF^{\star}	Score	Comments
Domain 1: Reliab	oility					
	Metric 1:	Methodology	High	× 1	1	NIOSH is expected to use reliable data collection and survey methods.
Domain 2: Repres	sentative					
	Metric 2:	Geographic Scope	High	$\times 1$	1	Data are based on U.S. dry cleaning machines.
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.
	Metric 4:	Temporal Representativeness	Medium	$\times 2$	4	Document published in 1997; more than 10 years old but generally less than 20 years.
	Metric 5:	Sample Size	Low	× 1	3	Data characterized with no statistics.
Domain 3: Access	zibility/Clar	ity				
Domain 5. Access	Metric 6:	Metadata Completeness	Low	× 1	3	Provides results but generally does not provide data sources, methods, or assumptions.
Domain 4. V:-1	:1:4:. and II-	a containte				
Domain 4: Variab	Metric 7:	Metadata Completeness	Low	× 1	3	Generally does not discuss variability or uncertainty.
Overall Quality E	Determinatio	${f n}^{\dagger}$	Medium		1.9	

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	use in four	Eisenberg, J.,Ramsey, J 2010. Health hazard evaluation report no. HETA 2008-0175-3111, Evaluation of 1-Bromopropane use in four New Jersey commercial dry cleaning facilities. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 3970603									
EXTRACTION Parameter			Data								
Life Cycle Stage: Route of Exposure:				Dry Cleaning Release Model2) Dry Cleaning Exposure Model Provides data used in inhalation exposure and release models.							
EVALUATION											
Domain		Metric	Rating	MWF^{\star}	Score	Comments					
Domain 1: Reliab	ility										
	Metric 1:	Methodology	High	× 1	1	NIOSH is expected to use reliable data collection methods.					
Domain 2: Repres	sentative										
	Metric 2:	Geographic Scope	High	$\times 1$	1	Data are based on U.S. dry cleaning machines.					
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.					
	Metric 4:	Temporal Representativeness	High	\times 2	2	Site visits conducted in 2008; less than 10 years old (from 2016).					
	Metric 5:	Sample Size	High	× 1	1	Limited number of samples, but individual data points allow characterization of distribution.					
Domain 3: Access	sibility/Clar	ity									
Domaii 5. Meess	Metric 6:	Metadata Completeness	High	× 1	1	Article fully documents its data sources, assessment methods, results, and assumptions.					
Domain 4: Variab	ility and U	ncertainty									
Domain 1. Valido	Metric 7:	Metadata Completeness	Medium	× 1	2	Discusses variability among the different sites.					
Overall Quality D	Overall Quality Determination †				1.1						

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Hazardous Waste Management Program in King County. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 3827371									
EXTRACTION										
Parameter			Data							
Life Cycle Stage: Route of Exposure:				1) Dry Cleaning Release Model2) Dry Cleaning Exposure Model3) Spot Cleaning Exposure Model Provides data used in inhalation exposure and release models.						
EVALUATION										
Domain		Metric	Rating	MWF^{\star}	Score	Comments				
Domain 1: Reliabi	ility Metric 1:	Methodology	High	× 1	1	King County has used reliable data collection and survey methods.				
Domain 2: Repres	ontativo									
Domain 2: Repres	Metric 2:	Geographic Scope	High	× 1	1	Data are based on U.S. dry cleaning machines (Washington).				
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.				
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	Site visits conducted in 2009-2010, with surveys conducted afterwards; less than 10 years old (from 2016).				
	Metric 5:	Sample Size	High	\times 1	1	Collected data are generally provided with robust statistics.				
Domain 3: Access	ibility/Clar Metric 6:	ity Metadata Completeness	High	× 1	1	Article fully documents its data sources, assessment methods, results, and assumptions.				
Domain 4: Variab	ility and Uı Metric 7:	ncertainty Metadata Completeness	High	× 1	1	Discusses uncertainty and variability in observed data.				
Overall Quality De	Overall Quality Determination [†]				1.0					

^{*} MWF = Metric Weighting Factor † If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	trichloroet	for Research and Technical A chylene in the textile cleaning in	dustry.	2007.	•	g chemicals: Alternatives to perchloroethylene and
Type of Data Source Hero ID	3045700	nal Exposure; Reports for Data	or Informa	ition Othe	er than	Exposure or Release Data;
EXTRACTION						
Parameter			Data			
Life Cycle Stage: Route of Exposure:						re Model2) Dry Cleaning Exposure Model alation exposure models.
EVALUATION						
Domain		Metric	Rating	MWF^{\star}	Score	Comments
Domain 1: Reliab	ility					
Domain 1. Teenab	Metric 1:	Methodology	High	× 1	1	CalEPA and EPA funded project expected to use reliable data collection methods.
Domain 2: Repres	sentative					
•	Metric 2:	Geographic Scope	High	$\times 1$	1	Data are based on U.S. dry cleaning machines (California)
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	Report published in 2007; less than 10 years old (from 2016).
	Metric 5:	Sample Size	Low	× 1	3	Data characterized with no statistics.
Domain 3: Access	ibility/Clar	itv				
	Metric 6:	Metadata Completeness	Low	× 1	3	Provides results but generally does not provide data sources, methods, or assumptions.
Domain 4: Variab	ility and Ur	ncertainty				
Domain 1. Variab	Metric 7:	· ·	Low	× 1	3	Generally does not discuss variability or uncertainty.
Overall Quality D	Overall Quality Determination [†]		Medium		1.7	

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	von Grote, J., Hürlimann, C., Scheringer, M., Hungerbühler, K 2006. Assessing occupational exposure to perchloroethylene in dry cleaning. Journal of Occupational and Environmental Hygiene. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 632592									
EXTRACTION	002002									
Parameter			Data							
Life Cycle Stage: Route of Exposure:			/ -	_		e Model2) Dry Cleaning Exposure Model alation exposure models.				
EVALUATION										
Domain		Metric	Rating	MWF^{\star}	Score	Comments				
Domain 1: Reliab	ilitz									
Domain 1. Itenab	Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.				
Domain 2: Repres	sentative									
Bolliam 2. Ropros	Metric 2:	Geographic Scope	Medium	\times 1	2	Data based on German dry cleaners (OECD country).				
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.				
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Work based on dissertation published in 2003. More than 10 but less than 20 years old.				
	Metric 5:	Sample Size	Medium	× 1	2	The various data elements used from the study are presented mostly as ranges or averages.				
Domain 3: Access	ibility/Clar	itv								
Domain 6. Treeess	Metric 6:	Metadata Completeness	Medium	× 1	2	Sources are cited, but does not provide details on how reported values were derived from cited sources.				
Domain 4: Variab	ility and Ur	ocertainty								
Domain 4. Variau	Metric 7:	Metadata Completeness	Medium	× 1	2	Variability in parameter values discussed, but no discussion of uncertainties.				
Overall Quality D	eterminatio	${f n}^\dagger$	Medium		1.7					

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Morris, M; Wolf, K. 2005. Evaluation of New and Emerging Technologies for Textile Cleaning. Institute for Research and Technical Assistance. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5176441								
EXTRACTION Parameter			Data						
Life Cycle Stage: Route of Exposure:				_	-	re Model2) Dry Cleaning Exposure Model alation exposure models.			
EVALUATION									
Domain		Metric	Rating	MWF^{\star}	Score	Comments			
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	CARB, CalEPA, and EPA funded project expected to use reliable data collection methods.			
Domain 2: Repres	sentative								
Bolliam 2. Tepro.	Metric 2:	Geographic Scope	High	\times 1	1	Data are based on U.S. dry cleaners (California).			
	Metric 3:	Applicability	High	$\times 2$	2	Spot cleaning data are applicable to the scope of the model.			
	Metric 4:	Temporal Representativeness	Medium	$\times 2$	4	Report published in 2005; more than 10 but less than 20 years old.			
	Metric 5:	Sample Size	Low	× 1	3	Data characterized with no statistics.			
Domain 3: Access	sibility/Clar	ritv							
	Metric 6:	Metadata Completeness	Low	× 1	3	Provides results but generally does not provide data sources, methods, or assumptions.			
Domain 4: Variab	oility and U	ncertainty							
	Metric 7:	Metadata Completeness	Low	× 1	3	Generally does not discuss variability or uncertainty.			
Overall Quality D	Oeterminatio	${ m nn}^\dagger$	Medium		1.9				

 $[\]star$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	exposures	with alternative "green" solvent	•	0,		ERC ban among dry cleaners leads to 1-bromopropane
Type of Data Source Hero ID	Occupation 3045119	nal Exposure; Reports for Data	or Informa	tion Othe	er than	Exposure or Release Data;
EXTRACTION Parameter			Data			
Life Cycle Stage: Route of Exposure	Dry Cleaning Exposure Model Provides data to estimate 1-BP based spot cleaner use rate in inhalation exposure model.					
EVALUATION						
Domain		Metric	Rating	MWF^{\star}	Score	Comments
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	State and academic research expected to use reliable data col-
	WICUIC 1.	Wichiodology	111611	^ I		lection methods.
Domain 2: Repres	sentative					
2011ani 21 100pros	Metric 2:	Geographic Scope	High	$\times 1$	1	Data are based on U.S. dry cleaners (New Jersey).
	Metric 3:	Applicability	High	$\times 2$	2	Spot cleaning data are applicable to the scope of the model.
	Metric 4:	Temporal Representativeness	High	\times 2	2	Data collected from 2008 to 2009; less than 10 years old (from 2016).
	Metric 5:	Sample Size	Low	\times 1	3	Data characterized with no statistics.
Domain 3: Access	ibility/Clar	ity				
Domain 6. Access	Metric 6:	Metadata Completeness	Medium	× 1	2	Sources are cited, but does not provide details on how reported values were derived from cited sources.
Domoin 4. Vanial	ilitre and II-	a containty				
Domain 4: Variab	Metric 7:	Metadata Completeness	Medium	× 1	2	Limited discussion of variability and uncertainty in results.
Overall Quality Determination †		High		1.4		

^{*} MWF = Metric Weighting Factor † If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID		ch International. 2013. Drysolv : nal Exposure; Reports for Data				· ·		
EXTRACTION Parameter			Data					
Life Cycle Stage: Route of Exposure:			1) Spot Cleaning Exposure Model2) Dry Cleaning Exposure Model Provides 1-BP concentration in 1-BP based spot cleaner.					
EVALUATION								
Domain		Metric	Rating	MWF*	Score	Comments		
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	Product manufacturer is expected to know the composition of their products.		
Domain 2: Repres	sentative							
Bomain 2. Teepres	Metric 2:	Geographic Scope	High	\times 1	1	Product available for sale in U.S.		
	Metric 3:	Applicability	High	$\times 2$	2	Spot cleaning data are applicable to the scope of the model.		
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	SDS issue date is 2013; less than 10 years old.		
	Metric 5:	Sample Size	Low	× 1	3	Data characterized with no statistics.		
Domain 3: Access	sibility/Clar	ity						
	Metric 6:	Metadata Completeness	High	\times 1	1	All needed metadata are provided.		
Domain 4: Variab	oility and Un Metric 7:	ncertainty Metadata Completeness	Medium	× 1	2	Range in concentration provided; unclear if this represents variability or uncertainty in the concentration.		
Overall Quality D	Overall Quality Determination [†]				1.3			

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	Docket, St	abject: Background information	document].		Eastern Research Group Inc 2005. [Letter from Eric Goehl and Jennifer O'Neil, Eastern Research group, Inc, to Dry Cleaning Docket, Subject: Background information document]. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data;								
Type of Data Source Hero ID	Occupation 3045690	nal Exposure; Reports for Data	or Informat	tion Othe	er than	Exposure or Release Data;							
EXTRACTION													
Parameter			Data										
Life Cycle Stage:			Dry Clea	ning Exp	osure M	Iodel							
Route of Exposure:			Provides	data used	d in inh	alation exposure models.							
EVALUATION													
Domain		Metric	Rating	MWF^{\star}	Score	Comments							
Domain 1: Reliab	ility												
	Metric 1:	Methodology	High	× 1	1	Data collected in support of EPA rulemaking.							
Domain 2: Repres	sentative												
•	Metric 2:	Geographic Scope	High	\times 1	1	Data are based on U.S. dry cleaners.							
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.							
	Metric 4:	Temporal Representativeness	Medium	$\times 2$	4	Data are more than 10 years old but less than 20 years old.							
	Metric 5:	Sample Size	High	× 1	1	Individual data points provided.							
Domain 3: Access	sibility/Clar	ity											
	Metric 6:	Metadata Completeness	High	× 1	1	Report fully documents its data sources, assessment methods, results, and assumptions. $$							
Domain 4: Variab	ility and U	ncertainty											
Domain 4. Variat	Metric 7:	Metadata Completeness	Medium	× 1	2	Limited discussion of variability and uncertainty in results.							
Overall Quality D	Overall Quality Determination [†]				1.3								

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	Massachus worksheet.	assachusetts Department of Environmental Protection. 2013. Alternative dry cleaning technologies comparative analysis orksheet. ccupational Exposure; Reports for Data or Information Other than Exposure or Release Data;								
Type of Data Source Hero ID	Occupatio 3045045	nal Exposure; Reports for Data	or Informa	tion Othe	er than	Exposure or Release Data;				
EXTRACTION Parameter			Data							
Life Cycle Stage:					re Model2) Dry Cleaning Exposure Model					
Route of Exposur	Route of Exposure:				-BP ba	sed spot cleaner for use in inhalation exposure				
EVALUATION										
Domain		Metric	Rating	MWF*	Score	Comments				
Domain 1: Reliab	ilitz									
Domaii 1. Renad	Metric 1:	Methodology	High	× 1	1	State and TURI expected to use reliable data collection methods.				
Domain 2: Repres	sentative									
Bollium 2. Idopio	Metric 2:	Geographic Scope	High	$\times 1$	1	Data are based on U.S. dry cleaners.				
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.				
	Metric 4:	Temporal Representativeness	High	$\times 2$	2	Worksheet published in 2013; less than 10 years old.				
	Metric 5:	Sample Size	Low	\times 1	3	Data characterized with no statistics.				
Domain 3: Access	sibility/Clar	ity								
Domain 9. Meess	Metric 6:	Metadata Completeness	Low	× 1	3	Provides results but generally does not provide data sources, methods, or assumptions.				
Domain 4: Variab	oility and H	ncertainty								
Domain 4. Variat	Metric 7:	Metadata Completeness	Low	× 1	3	Generally does not discuss variability or uncertainty.				
Overall Quality D)eterminatio	${f n}^\dagger$	Medium		1.7					

 $^{^{\}star}$ MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation:	Von Grote, J. 2003. Occupational Exposure Assessment in Metal Degreasing and Dry Cleaning -Influences of Technology Innovation and Legislation. A dissertation submitted to the Swiss Federal Institute of Technology Z"rich for the degree of								
Type of Data Source Hero ID	Doctor of Natural Sciences. Swiss Federal Institute of Technology Z"rich. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 5176439								
EXTRACTION Parameter			Data						
Life Cycle Stage: Route of Exposure:			Dry Cleaning Exposure Model Cited for:1) Residual solvent on garments2) Duration of finishing/ pressing3) Size of machine cylinders						
EVALUATION									
Domain		Metric	Rating	MWF*	Score	Comments			
Domain 1: Reliab	oility Metric 1:	Methodology	High	× 1	1	Academic PhD dissertation expected to use reliable data collection and analysis methods.			
Di 0. D	4-4:								
Domain 2: Repres	Metric 2:	Geographic Scope	Medium	× 1	2	Data based on German dry cleaners (OECD country).			
	Metric 3:	Applicability	High	$\times 2$	2	Dry cleaning data are applicable to the scope of the model.			
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Dissertation published in 2003. More than 10 but less than 20 years old.			
	Metric 5:	Sample Size	Medium	× 1	2	The various data elements used from the study are presented mostly as ranges or averages.			
Domain 3: Access	sibility/Clar	itv							
Domain 9. Access	Metric 6:	Metadata Completeness	Medium	× 1	2	Sources are cited, but does not provide details on how reported values were derived from cited sources.			
Domain 4: Variab	oility and Ur Metric 7:	ncertainty Metadata Completeness	Medium	× 1	2	Variability in parameter values discussed, but no discussion of uncertainties.			
Overall Quality Determination [†]			Medium		1.7				

^{*} MWF = Metric Weighting Factor † If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Source Citation: Type of Data Source Hero ID	Von Grote, J.,J. C. Hurlimann, Scheringer, M., Hungerbuhler, K 2003. Reduction of Occupational Exposure to Perchloroethylene and Trichloroethylene in Metal Degreasing over the Last 30 years: Influence of Technology Innovation and Legislation. Journal of Exposure Analysis and Environmental Epidemiology. Occupational Exposure; Reports for Data or Information Other than Exposure or Release Data; 3045042						
EXTRACTION Parameter			Data				
Life Cycle Stage: Route of Exposure:			Vapor and Cold Degreasing Exposure Models Cited for far-field volumes for inhalation exposure models.				
EVALUATION							
Domain		Metric	Rating	MWF*	Score	Comments	
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	Article is published in peer-reviewed scientific journal.	
Domain 2: Repres	antatira						
Domain 2. Kepres	Metric 2:	Geographic Scope	Medium	× 1	2	Data based on German facilities (OECD country).	
	Metric 3:	Applicability	High	$\times 2$	2	Degreasing facility data are applicable to the scope of the model.	
	Metric 4:	Temporal Representativeness	Medium	\times 2	4	Work based on dissertation published in 2003. More than 10 but less than 20 years old.	
	Metric 5:	Sample Size	Medium	× 1	2	The various data elements used from the study are presented mostly as ranges or averages.	
Domain 2. Access	::b::1:4/Closs	:4					
Domain 3: Access	Metric 6:	Metadata Completeness	Medium	× 1	2	Sources are cited, but does not provide details on how reported values were derived from cited sources.	
	*1**						
Domain 4: Variab	oility and Ur Metric 7:		Medium	× 1	2	Variability in parameter values discussed, but no discussion of uncertainties.	
Overall Quality Determination †			Medium		1.7		

^{*} MWF = Metric Weighting Factor

[†] If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .

Facility

Source Citation: Type of Data Source Hero ID	U.S. EPA. 2017. Public database 2016 chemical data reporting (May 2017 release). Facility; Reports for Data or Information Other than Exposure or Release Data; 3827204						
EXTRACTION Parameter			Data				
Life Cycle Stage: Life Cycle Description (Subcategory of Use): Total Annual U.S. Volume (and percent of PV): Number of Sites: Possible Physical Form: Chemical Concentration:			Manufacture and Import Manufacture and Import Provides U.S. domestic manufactured and imported PV and percent PV to downstream uses. Provides number of manufacturing and import sites. Provides physical form. Provides concentration.				
EVALUATION Domain		Metric	Rating	MWF*	Score	Comments	
Domain 1: Reliab	ility Metric 1:	Methodology	High	× 1	1	EPA is a trusted source.	
Domain 2: Repres	Metric 2: Metric 3: Metric 4: Metric 5:	Geographic Scope Applicability Temporal Representativeness Sample Size	High High High Medium	$\begin{array}{c} \times \ 1 \\ \times \ 2 \\ \times \ 2 \\ \times \ 1 \end{array}$	1 2 2	CDR is U.S. based data. CDR covers chemical manufacturers and importers, which are in scope for all chemicals. EPA used data from the 2016 CDR, which includes data reported for 2015. Due to reporting threshold, statistical representativeness is un-	
Domain 3: Access			Medium	× 1	2	Submissions do not include method of how production volumes were determined. CDR industry sector codes, industrial processing and use codes, industrial function codes, and commercial product codes provide good metadata; but lack of clarifying information and narratives and occasional misreportings limit clarity of data.	
Domain 4: Variab	ility and Ur Metric 7:	ncertainty Metadata Completeness	Low	× 1	3	CDR data do not address variability or uncertainty in submitter provided data.	
		Cor	ntinued on r	next page	9		

- continued from previous page

command from previous page									
Source Citation: Type of Data Source Hero ID	U.S. EPA. 2017. Public database 2016 chemical data reporting (May 2017 release). Facility; Reports for Data or Information Other than Exposure or Release Data; 3827204								
EVALUATION Domain	Metric	Rating	MWF* Score	Comments					
Overall Quality I	Determination [†]	High	1.4						

^{*} MWF = Metric Weighting Factor \dagger If any individual metrics are deemed Unacceptable, then the overall rating is also unacceptable. Otherwise, the overall rating is based on the following scale: High: ≥ 1 to < 1.7; Medium: ≥ 1.7 to < 2.3; Low: ≥ 2.3 to ≤ 3 .