


# Final Scope of the Risk Evaluation for 1,1,2-Trichloroethane

## CASRN 79-00-5



August 2020

## TABLE OF CONTENTS

A	KNOWLEDGEMENTS	7
A	BREVIATIONS AND ACRONYMS	8
E	ECUTIVE SUMMARY	11
1	INTRODUCTION	14
2	SCOPE OF THE EVALUATION	14
	.1 Reasonably Available Information	14
	2.1.1 Search of Gray Literature	
	2.1.2 Search of Literature from Publicly Available Databases (Peer-Reviewed Literature)	
	2.1.3 Search of TSCA Submissions	
	.2 Conditions of Use	
	2.2.1 Categories and Subcategories of Conditions of Use Included in the Scope of the Risk	
	Evaluation	
	2.2.2 Activities Excluded from the Scope of the Risk Evaluation	
	2.2.3 Production Volume	
	2.2.4 Overview of Conditions of Use and Lifecycle Diagram	
	3 Exposures	
	2.3.1 Physical and Chemical Properties	
	<ul><li>2.3.2 Environmental Fate and Transport</li><li>2.3.3 Releases to the Environment</li></ul>	
	2.3.4 Environmental Exposures	
	2.3.4 Environmental Exposures	
	2.3.6 Consumer Exposures	
	2.3.7 General Population Exposures	
	.4 Hazards (Effects)	
	2.4.1 Environmental Hazards	
	2.4.2 Human Health Hazards	
	.5 Potentially Exposed or Susceptible Subpopulations	
	.6 Conceptual Models	
	2.6.1 Conceptual Model for Industrial and Commercial Activities and Uses	38
	2.6.2 Conceptual Model for Consumer Activities and Uses	41
	2.6.3 Conceptual Model for Environmental Releases and Wastes: Potential Exposures and	
	Hazards (Regulatory Overlay)	
	2.6.3.1 Exposure Pathways and Risks Addressed by Other EPA Administered Statutes	
	2.6.3.2 Conceptual Model for Environmental Releases and Wastes: Potential Exposures	
	Hazards	
	7 Analysis Plan	
	<ul><li>2.7.1 Physical and Chemical Properties and Environmental Fate</li><li>2.7.2 Exposure</li></ul>	
	2.7.2 Exposure	
	2.7.2.1 Environmental Refeases	
	2.7.2.2 Environmental Exposures	
	2.7.2.4 Consumer Exposures	
	2.7.2.5 General Population	
	1 I	

2.7.3 Ha	zards (Effects)	
2.7.3.1		
2.7.3.2		
	mmary of Risk Approaches for Characterization	
	Review	
REFERENCI	ES	67
APPENDICE	S	73
Appendix A	ABBREVIATED METHODS FOR SEARCHING AND SCREENING	
A.1 Litera	ture Search of Publicly Available Databases	73
	arch Term Genesis and Chemical Verification	
	blicly Available Database Searches	
A.1.2.1	Query Strings for the Publicly Available Database Searches on 1,1,2-Trichloro	
A 1 2 2	Data Prioritization for Environmental Hazard, Human Health Hazard, Fate and	
	Chemistry	•
A.1.2.3	Data Prioritization for Occupational Exposures and Environmental Releases and	nd Gen
	Pop, Consumer and Environmental Exposures	
	Reviewed Screening Process	
	clusion/Exclusion Criteria	
	PECO for Environmental and Human Health Hazards	
	PECO for Consumer, Environmental, and General Population Exposures	
	RESO for Occupational Exposure and Environmental Releases	
	PESO for Fate and Transport	
	Generation of Hazard Heat Maps	
	Literature Search and Screening Strategies	
	tial Screening of Sources using Decision Logic Tree	
	CA Submission Searching and Title Screening	
	ay Literature Search Results for 1,1,2-Trichloroethane	
	PHYSICAL AND CHEMICAL PROPERTIES OF 1,1,2-TRICHLOROET	
••	ENVIRONMENTAL FATE AND TRANSORT PROPERTIES OF 1,1,2-	
ipponum e	TRICHLOROETHANE	
Appendix D	REGULATORY HISTORY	101
D.1 Feder	al Laws and Regulations	101
D.2 State	Laws and Regulations	107
	national Laws and Regulations	
Appendix E	PROCESS, RELEASE AND OCCUPATIONAL EXPOSURE INFORMAT	<b>FION 110</b>
E.1 Proce	ss Information	110
	anufacture (Including Import)	
E.1.1.1	Manufacture	
E.1.1.2	Import	
	ocessing and Distribution	
E.1.2.1	Processing as a Reactant or Intermediate	110

E.1.2.2	Recycling	111
E.1.3 Inc	lustrial and Commercial Uses	111
E.1.3.1	Industrial Use: Adhesives and Sealants	111
E.1.3.2	Commercial Use: Adhesives and Sealants	111
E.1.3.3	Commercial Use: Laboratory Use	111
E.1.4 Dis	sposal	111
	ninary Occupational Exposure Data	
	SUPPORTING INFORMATION - CONCEPTUAL MODEL FOR IND AND COMMERCIAL ACTIVITIES AND USES	
	SUPPORTING INFORMATION - CONCEPTUAL MODEL FOR CON ACTIVITIES AND USES	
11	SUPPORTING INFORMATION – CONCEPTUAL MODEL FOR ENVIRONMENTAL RELEASES AND WASTES	118

## LIST OF TABLES

Table 2-1. Results of Title Screening of Submissions to EPA under Various Sections of TSCA a	26
Table 2-2. Categories and Subcategories of Conditions of Use Included in the Scope of the Risk	
Evaluation	27
Table 2-3. Physical and Chemical Properties of 1,1,2-Trichloroethane	31
Table 2-4. Summary of 1,1,2-Trichloroethane TRI Production-Related Waste Managed in 2018	34
Table 2-5. Summary of Releases of 1,1,2-Trichloroethane to the Environment During 2018	34
Table 2-6. Categories and Sources of Environmental Release Data	55
Table 2-7. Potential Sources of Occupational Exposure Data	58

## LIST OF FIGURES

Figure 2-1. Gray Literature Tags by Discipline for 1,1,2-Trichloroethane
Figure 2-2. Peer-reviewed Literature Inventory Tree – Physical and Chemical Properties Search Results
for 1,1,2-Trichloroethane
Figure 2-3. Peer-reviewed Literature Inventory Tree - Fate and Transport Search Results for 18
Figure 2-4. Peer-reviewed Literature Inventory Heat Map – Fate and Transport Search Results for 1,1,2-
Trichloroethane
Figure 2-5. Peer-reviewed Literature Inventory Tree - Engineering Search Results for
Figure 2-6. Peer-reviewed Literature Inventory Heat Map - Engineering Search Results for 21
Figure 2-7. Peer-reviewed and Gray Literature Inventory Tree - Exposure Search Results for
Figure 2-8. Peer-reviewed and Gray Literature Inventory Heat Map - Exposure Search Results for 1,1,2-
Trichloroethane
Figure 2-9. Peer-reviewed Literature Inventory Tree - Human Health and Environmental Hazards Search
Results for 1,1,2-Trichloroethane
Figure 2-10. Peer-reviewed Literature Inventory Heat Map – Human Health and Environmental Hazards
Search Results for 1,1,2-Trichloroethane
Figure 2-11. 1,1,2-Trichloroethane Life Cycle Diagram
Figure 2-12. Box and Whisker Plots of Reported Physical and Chemical Property Values
Figure 2-13. 1,1,2-Trichloroethane Conceptual Model for Industrial and Commercial Activities and
Uses: Worker and Occupational Non-User Exposures and Hazards
Figure 2-14. 1,1,2-Trichloroethane Conceptual Model for Consumer Activities and Uses: Consumer
Exposures and Hazards
Figure 2-15. 1,1,2-Trichloroethane Conceptual Model for Environmental Releases and Wastes:
Environmental and General Population Exposures and Hazards (Regulatory Overlay) 44
Figure 2-16. 1,1,2-Trichloroethane Conceptual Model for Environmental Releases and Wastes:
Environmental Exposures and Hazards

## LIST OF APPENDIX TABLES

Table_Apx A-1. Sources of Verification for Chemical Names and Structures	73
Table_Apx A-2. Summary of Data Sources, Search Dates and Number of Peer-Reviewed Literature	
Search Results for 1,1,2-Trichloroethane	75
Table_Apx A-3. Hazards Title and Abstract and Full-text PECO Criteria for 1,1,2-Trichloroethane	83
Table_Apx A-4. Major Categories of Potentially Relevant Supplemental Materials for 1,1,2-	
Trichloroethane	84
Table_Apx A-5. General PECO for Consumer, Environmental, and General Population Exposures	85

Table_Apx A-6. Pathways Identified as Supplemental for 1,1,2-Trichloroethane <sup>a</sup>	85
Table_Apx A-7. Inclusion Criteria for Data Sources Reporting Environmental Release and C	Occupational
Exposure Data	-
Table_Apx A-8. Engineering, Environmental Release and Occupational Data Necessary to D	Develop the
Environmental Release and Occupational Exposure Assessments	
Table_Apx A-9. Inclusion Criteria for Data or Information Sources Reporting Environmenta	
Transport Data	
Table_Apx A-10. Fate Endpoints and Associated Processes, Media and Exposure Pathways	
in the Development of the Environmental Fate Assessment	
Table_Apx A-11. Decision Logic Tree Overview	
Table_Apx A-12. List of Gray Literature Sources for 1,1,2-Trichloroethane	
Table_Apx B-1. Summary Statistics for Reviewed Physical Properties	
Table_Apx C-1. Environmental Fate and Transport Properties of 1,1,2-Trichloroethane	
Table_Apx D-1. Federal Laws and Regulations	101
Table_Apx D-2. State Laws and Regulations	107
Table_Apx D-3. Regulatory Actions by other Governments, Tribes, and International Agree	ments 108
Table_Apx E-1. Summary of Industry Sectors with 1,1,2-Trichloroethane Personal Monitori	ng Air
Samples Obtained from OSHA Inspections Conducted since 1984	
Table_Apx E-2. Potentially Relevant Data Sources for Exposure Monitoring and Area Moni	toring Data
from NIOSH Health Hazard Evaluations (https://www.cdc.gov/niosh/hhe/)	112
Table_Apx F-1. Worker and Occupational Non-User Exposure Conceptual Model Supportin	g Table 113
Table_Apx G-1. Consumer Exposure Conceptual Model Supporting Table	117
Table_Apx H-1. General Population and Environmental Exposure Conceptual Model Suppo	rting Table

### LIST OF APPENDIX FIGURES

### **ACKNOWLEDGEMENTS**

This report was developed by the United States Environmental Protection Agency (U.S. EPA), Office of Chemical Safety and Pollution Prevention (OCSPP), Office of Pollution Prevention and Toxics (OPPT).

#### Acknowledgements

The OPPT Assessment Team gratefully acknowledges participation or input from Intra-agency reviewers that included multiple offices within EPA, Inter-agency reviewers that included multiple Federal agencies, and assistance from EPA contractors Abt Associates (Contract No. EP-W-16-009), ERG (Contract No. EP-W-12-006), GDIT (Contract No. HHSN316201200013W), ICF (Contract No.68HERC19D0003), SRC (Contract No. 68HERH19F0213), and Versar (Contract No. EP-W-17-006). EPA also acknowledges the contributions of technical experts from EPA's Office of Research and Development.

#### Docket

Supporting information can be found in public docket: [EPA-HQ-OPPT-2018-0421].

#### Disclaimer

Reference herein to any specific commercial products, process or service by trade name, trademark, manufacturer or otherwise does not constitute or imply its endorsement, recommendation or favoring by the United States Government.

## ABBREVIATIONS AND ACRONYMS

ACGIH	American Conference of Governmental Industrial Hygienists
ADME	Absorption, Distribution, Metabolism, and Excretion
AIA	Aerospace Industries Association
AMTIC	Ambient Monitoring Technology Information Center
ATSDR	Agency for Toxic Substances and Disease Registry
BAF	Bioaccumulation Factor
BCF	Bioconcentration Factor
BMF	Biomagnification factor
BOD	Biochemical Oxygen Demand
BP	Boiling Point
BSER	Best System of Emission Reduction
<b>BW</b> <sup>3/4</sup>	Body Weight <sup>3</sup> / <sub>4</sub> Extrapolation
CAA	Clean Air Act
CalEPA	California Environmental Protection Agency
CASRN	Chemical Abstracts Service Registry Number
CBI	Confidential Business Information
CDC	Centers for Diseases Control and Prevention
CDR	Chemical Data Reporting
CEHD	Chemical Exposure Health Data
CEM	Consumer Exposure Model
CEPA	Canadian Environmental Protection Act
CERCLA	Comprehensive Environmental Response, Compensation and Liability Act
CFR	Code of Federal Regulations
-	Chemical Screening Tool for Occupational Exposures and Releases
CHRIP	Chemical Risk Information Platform
COC	Concentration of Concern
CPCat	Chemical and Product Categories
CRC	Coordinating Research Council
CSCL	Chemical Substances Control Law
CWA	Clean Water Act
	C Clean Water Act- Ambient Water Quality Criteria
DMR	Discharge Monitoring Report
DOT	Department of Transportation
EC	Engineering Control(s)
ECHA	European Chemicals Agency
EPA	Environmental Protection Agency
EPCRA	Emergency Planning and Community Right-to-Know Act
EPI	Estimation Programs Interface
ERG	Eastern Research Group
ESD	Emission Scenario Document
ESDU	Engineering Sciences Data Unit
EU	European Union
FDA	Food and Drug Administration
FFDCA	Federal Food, Drug, and Cosmetic Act
FR	Federal Register
FYI	For your information
	-

CACT	Conservables Associately Constant Track and a serv
GACT	Generally Available Control Technology
GDIT	General Dynamics Information Technology
GESTIS	Substance Database contains information for the safe handling of hazardous substances
00	and other chemical substances at work
GS	Generic Scenario
HAP	Hazardous Air Pollutant
HERO	Health and Environmental Research Online
HHE	Health Hazard Evaluation
HLC	Henry's Law Constant
HMTA	Federal Hazardous Materials Transportation Act
HPLC	High Performance Liquid Chromatography
HSDB	Hazardous Substances Data Bank
IARC	International Agency for Research on Cancer
ICF	ICF is a global consulting services company
IMAP	Inventory Multi-Tiered Assessment and Prioritisation (Australia)
IRIS	Integrated Risk Information System
ISHA	Industrial Safety and Health Act
K <sub>OC</sub>	Organic Carbon: Water Partition Coefficient
K <sub>OW</sub>	Octanol: Water Partition Coefficient
LOAELs	Lowest Observed Adverse Effect Level
MACT	Maximum Achievable Control Technology
MCL	Maximum Contaminant Level
MCLG	Maximum Contaminant Level Goal
MITI	Ministry of International Trade and Industry
MOA	Mode of Action
MP	Melting Point
MSW	Municipal Solid Waste
NATA	National-scale Air Toxics Assessment
NBOS	National Bureau of Standards
NCI	National Cancer Institute
NEI	National Emissions Inventory
NESHAP	National Emission Standards for Hazardous Air Pollutants
NHANES	National Health and Nutrition Examination Survey
NICNAS	National Industrial Chemicals Notification and Assessment Scheme (Australia)
NIOSH	National Institute for Occupational Safety and Health
NITE	National Institute of Technology and Evaluation
NLM	National Library of Medicine
NOAELs	No Observed Adverse Effect Level
NOEC	No Observed Effect Concentration
NPDES	National Pollutant Discharge Elimination System
NPDWR	National Primary Drinking Water Regulation
NPL	National Priorities List
NPRI	National Pollutant Release Inventory
NSPS	New Source Performance Standards
NTP	National Toxicology Program
OCSPP	Office of Chemical Safety and Pollution Prevention
OECD	Organisation for Economic Co-operation and Development

OEHHA	Office of Environmental Health Hazard Assessment (California)
ONU	Occupational Non-User
OPPT	Office of Pollution Prevention and Toxics
OSHA	Occupational Safety and Health Administration
OW	EPA's Office of Water
PBPK	Physiologically Based Pharmacokinetic
PBT	Persistent Bioaccumulation, and Toxic
PECO	Population, Exposure, Comparator, Outcome
PEL	Permissible Exposure Limit
PESO	Pathways and Processes, Exposures, Setting or Scenario and Outcomes
PESS	Potentially Exposed or Susceptible Subpopulation
PODs	Points of Departure
POTW	Publicly Owned Treatment Works
PPE	Personal Protective Equipment
RCRA	Resource Conservation and Recovery Act
REACH	Registration, Evaluation, Authorisation and Restriction of Chemicals (European Union)
REL	Recommended Exposure Limit
RESO	Receptors, Exposure, Setting or Scenario and Outcomes
RQs	Risk Quotients
SARA	Superfund Amendments and Reauthorization Act
SDS	Safety Data Sheet
SDWA	Safe Drinking Water Act
SIDS	Screening Information Dataset
SNAc	Significant New Activity
SRC	SRC Inc., formerly Syracuse Research Corporation
SYKE	The Finnish Environment Institute
TBD	To be determined
TCA	Trichloroethane
TG	Test Guideline
TIAB	Title and Abstract
TLV	Threshold Limit Value
TMF	Trophic Magnification Factors
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TSD	Technical Support Document
TTO	Total Toxic Organics
TURA	Toxic Use Reduction Act
TWA	Time-weighted average
UCMR	Unregulated Contaminants Monitoring Rule
UIC	Underground Injection Control
USGS	United States Geological Survey
VOC	Volatile Organic Compound
VP	Vapor Pressure
WS	Water Solubility
WWT	Wastewater Treatment

## **EXECUTIVE SUMMARY**

In December 2019, EPA designated 1,1,2-trichloroethane (CASRN 79-00-5) as a high priority substance for risk evaluation following the prioritization process as required by Section 6(b) of the Toxic Substances Control Act (TSCA) and implementing regulations (40 CFR Part 702) (Docket ID: <u>EPA-HQ-OPPT-2019-0131</u>). The first step of the risk evaluation process is the development of the draft scope document. EPA published the *Draft Scope of the Risk Evaluation for 1,1,2-Trichloroethane CASRN 95-50-1* (EPA Document No. EPA-740-R-20-003) (<u>U.S. EPA, 2020c</u>) and provided a 45-day comment period on the draft scope per 40 CFR 702.41(c)(7). EPA has considered comments received (Docket ID: <u>EPA-HQ-OPPT-2018-0421</u>) during the public comment period to inform the development of this final scope document, and public comments received will continue to inform the development of the risk evaluation for 1,1,2-trichloroethane. This document fulfills the TSCA requirement to issue a final scope document per TSCA Section 6(b)(4)(D) and as described in 40 CFR 702.41(c)(8). The scope for 1,1,2trichloroethane includes the following information: the conditions of use, potentially exposed or susceptible subpopulations (PESS), hazards, and exposures that EPA plans to consider in this risk evaluation, along with a description of the reasonably available information, conceptual model, analysis plan and science approaches, and plan for peer review for this chemical substance.

*General Information*. 1,1,2-Trichloroethane (CASRN 79-00-5) is a colorless, sweet-smelling liquid. It is generally nonflammable, can be dissolved in water, and evaporates easily (volatilization half-life of 35 minutes). Although this liquid is nonflammable, containers of 1,1,2-trichloroethane may explode in the heat of a fire. It is used as a solvent and as an intermediate in the production of other chemicals. 1,1,2-Trichloroethane is sometimes present as an impurity in other chemicals, and it may be formed when another chemical breaks down in the environment under conditions where there is no air.

**Reasonably Available Information.** EPA leveraged the data and information sources already described in the *Proposed Designation of 1,1,2-trichloroethane (CASRN 79-00-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019d) to inform the development of this scope document. Furthermore, EPA conducted a comprehensive search to identify and screen multiple evidence streams (*i.e.*, chemistry, fate, release and engineering, exposure, hazard), and the search and screening results are provided in Section 2.1. EPA used the systematic review process described in Appendix A to search for and screen reasonably available information, including information already in EPA's possession, for inclusion in the risk evaluation. This information includes the hazards, exposures, PESS, and conditions of use that may help inform the risk evaluation for 1,1,2-trichloroethane. EPA has focused on the data collection phase (consisting of data search, data screening, and data extraction) during the preparation of the scope document, whereas the data evaluation and integration stages will occur during the development of the risk evaluation and thus are not part of the scoping activities described in this document. EPA will consider additional information identified following publication of this scope document, as appropriate, in developing the risk evaluation, including the Chemical Data Reporting Rule (CDR) information the Agency will receive by the end of November 2020.

*Conditions of Use.* EPA plans to evaluate manufacturing (including importing), processing, distribution in commerce, industrial, commercial and consumer uses, and disposal of 1,1,2-trichloroethane in the risk evaluation. 1,1,2-Trichloroethane is manufactured (including imported) in the United States. The conditions of use include processing as a reactant and other processing activities such as recycling. Industrial and commercial uses identified include adhesives and sealants and use as a laboratory chemical. EPA plans to evaluate the use of 1,1,2-trichloroethane produced as a byproduct from the manufacture of 1,2-dichloroethane (CASRN 107-06-2) in the risk evaluation for 1,2-dichloroethane and

not in the risk evaluation for 1,1,2-trichloroethane. The only consumer use reported is in adhesives and sealants. EPA identified these conditions of use from information reported to EPA through CDR and Toxics Release Inventory (TRI) reporting, published literature, and consultation with stakeholders for both uses currently in production and uses whose production may have ceased. EPA revised the conditions of use in the final scope of the risk evaluation based on additional information and public comments (Docket ID: EPA-HQ-OPPT-2018-0421) on the draft scope document for 1,1,2-trichloroethane. Section 2.2 provides details about the conditions of use within the scope of the risk evaluation.

*Conceptual Model.* The conceptual models for 1,1,2-trichloroethane are presented in Section 2.6. Conceptual models are graphical depictions of the actual or predicted relationships of conditions of use, exposure pathways (*e.g.*, media), exposure routes (*e.g.*, inhalation, dermal, oral), hazards and receptors throughout the life cycle of the chemical substance. EPA considered reasonably available information as well as public comments received on the draft scope document for 1,1,2-trichloroethane in finalizing the exposure pathways, exposure routes, and hazards EPA plans to evaluate in the risk evaluation. As a result, EPA plans to focus the risk evaluation for 1,1,2-trichloroethane on the following exposures, hazards and receptors.

• *Exposures (Pathways and Routes), Receptors and PESS.* EPA plans to evaluate releases to the environment as well as human and environmental exposures resulting from the conditions of use of 1,1,2-trichloroethane that EPA plans to consider in the risk evaluation. Exposures for 1,1,2-trichloroethane are discussed in Section 2.3. Additional information gathered through the results of systematic review searches will also informed expected exposures.

EPA's plan for evaluating environmental exposure pathways in the scope of the risk evaluation considers whether and how other EPA administered statutes and regulatory programs cover the presence of 1,1,2-trichloroethane in media pathways falling under the jurisdiction of those authorities. Section 2.6.3.1 discusses pathways under the jurisdiction of other EPA-administered laws. In Section 2.6.3.2, EPA presents the conceptual model describing the identified exposures (pathways and routes), receptors and hazards associated with the conditions of use of 1,1,2-trichloroethane within the scope of the risk evaluation. Based on pathways under the jurisdiction of other EPA-administered laws, EPA does not plan to evaluate the general population as a receptor in the scope of the risk evaluation (see Figure 2-16).

EPA considered reasonably available information and comments received on the draft scope for 1,1,2-trichloroethane in determining the human and environmental exposure pathways, routes, and receptors for inclusion in the final scope. EPA plans to evaluate\_the following human and environmental exposure pathways, routes, and receptors in the scope of the risk evaluation:

- *Occupational exposure:* EPA plans to evaluate exposures to workers and occupational non-users (ONUs) via the inhalation route and exposures to workers via the dermal route associated with manufacturing, processing, use or disposal of 1,1,2-trichloroethane.
- Consumer and bystander exposure: EPA plans to evaluate inhalation exposures to 1,1,2trichloroethane for consumers and bystanders and dermal exposures to 1,1,2trichloroethane for consumers during use of adhesives and sealants.
- *PESS:* EPA plans to include children, women of reproductive age (*e.g.*, pregnant women), workers and consumers as receptors and PESS in the risk evaluation.

- *Environmental exposure:* EPA plans to evaluate exposure to 1,1,2-trichloroethane for aquatic receptors.
- *Hazards*. Hazards for 1,1,2-trichloroethane are discussed in Section 2.4. EPA completed preliminary reviews of information (*e.g.*, federal and international government chemical assessments) to identify potential environmental and human health hazards for 1,1,2-trichloroethane as part of the prioritization (U.S. EPA, 2019d) and scoping process (U.S. EPA, 2020c). EPA also considered reasonably available information collected through systematic review methods as outlined in Appendix A and public comments received on the draft scope for 1,1,2-trichloroethane in determining the broad categories of environmental and human health hazard effects to be evaluated in the risk evaluation. EPA will use systematic review methods to evaluate the epidemiological and toxicological literature for 1,1,2-trichloroethane.

EPA plans to evaluate all potential environmental and human health hazard effects identified for 1,1,2-trichloroethane in Sections 2.4.1 and 2.4.2, respectively. Identified through the data screening phase of systematic review, the potential environmental hazard effects and related information that EPA plans to consider for the risk evaluation include: developmental, mortality, musculoskeletal, reproductive and respiratory effects for 1,1,2-trichloroethane. Similarly, the potential human health hazard effects and related information identified through prioritization and the data screening phase of systematic review for 1,1,2-trichloroethane that EPA plans to consider for the risk evaluation include: ADME, PBPK, cancer, cardiovascular, developmental, endocrine, gastrointestinal, hematological and immune, hepatic, mortality, musculoskeletal, neurological, nutritional and metabolic, ocular and sensory, renal, reproductive, respiratory and skin and connective tissue effects.

*Analysis Plan.* The analysis plan for 1,1,2-trichloroethane is presented in Section 2.7. The analysis plan outlines the general science approaches that EPA plans to use for the various information streams (*i.e.*, chemistry, fate, release and engineering, exposure, hazard) supporting the risk evaluation. The analysis plan is based on EPA's knowledge of 1,1,2-trichloroethane to date, which includes a review of identified information as described in Section 2.1. Should additional data or approaches become reasonably available, EPA may consider them for the risk evaluation

*Peer Review.* The draft risk evaluation for 1,1,2-trichloroethane will be peer reviewed. Peer review will be conducted in accordance with relevant and applicable methods for chemical risk evaluations, including using EPA's Peer Review Handbook (U.S. EPA, 2015b) and other methods consistent with Section 26 of TSCA (see 40 CFR 702.45).

## **1 INTRODUCTION**

This document presents the scope of the risk evaluation to be conducted for 1,1,2-trichloroethane under the Frank R. Lautenberg Chemical Safety for the 21st Century Act. The Frank R. Lautenberg Chemical Safety for the 21st Century Act amended TSCA on June 22, 2016. The new law includes statutory requirements and deadlines for actions related to conducting risk evaluations of existing chemicals.

Under TSCA § 6(b), the Environmental Protection Agency (EPA) must designate chemical substances as high-priority substances for risk evaluation or low-priority substances for which risk evaluations are not warranted at the time, and upon designating a chemical substance as a high-priority substance, initiate a risk evaluation on the substance. TSCA § 6(b)(4) directs EPA to conduct risk evaluations for existing chemicals, to "*determine whether a chemical substance presents an unreasonable risk of injury to health or the environment, without consideration of costs or other nonrisk factors, including an unreasonable risk to a potentially exposed or susceptible subpopulation identified as relevant to the risk evaluation by the Administrator under the conditions of use.*"

TSCA § 6(b)(4)(D) and implementing regulations require that EPA publish the scope of the risk evaluation to be conducted, including the hazards, exposures, conditions of use and PESS that the Administrator expects to consider, within 6 months after the initiation of a risk evaluation. In addition, a draft scope is to be published pursuant to 40 CFR 702.41. In December 2019, EPA published a list of 20 chemical substances that have been designated high-priority substances for risk evaluations (Docket ID: EPA-HQ-OPPT-2019-0131) (84 FR 71924, December 30, 2019), as required by TSCA § 6(b)(2)(B), which initiated the risk evaluation process for those chemical substances. 1,1,2-Trichloroethane is one of the chemicals designated as a high priority substance for risk evaluation. On April 23, 2020, EPA published the *Draft Scope of the Risk Evaluation for 1,1,2-Trichloroethane* (EPA Document No. (85 FR 22733, April 23, 2020) (U.S. EPA, 2020c) for a 45-day public comment period. After reviewing and considering the public comments received (Docket ID: EPA-HQ-OPPT-2018-0421) on the draft scope document, EPA is now publishing this final scope document pursuant to 40 CFR 702.41(c)(8).

## **2** SCOPE OF THE EVALUATION

### 2.1 Reasonably Available Information

EPA conducted a comprehensive search for reasonably available information<sup>1</sup> to support the development of this draft scope document for 1,1,2-trichloroethane. EPA leveraged the data and information sources already collected in the documents supporting the high-priority substance designations. In addition, EPA searched for additional data and information on physical and chemical properties, environmental fate, engineering, exposure, environmental and human health hazards that could be obtained from in the following general categories of sources:

- 1. Databases containing publicly available, peer-reviewed literature;
- 2. Gray literature, which is defined as the broad category of data/information sources not found in standard, peer-reviewed literature databases;

<sup>&</sup>lt;sup>1</sup>*Reasonably available information* means information that EPA possesses or can reasonably generate, obtain, and synthesize for use in risk evaluations, considering the deadlines specified in TSCA Section 6(b)(4)(G) for completing such evaluation. Information that meets the terms of the preceding sentence is reasonably available information whether or not the information is confidential business information, that is protected from public disclosure under TSCA Section 14 (40 CFR 702.33).

3. Data and information submitted under TSCA Sections 4, 5, 8(e), and 8(d), as well as "for your information" (FYI) submissions.

Following the comprehensive search, EPA performed a title and abstract screening to identify information potentially relevant for the risk evaluation process. This step also classified the references into useful categories or tags to facilitate the sorting of information through the systematic review process.

Search terms were used to search each of the literature streams and gather 1,1,2-trichloroethane studies. These terms and the methods used to develop them are listed in Appendix A. The studies resulting from the search process were loaded into the EPA Health and Environmental Research Online (HERO) database and then prioritized to screen first the literature likely relevant for each of the disciplines: fate, physical and chemical properties, engineering, exposure and hazard. The tools and methods used to manage the screening process are also outlined in Appendix A. The studies resulting from the search underwent a title/abstract screening process, which tagged them by topic or category. Following this, a determination was made to move studies forward into full-text screening. The criteria used in the screening process for each discipline are found in the population, exposure, comparator, outcome (PECO) statements listed in Appendix A. The screening process results are presented in the form of literature inventory trees and heat maps in Section 2.1.2. The screening process was conducted based on EPA's planning, execution and assessment activities outlined in Appendix A.


EPA has focused on the data collection phase (consisting of data search, data screening, and data extraction) during the preparation of the scope document, whereas the data evaluation and integration stages will occur during the development of the risk evaluation and thus are not part of the scoping activities described in this document.

The subsequent sections summarize the data collection activities completed up to date for the general categories of sources and topic areas (or disciplines) using systematic review methods.

#### 2.1.1 Search of Gray Literature

EPA surveyed the gray literature<sup>2</sup> and identified 173 search results relevant to EPA's risk evaluation needs for 1,1,2-trichloroethane. Appendix A.3.4 lists the gray literature sources that yielded 173 discrete data or information sources relevant to 1,1,2-trichloroethane. EPA further categorized the data and information into the various topic areas (or disciplines) supporting the risk evaluation (*e.g.*, physical and chemical properties, environmental fate, ecological hazard, human health hazard, exposure, engineering), and the breakdown is shown in Figure 2-1.. EPA will consider additional reasonably available information from gray literature if it becomes available during the risk evaluation phase.

 $<sup>^2</sup>$  Gray literature is defined as the broad category of data/information sources not found in standard, peer-reviewed literature databases (*e.g.*, PubMed and Web of Science). Gray literature includes data/information sources such as white papers, conference proceedings, technical reports, reference books, dissertations, information on various stakeholder websites, and other databases.



#### Figure 2-1. Gray Literature Tags by Discipline for 1,1,2-Trichloroethane


The percentages across disciplines do not add up to 100%, as each source may provide data or information for various topic areas (or disciplines).

## 2.1.2 Search of Literature from Publicly Available Databases (Peer-Reviewed Literature)

EPA has begun the systematic review process and has conducted searching and screening of the reasonably available literature using the process outlined in Appendix A. This includes performing a comprehensive search of the reasonably available peer review literature on physical and chemical properties, environmental fate and transport, engineering (environmental release and occupational exposure), exposure (environmental, general population and consumer) and environmental and human health hazards of 1,1,2-trichloroethane. Eligibility criteria were applied in the form of PECO statements (see Appendix A). Included references met the PECO criteria, whereas excluded references did not meet the criteria (*i.e.*, not relevant), and supplemental material was considered as potentially relevant (see Appendix A.2). EPA plans to analyze the reasonably available information identified for each discipline during the development of the risk evaluation.


EPA created literature inventory trees to graphically illustrate the flow of data and information sources following full-text screening (see Figure 2-2, Figure 2-3, Figure 2-5, Figure 2-7, and Figure 2-9). For the physical and chemical, fate, engineering and hazard literature, EPA used the Health Assessment Workplace Collaborative (HAWC) tool to develop web-based literature inventory trees illustrating, through interactive links, studies that were included or excluded. These literature inventory trees enhance the transparency of the decisions resulting from the screening process described in Appendix A. For each of the corresponding disciplines, the literature was tagged to be included for evaluation during the risk evaluation. Literature inventory trees for physical and chemical properties are provided as static diagrams (Figure 2-2). For all other disciplines, static screen captures are provided in addition to links to the interactive trees, which are provided in their corresponding captions. The links show individual studies that were tagged as included, excluded or supplemental. Supplemental studies did not meet all inclusion criteria but may be considered during the risk evaluation as supporting information (see Appendix A). These studies can be accessed through the hyperlink provided in the associated caption below each figure. In some figures, the sum of the numbers for the various sub-categories may be larger than the broader category because some studies may be included under multiple sub-categories. In other

cases, the sum of the various sub-categories may be smaller than the main category because some studies may not be depicted in the sub-categories if their relevance to the risk evaluation was unclear. In addition, EPA tabulated the number and characteristics of the data and information sources included in the full-text screening process in the form of a literature heat map for the fate, engineering, exposure and hazard information (see Figure 2-4, Figure 2-6, Figure 2-8, Figure 2-10). For each of these four disciplines, a static image of the literature inventory heat map is provided, and a link to the interactive version presented in HAWC is included in the caption below each diagram.



## Figure 2-2. Peer-reviewed Literature Inventory Tree – Physical and Chemical Properties Search Results for 1,1,2-Trichloroethane

Data in this static figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of June 2, 2020. TIAB refers to "title and abstract" screening.




## Figure 2-3. Peer-reviewed Literature Inventory Tree – Fate and Transport Search Results for 1,1,2-Trichloroethane

Click <u>here</u> to view the interactive literature inventory tree. Data in this figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of June 2, 2020. Additional data may be added to the interactive version as they become available.

			Med	lia		
Endpoint	Air	Soil, Sediment	Wastewater, Biosolids	Water	Other	Grand Total
Bioconcentration		0	1	1		2
Biodegradation		3	1	6		9
Hydrolysis	2	2		5		6
Photolysis	4	1		2		4
Sorption	1	6		7	1	7
Volatilization	1	2	1	3		3
Wastewater Treatment			2	1		2
Other	1	1	2	2	1	3
Grand Total	6	13	5	18	1	26

## Figure 2-4. Peer-reviewed Literature Inventory Heat Map – Fate and Transport Search Results for 1,1,2-Trichloroethane

Click <u>here</u> to view the interactive version for additional study details. The column totals, row totals, and grand totals indicate total numbers of unique references, as some references may be included in multiple cells. The various shades of color visually represent the amount of relevant references identified by media or endpoint. The darker the color, the more references are available for a given media or endpoint. Data in this figure represents references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of June 2, 2020. Additional data may be added to the interactive version as they become available.




## **Figure 2-5. Peer-reviewed Literature Inventory Tree - Engineering Search Results for 1,1,2-Trichloroethane**

Click <u>here</u> to view the interactive literature inventory tree. Data in this figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of August 5, 2020. Additional data may be added to the interactive version as they become available.

Data Type 💈	Evidence Tags	
	Description of release source	19
	Release frequency	1
Environmental	Release or emission factors	16
Releases	Release quantity	15
	Waste treatment methods and pollution control	16
	Total	34
	Chemical concentration	5
	Life cycle description	6
General	No evidence tag	7
Engineering	Number of sites	10
Assessment	Process description	15
Assessment	Production, import, or use volume	17
	Throughput	5
	Total	41
	Area sampling data	14
	Dermal exposure data	6
	Engineering control	3
	Exposure duration	9
	Exposure frequency	4
	Exposure route	20
Occupational	No evidence tag	5
Exposures	Number of workers	10
	Particle size characterization	
	Personal protective equipment	4
	Personal sampling data	9
	Physical form	12
	Worker activity description	16
	Total	35
Grand Total		69

## Figure 2-6. Peer-reviewed Literature Inventory Heat Map - Engineering Search Results for 1,1,2-Trichloroethane

Click <u>here</u> to view the interactive version for additional study details. Data in this figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of August 5, 2020. Additional data may be added to the interactive version as they become available.




## Figure 2-7. Peer-reviewed and Gray Literature Inventory Tree - Exposure Search Results for 1,1,2-Trichloroethane

Click <u>here</u> to view the interactive literature inventory tree. Data in this figure represent all references obtained from the publicly available databases search (see Appendix A.1.2.), and gray literature references search (see Appendix A.3) that were included during full-text screening as of July 31, 2020. Additional data may be added to the interactive version as they become available.

	Data Type							
Media (group)	Monitoring Study	Modeling Study	Completed Assessment	Experimental Study	Epidemiological Study	Database	Survey	Grand Total
Ambient Air								
Biosolids/Sludge			1					1
Drinking Water								
Groundwater								
Land Disposal/Landfill								
Sediment	2	1	2					3
Soil	4	1	1					5
Surface Water	2	1 1	2			1		5 4 1
Wastewater			1					1
Aquatic Species	1							1
Terrestrial Species								
Consumer	10	3	2	12				22
Dietary	9	1	1	2	1	1	2	10
Dust								
Exposure Factors	6	6	2	1		3	2	12
Exposure Pathway	8	2	3	1		1	2	10
Human Biomonitoring	33	3	5	4	4	6	3	38
Indoor Air	95	9	18	11	6	3	10	112
Isomers	109	12	12	17	9	8	13	129
Use Information	3	1	4					6
No Evidence Type								
Grand Total	130	16	23	19	12	12	15	158

## Figure 2-8. Peer-reviewed and Gray Literature Inventory Heat Map –Exposure Search Results for 1,1,2-Trichloroethane

Click <u>here</u> to view the interactive version for additional study details. The column totals, row totals, and grand totals indicate total numbers of unique references only, as some references may be included in multiple cells. The various shades of color visually represent the amount of relevant references identified by exposure media or data type. The darker the color, the more references are available for a given exposure media or data type. Data in this figure represent all references obtained from the publicly available databases search (see Appendix A.1.2), and gray literature references search (see Appendix A.3) that were included during full-text screening as of July 31, 2020. Additional data may be added to the interactive version as they become available.



## Figure 2-9. Peer-reviewed Literature Inventory Tree - Human Health and Environmental Hazards Search Results for 1,1,2-Trichloroethane

Click <u>here</u> to view the interactive literature inventory tree. Data in this figure represent references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of June 2, 2020. Additional data may be added to the interactive version as they become available.

Health Outcomes	Human	Animal - Human Health Model	Animal - Environmental Model	Plant	Grand Total
ADME	6	20			25
Cancer	5	6			11
Cardiovascular	3	1			4
Developmental	7	8	5	1	21
Endocrine	1	4			5
Gastrointestinal	3	2			5
Hematological and Immune	1	13			14
Hepatic		12			12
Mortality	2	2	4		8
Musculoskeletal	1	1	1		3
Neurological	4	5			9
Nutritional and Metabolic	1	2			3
Ocular and Sensory	2	3			5
РВРК		1			1
Renal	1	6			7
Reproductive	5	3	5		13
Respiratory	2	15	1		18
Skin and Connective Tissue	3	4			6
No Tag	1	4	8	1	14
Grand Total	15	42	17	2	75

#### Evidence Type

## Figure 2-10. Peer-reviewed Literature Inventory Heat Map – Human Health and Environmental Hazards Search Results for 1,1,2-Trichloroethane

Click <u>here</u> to view the interactive version for additional study details. The numbers indicate the number of studies with TIAB keywords related to a particular health outcome, not the number of studies that observed an association with 1,1,2-trichloroethane. Evidence types were manually extracted, and Health Systems were determined via machine learning. Therefore, the studies examining multiple Health Outcomes and Evidence types, connections between health outcome, and evidence type may not be accurately represented. If a study evaluated multiple health outcomes or included multiple populations or study designs, it is shown here multiple times. Data in this figure represents references obtained from the publicly available databases search (see Appendix A.1.2) that were included during full-text screening as of June 2, 2020. Additional data may be added to the interactive version as they become available.

### 2.1.3 Search of TSCA Submissions

Table 2-1 presents the results of screening the titles of data sources and reports submitted to EPA under various Sections of TSCA. EPA screened a total of 128 submissions using PECO or similar statements that identify inclusion/exclusion criteria specific to individual disciplines. Details of the criteria are presented in Appendix A.2.1. EPA identified 98 submissions that met the inclusion criteria in these

statements and identified 14 submissions with supplemental data.<sup>3</sup> EPA excluded 16 submissions because the reports were identified as one of the following:

- Published report that would be identified via other peer or gray literature searches
- Draft report of a final available submitted report
- Environmental impact statement for proposed equipment
- Status report
- Letter of intent
- Ranking of chemicals for proposed evaluation
- Submission on a different chemical
- Progress report
- Route-to-route extrapolation of human health hazard with no original data.

Tuble 2 11 Rebuild of The Screening of Submissions to 21 If ander + artous Sections of The Str							
Discipline	Included	Supplemental <sup>b</sup>					
Physical and Chemical Properties	0	0					
Environmental Fate and Transport	12	0					
Environmental and General Population Exposure	56	2					
Occupational Exposure/Release Information	14	0					
Environmental Hazard	3	2					
Human Health Hazard	26	10					

#### Table 2-1. Results of Title Screening of Submissions to EPA under Various Sections of TSCA <sup>a</sup>

<sup>a</sup> Individual submissions may be relevant to multiple disciplines.

<sup>b</sup> Included submissions may contain supplemental data for other disciplines, which will be identified at full-text review.

### 2.2 Conditions of Use

As described in the *Proposed Designation of 1,1,2-trichloroethane (CASRN 79-00-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019d), EPA assembled information from the CDR and TRI programs to determine conditions of use<sup>4</sup> or significant changes in conditions of use of the chemical substance. Once the 2020 CDR reporting period ends in November 2020, EPA will utilize the most recent CDR information. EPA also consulted a variety of other sources to identify uses of 1,1,2-trichloroethane, including: published literature, company websites, and government and commercial trade databases and publications. To identify formulated products containing1,1,2-trichloroethane, EPA searched for safety data sheets (SDS) using internet searches, EPA Chemical and Product Categories (CPCat) (U.S. EPA, 2019c) data, and other resources in which SDSs could be found. SDSs were cross-checked with company websites to make sure that each product SDS was current. In addition, EPA incorporated communications with companies, industry groups, and public comments to supplement the use information.

EPA identified and described the categories and subcategories of conditions of use that EPA plans to consider in the risk evaluation (Section 2.2.1; Table 2-2). The conditions of use included in the scope are those reflected in the life cycle diagrams and conceptual models.

<sup>&</sup>lt;sup>3</sup> EPA may further consider some supplemental or excluded references depending on the reasons for tagging as supplemental or excluded.

<sup>&</sup>lt;sup>4</sup> *Conditions of use* means the circumstances, as determined by the Administrator, under which a chemical substance is intended, known, or reasonably foreseen to be manufactured, processed, distributed in commerce, used, or disposed of (TSCA § 3(4)).

After gathering reasonably available information related to the manufacture, processing, distribution in commerce, use, and disposal of 1,1,2-trichloroethane, EPA identified those activities for 1,1,2-trichloroethane the Agency determined not to be conditions of use or are otherwise excluded from the scope of the risk evaluation. These excluded activities are described in Section 2.2.2.

## 2.2.1 Categories and Subcategories of Conditions of Use Included in the Scope of the Risk Evaluation

Table 2-2 lists the conditions of use that are included in the scope of the risk evaluation.

Life Cycle Stage <sup>a</sup>	Category <sup>b</sup>	Subcategory <sup>c</sup>	References
Manufacturing	Domestic manufacture	Domestic manufacture	<u>U.S. EPA (2019a)</u>
	Import	Import	<u>U.S. EPA (2019a)</u>
Processing	Processing as a reactant	Intermediate in: Plastic manufacturing; Petrochemical manufacturing; All other chemical product manufacturing	<u>U.S. EPA (2019a)</u>
	Recycling	Recycling	U.S. EPA (2019a)
Distribution in commerce	Distribution in commerce	Distribution in commerce	<u>U.S. EPA (2019a)</u>
Industrial Use	Adhesives and sealants	Adhesive and sealants	EPA-HQ-OPPT-2018- 0421-0006
Commercial Use	Adhesives and sealants	Adhesives and sealants	EPA-HQ-OPPT-2018- 0427-0003; EPA-HQ- OPPT-2018-0421-0013
	Other use	Laboratory chemical	<u>U.S. EPA (2019a)</u>
Consumer Use	Adhesives and sealants	Adhesives and sealants	EPA-HQ-OPPT-2018- 0427-0003; EPA-HQ- OPPT-2018-0421-0013
Disposal	Disposal	Disposal	

 Table 2-2. Categories and Subcategories of Conditions of Use Included in the Scope of the Risk

 Evaluation

a. Life Cycle Stage Use Definitions (40 CFR § 711.3)

 "Industrial use" means use at a site at which one or more chemicals or mixtures are manufactured (including imported) or processed.

 "Commercial use" means the use of a chemical or a mixture containing a chemical (including as part of an article) in a commercial enterprise providing saleable goods or services.

- "Consumer use" means the use of a chemical or a mixture containing a chemical (including as part of an article, such as furniture or clothing) when sold to or made available to consumers for their use.

Although EPA has identified both industrial and commercial uses here for purposes of distinguishing scenarios in this document, the Agency interprets the authority over "any manner or method of commercial use" under TSCA Section 6(a)(5) to reach both.

b. These categories of conditions of use appear in the Life Cycle Diagram, reflect CDR codes, and broadly represent conditions of use of 1,1,2-trichloroethane in industrial and/or commercial settings and for consumer uses.

c. These subcategories reflect more specific conditions of use of 1,1,2-trichloroethane.

Life Cycle Stage <sup>a</sup>	Category <sup>b</sup>	Subcategory <sup>c</sup>	References					
- The draft scope document included an industrial condition of use for non-incorporative activities – Other in all other chemical product and preparation manufacturing. Further discussion with the reporting company determined this non-								
		ne as an unintended byproduct during						
dichloroethean (CASR)	N 107-06-2) ( <u>EPA-HQ-OPP</u>	<u>T-2018-0421-0029</u> ). Public comment	ts submitted to EPA in the					
docket (EPA-HQ-OPP	docket (EPA-HQ-OPPT-2018-0421-0027) additionally indicated 1,1,2-trichloroethane is produced as a reaction							
byproduct, particularly in the manufacture and production of 1,2-dichloroethane (CASRN 107-06-2). EPA is not								
including this activity in the risk evaluation for 1,1,2-trichloroethane, but will address it in the risk evaluation for 1,2-								
dichloroethane and has	therefore removed the non-	incorporative activities condition of u	se from the final scope for					
1,1,2-trichloroethane.								

### 2.2.2 Activities Excluded from the Scope of the Risk Evaluation

As explained in the final rule for *Procedures for Chemical Risk Evaluation Under the Amended Toxic Substances Control Act* (82 FR 33726, July 20, 2017), TSCA Section 6(b)(4)(D) requires EPA to identify the hazards, exposures, conditions of use, and the PESS the Administrator expects to consider in a risk evaluation, suggesting that EPA may exclude certain activities that it determines to be conditions of use on a case-by-case basis (82 FR 33736, 33729; July 20, 2017). TSCA Section 3(4) also grants EPA discretion to determine the circumstances that are appropriately considered to be conditions of use for a particular chemical substance<sup>5</sup>. As a result, EPA does not plan to include in this scope or in the risk evaluation activities that the Agency does not consider to be conditions of use or for which EPA is exercising discretionary authority provided by TSCA Section 6(b)(4)(D).

No activities were excluded for 1,1,2-trichloroethane.

### 2.2.3 Production Volume

As reported to EPA during the 2016 CDR submission period and described here as a range to protect production volumes that were claimed as confidential business information (CBI), total production volume of 1,1,2-trichloroethane in 2015 was between 100 million and 250 million pounds (U.S. EPA, 2020a). EPA also uses pre-2015 CDR production volume information, as detailed in the *Proposed Designation of 1,1,2-trichloroethane (CASRN 79-00-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019d) and will include future production volume information from the 2020 CDR reporting period in the risk evaluation to support the exposure assessment.

#### 2.2.4 Overview of Conditions of Use and Lifecycle Diagram

Figure 2-11 provides the lifecycle diagram for 1,1,2-trichloroethane. The life cycle diagram is a graphical representation of the various life stages of the industrial, commercial and consumer use categories included within the scope of the risk evaluation. The information in the life cycle diagram is

<sup>&</sup>lt;sup>5</sup> *Chemical substance* means any organic or inorganic substance of a particular molecular identity, including any combination of such substances occurring in whole or in part as a result of a chemical reaction or occurring in nature, and any element or uncombined radical. Chemical substance does not include (1) any mixture; (2) any pesticide (as defined in the Federal Insecticide, Fungicide, and Rodenticide Act) when manufactured, processed, or distributed in commerce for use as a pesticide; (3) tobacco or any tobacco product; (4) any source material, special nuclear material, or byproduct material (as such terms are defined in the Atomic Energy Act of 1954 and regulations issued under such Act); (5) any article the sale of which is subject to the tax imposed by Section 4181 of the Internal Revenue Code of 1954 (determined without regard to any exemptions from such tax provided by Section 4182 or 4221 or any other provision of such Code), and; (6) any food, food additive, drug, cosmetic, or device (as such terms are defined in Section 201 of the Federal Food, Drug, and Cosmetic Act) when manufactured, processed, or distributed in commerce for use as a food, food additive, drug, cosmetic, or device (TSCA § 3(2)).

grouped according to the CDR processing codes and use categories (including functional use codes for industrial uses and product categories for industrial, commercial and consumer uses). Appendix E contains additional descriptions (*e.g.*, process descriptions, worker activities, process flow diagrams) for each manufacture, processing, distribution in commerce, use and disposal category.




Figure 2-11. 1,1,2-Trichloroethane Life Cycle Diagram

### 2.3 Exposures

For TSCA exposure assessments, EPA plans to analyze exposures and releases to the environment resulting from the conditions of use within the scope of the risk evaluation for 1,1,2-trichloroethane. Release pathways and routes will be described to characterize the relationship or connection between the conditions of use of the chemical and the exposure to human receptors, including PESS and environmental receptors. EPA plans to consider, where relevant, the duration, intensity (concentration), frequency and number of exposures in characterizing exposures to 1,1,2-trichloroethane.

### 2.3.1 Physical and Chemical Properties

Consideration of physical and chemical properties is essential for a thorough understanding or prediction of environmental fate (*i.e.*, transport and transformation) and the eventual environmental concentrations. It can also inform the hazard assessment. Table 2-3 summarizes the physical and chemical property values preliminarily selected for use in the risk evaluation from among the range of reported values collected as of June 2020. This table differs from that presented in the *Proposed Designation of 1,1,2-Trichloroethane (CASRN 79-00-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019d) and may be updated as EPA continues to evaluate and integrate additional information through systematic review methods. Figure 2-12 summarizes the distribution of reported values for eight physical and chemical properties routinely used in existing chemical risk evaluations. Appendix B presents summary statistics for reported physical and chemical property values that were extracted and evaluated as of June 2020 are presented in the supplemental file *Data Extraction and Data Evaluation Tables for Physical and Chemical Property Studies* (EPA-HQ-OPPT-2018-0421).

Property or Endpoint	Value <sup>a</sup>	Reference	Data Quality Rating
Molecular formula	C <sub>2</sub> H <sub>3</sub> Cl <sub>3</sub>	NA	NA
Molecular weight	133.41 g/mol	NA	NA
Physical state	Liquid	O'Neil et al. (2013)	High
Physical properties	Clear, colorless liquid, pleasant odor	<u>NLM (2018)</u>	High
Melting point	-36.5°C	Rumble (2018)	High
Boiling point	114°C	U.S. EPA (2019b)	High
Density	1.43216 g/cm <sup>3</sup> at 298.15 K	Sathyanarayana et al. (2006)	High
Vapor pressure	23 mm Hg at 25°C	NLM (2018)	High
Vapor density	4.21 g/L at 760 mm Hg	NLM (2018)	High
Water solubility	4590 mg/L at 25°C	NLM (2018)	High
Octanol/water partition coefficient (log Kow)	1.89 at 20°C	<u>NLM (2018)</u>	High
Henry's Law constant	$8.24 \times 10^{-4}$ atm·m <sup>3</sup> /mol at 25°C	<u>NLM (2018)</u>	High
Flash point	114°C	<u>RSC (2019)</u>	Medium
Auto flammability	Not available		

 Table 2-3. Physical and Chemical Properties of 1,1,2-Trichloroethane

Property or Endpoint	Property or Endpoint Value <sup>a</sup>		Data Quality Rating
Viscosity	1.49846 cP at 20-25°C	Elsevier (2019)	High
Refractive index	1.4711	<u>NLM (2018)</u>	High
Dielectric constant	7.29	Elsevier (2019)	High

<sup>a</sup> Measured unless otherwise noted.

NA = Not applicable

Figure 2-12 displays a summary of the data collected as of June 2020 for eight physical and chemical values routinely used in TSCA existing chemical risk evaluations. The box and whisker plots for each endpoint illustrate the mean (average, indicated by the blue diamond) and the 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup> (median), 75<sup>th</sup>, and 90<sup>th</sup> percentiles. All individual data points are indicated by black squares, and value preliminarily selected for use in the risk evaluation is overlaid (indicated by the orange circle) to provide context for where it lies within the distribution of the dataset. The number of unique primary data sources is indicated below each box and whisker plot. If multiple sources presented equivalent values and cited the same primary source, only one of those was included in the statistical calculations. As a result, the number of sources listed in Figure 2-12 may differ from the total number of data sources presented in Figure 2-2.




Figure 2-12. Box and Whisker Plots of Reported Physical and Chemical Property Values

### 2.3.2 Environmental Fate and Transport

Understanding of environmental fate and transport processes assists in the determination of the specific exposure pathways and potential human and environmental receptors that need to be assessed in the risk evaluation for 1,1,2-trichloroethane. EPA plans to use the environmental fate characteristics described in Appendix C to support the development of the risk evaluation for 1,1,2-trichloroethane. The values for the environmental fate properties may be updated as EPA evaluates and integrates additional information into the risk evaluation through systematic review methods.

### 2.3.3 Releases to the Environment

Releases to the environment from conditions of use are a component of potential exposure and may be derived from reported data that are obtained through direct measurement, calculations based on empirical data and/or assumptions and models.

A source of information that EPA plans to consider in evaluating exposure is data reported under the Toxics Release Inventory (TRI) program. EPA's TRI database contains information on chemical waste management activities that are disclosed by industrial and federal facilities, including quantities released into the environment (*i.e.*, to air, water, and disposed of to land), treated, burned for energy, recycled, or transferred off-site to other facilities for these purposes.

Under the Emergency Planning and Community Right-to-Know Act (EPCRA) Section 313, 1,1,2trichloroethane is a TRI-reportable substance effective January 01, 1987 (40 CFR 372.65). For TRI reporting<sup>6</sup>, facilities in covered sectors in the United States are required to disclose releases and other waste management activity quantities of 1,1,2-trichloroethane under the CASRN 79-00-5 if they manufacture (including import) or process more than 25,000 pounds or otherwise use more than 10,000 pounds of the chemical in a given year by July 1 of the following year. Table 2-4 provides production-related waste management data for 1,1,2-trichloroethane reported by facilities to the TRI program for reporting year 2018.<sup>7</sup> As shown in the table, 27 facilities reported a total of approximately 99 million pounds of 1,1,2-trichloroethane waste managed. Approximately 45 and 54 million pounds were reported as having been recycled or treated, respectively, and the majority of which occurred on site. These two waste management activities account for 99.8% of the total production-related waste quantities reported for 1,1,2-trichloroethane for 2018. Quantities of 1,1,2-trichloroethane in magnitude.

Year	Number of Facilities	Recycled (lbs)	Recovered for Energy (lbs)	Treated (lbs)	Released <sup>a,b,c</sup> -(lbs) <sup>a,b,e</sup> (lbs)	Total Production Related Waste (lbs)
2018	27	44,571,667	61,794	54,484,826	65,369	99,183,656
Data source: U	J.S. EPA (2019e	)				

<sup>a</sup> Terminology used in these columns may not match the more detailed data element names used in the TRI public data and analysis access points. <sup>b</sup> Does not include releases due to one-time event not associated with production such as remedial actions or earthquakes. <sup>c</sup> Counts all releases including release quantities transferred and release quantities disposed of by a receiving facility reporting to TRI.

Table 2-5 provides a summary of the quantities of 1,1,2-trichloroethane disposed of or otherwise released to the environment during 2018. Releases to air comprised 71% of the 65,369 pounds of total disposal and other releases of 1,1,2-trichloroethane. About 67% of these air emissions originated from point sources, with the remainder from fugitive sources. Land disposal accounted for roughly 28% of 1,1,2-trichloroethane releases, the vast majority of which took place on site to Resource Conservation and Recovery Act (RCRA) Subtitle C landfills. The remaining portion of the total releases came primarily from a transfer off site to a waste broker for disposal (502 pounds).

	Air R		eleases	Water Releases (lbs)	]	Land Dispos	al	Other	
	Number of Facilities	Stack Air Releases (lbs)	Fugitive Air Releases (lbs)		Class I Under- ground Injection (lbs)	RCRA Subtitle C Landfills (lbs)	All other Land Disposal <sup>a</sup> (lbs)	Releases <sup>a</sup> (lbs)	Total Releases <sup>b, c</sup> (lbs)
Totals	27	30,822	15,470		96	18,434	5	502	65 260
Totals	21	46,	291	40.54		18,535		302	65,369

Table 2-5. Summary of Releases of 1,1,2-Trichloroethane to the Environment During 2018

Data source: U.S. EPA (2019e)

<sup>a</sup> Terminology used in these columns may not match the more detailed data element names used in the TRI public data and analysis access points. <sup>b</sup> These release quantities do include releases due to one-time events not associated with production such as remedial actions or earthquakes.

Counts release quantities once at final disposition, accounting for transfers to other TRI reporting facilities that ultimately dispose of the chemical waste.

<sup>&</sup>lt;sup>6</sup> For TRI reporting criteria see <u>https://www.epa.gov/toxics-release-inventory-tri-program/basics-tri-reporting</u>

<sup>&</sup>lt;sup>7</sup> Reporting year 2018 is the most recent TRI data available. Data presented in Table 2-4 were queried using TRI Explorer and uses the 2018 National Analysis data set (released to the public in November 2019). This dataset includes revisions for the years 1988 to 2018 processed by EPA.

While production-related waste managed shown in Table 2-4 excludes any quantities reported as catastrophic or one-time releases (TRI Form R Section 8 data), release quantities shown in Table 2-5 include both production-related and non-production-related quantities. In the case of 1,1,2-trichloroethane, the total release quantities listed in Table 2-4 and Table 2-5 are the same, but for other TRI chemicals these respective release quantities may differ slightly and may further reflect differences in TRI calculation methods for reported release range estimates (U.S. EPA, 2017).

EPA plans to review these data in conducting the exposure assessment component of the risk evaluation for 1,1,2-trichloroethane.

### 2.3.4 Environmental Exposures

The manufacturing, processing, distribution, use and disposal of 1,1,2-trichloroethane can result in releases to the environment and exposure to aquatic and terrestrial receptors (biota). Environmental exposures to biota are informed by releases into the environment, overall persistence, degradation, and bioaccumulation, and partitioning across different media. Concentrations of chemical substances in biota provide evidence of exposure. EPA plans to review available environmental exposure data for 1,1,2-trichloroethane.

EPA plans to review available environmental monitoring data in the risk evaluation. EPA's Ambient Monitoring Technology Information Center Air Toxics database has identified 1,1,2-trichloroethane in air (U.S. EPA, 1990). USGS's Monitoring Data – National Water Quality Monitoring Council has identified 1,1,2-trichloroethane in ground water, sediment, soil, surface water and biota (USGS, 1991a, <u>b</u>, <u>c</u>, <u>d</u>, <u>e</u>, <u>f</u>, <u>g</u>).

Estimated background levels of 0.00039  $\mu$ g/m<sup>3</sup> were modeled from emissions data available in 2011 in which reported National Emissions Inventory (NEI) was 48 tons per year (<u>U.S. EPA, 2015c</u>). 1,1,2-Trichloroethane can form in landfills as its parent compound, 1,1,2,2-tetrachloroethane, is broken down. When released into the environment, the majority of 1,1,2-trichloroethane ends up in the air although some may enter groundwater (<u>ATSDR, 1989</u>).

### 2.3.5 Occupational Exposures

EPA plans to evaluate worker activities where there is a potential for exposure under the various conditions of use (manufacturing, processing, industrial/commercial uses, and disposal) described in Section 2.2. In addition, EPA plans to evaluate exposure to occupational non-users (ONUs), *i.e.*, workers who do not directly handle the chemical but perform work in an area where the chemical is present. EPA also plans to consider the effect(s) that engineering controls (EC) and/or personal protective equipment (PPE) have on occupational exposure levels as part of the risk evaluation.

Examples of worker activities associated with the conditions of use within the scope of the risk evaluation for 1,1,2-trichloroethane that EPA may analyze include, but are not limited to:

- Unloading and transferring 1,1,2-trichloroethane to and from storage containers and process vessels;
- Handling and disposing of waste containing 1,1,2-trichloroethane;
- Cleaning and maintaining equipment;
- Sampling chemicals, formulations, or products containing 1,1,2-trichloroethane for quality control;
- Repackaging chemicals, formulations, or products containing 1,1,2-trichloroethane;

• Performing other work activities in or near areas where 1,1,2-trichloroethane is used.

1,1,2-Trichloroethane has a vapor pressure of approximately 23 mmHg at 25°C (see Appendix B). Based on the chemical's high volatility, EPA anticipates that workers and ONUs will be exposed via the inhalation route. EPA plans to analyze inhalation exposure to vapor in occupational exposure scenarios where 1,1,2-trichloroethane is used and handled in open systems; the extent of exposure could vary from facility to facility depending on many factors including but not limited to EC, type of facility, and facility design. Based on the conditions of use presented in Section 2.2, EPA plans to also analyze the potential for inhalation exposure to mist in occupational exposure scenarios where the chemical is spray applied (*e.g.*, adhesives in the aerospace industry). EPA also plans to analyze worker exposure to liquids via the dermal route. EPA does not plan to analyze dermal exposure for ONUs because they do not directly handle 1,1,2-trichloroethane.

EPA generally does not evaluate occupational exposures through the oral route. Workers and ONUs may inadvertently ingest inhaled particles that deposit in the upper respiratory tract. In addition, workers may transfer chemicals from their hands to their mouths. The frequency and significance of this exposure route are dependent on several factors including the physical and chemical properties of the substance during worker activities, the visibility of the chemicals on the hands while working, workplace training and practices, and personal hygiene that is difficult to predict (<u>Cherrie et al., 2006</u>). EPA will consider the relevance of this exposure route on a case-by-case basis, taking into consideration the aforementioned factors and any reasonably available information, and may assess oral exposure for workers for certain COUs and worker activities where warranted.

The United States has several regulatory and non-regulatory exposure limits for 1,1,2-trichloroethane: The Occupational Safety and Health Administration (OSHA) permissible exposure limit (PEL) (<u>OSHA</u>, <u>2009</u>) of 10 ppm (or 45 mg/m<sup>3</sup>) over an 8-hour workday, time weighted average (TWA). This chemical also has a National Institute for Occupational Safety and Health (NIOSH) Recommended Exposure Limit (REL) (<u>NIOSH</u>, <u>2005</u>) of 10 ppm (45 mg/m<sup>3</sup>) TWA. The American Conference of Governmental Industrial Hygienists (ACGIH) sets the Threshold Limit Value (TLV) at 10 ppm TWA.

### 2.3.6 Consumer Exposures

No consumer conditions of use information for 1,1,2-trichloroethane were found in the 2012 or 2016 CDR. Public comments submitted by the American Coatings Association indicated that 1,1,2-trichloroethane is used in adhesives and sealants available for consumer use.

Consumers are primarily exposed via inhalation from use of adhesives (<u>ATSDR, 2010</u>). Consumers using or disposing of adhesives or sealants may be exposed to 1,1,2-trichloroethane through direct liquid contact which may lead to a dermal exposure, or through vapor emissions which may lead to inhalation exposure, given its volatility at room temperature. Bystanders present during the consumer use of adhesives and sealants or disposal of 1,1,2-trichloroethane may also be exposed to vapor emissions leading to an inhalation exposure. Bystanders are not expected to come in contact with liquid adhesives and sealant. Known adhesive products are supplied in tubes. Because the product will not be sprayapplied, mist generation is expected to be negligible.

Based on these potential sources and pathways of exposure, EPA plans to analyze inhalation and dermal routes of exposures to consumers, and inhalation routes of exposure for bystanders that may result from the conditions of use of 1,1,2-trichloroethane. EPA does not plan to evaluate consumer exposures to 1,1,2-trichloroethane via the oral route as the exposure is not expected.

#### 2.3.7 General Population Exposures

Releases of 1,1,2-trichloroethane from certain conditions of use, such as manufacturing, processing, or disposal, may result in general population exposures via drinking water ingestion, dermal contact, and inhalation from air releases. The main route of exposure for humans is via inhalation (<u>OECD, 2018</u>); (<u>ATSDR, 2010</u>). Small amounts of 1,1,2-trichloroethane have been detected in drinking water (<u>CalEPA, 2006</u>).

A biomonitoring survey found that more than 95 percent of blood samples collected from the general population were below the level of detection for 1,1,2-trichloroethane (<u>ATSDR, 2010</u>). Blood concentrations of 1,1,2-trichloroethane were below the level of detection in 2783 individuals who participated in the National Health and Nutrition Examination Survey (NHANES) 2011-2012 subsample of the U.S. population (<u>CDC, 2013</u>). The general population pathways in the scope of this evaluation are described in Sections 2.6.3 and 2.7.2.5.

### 2.4 Hazards (Effects)

#### 2.4.1 Environmental Hazards

EPA considered reasonably available information (*e.g.*, federal and international government chemical assessments) on 1,1,2-trichloroethane as well as public comments received on the *Proposed Designation of* 1,1,2-*trichloroethane* (*CASRN* 79-00-5) *as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019d) and draft scope for 1,1,2-trichloroethane (U.S. EPA, 2020c) to identify potential environmental hazards. During prioritization, EPA identified environmental hazard effects for aquatic and terrestrial organisms.

Since prioritization, EPA applied automated techniques during the data screening phase of systematic review to identify the following potential environmental hazards and related information that may be considered for the risk evaluation (as explained in Appendix A): developmental, mortality, musculoskeletal, reproductive, and respiratory (Figure 2-10). A summary of references identified during the screening step of systematic review is included in the interactive literature inventory trees (Figure 2-9). As EPA continues to evaluate reasonably available and relevant hazard information identified through systematic review, EPA may update the list of potential hazard effects to be analyzed in the risk evaluation.

#### 2.4.2 Human Health Hazards

EPA considered reasonably available information (*e.g.*, federal and international government chemical assessments) on 1,1,2-trichloroethane as well as public comments received on the *Proposed Designation* of 1,1,2-trichloroethane (CASRN 79-00-5) as a High-Priority Substance for Risk Evaluation (U.S. EPA, 2019d) and draft scope for 1,1,2-trichloroethane (U.S. EPA, 2020c) to identify potential human health hazards. During prioritization, EPA identified the following potential human health hazards and related information: acute, repeat dose, genetic, toxicokinetic, irritation/corrosion, cancer, immune and neurological effects.

Since prioritization, EPA applied automated techniques during the data screening phase of systematic review to identify the following additional potential human health hazards and related information that may be considered for the risk evaluation (as explained in Appendix A): ADME, PBPK, cardiovascular, developmental, endocrine, gastrointestinal, hepatic, mortality, musculoskeletal, nutritional and metabolic, ocular and sensory, renal, reproductive, respiratory and skin and connective tissue (Figure 2-10). A summary of references identified during the screening step of systematic review is included in

the interactive literature inventory trees (Figure 2-9). As EPA continues to evaluate reasonably available and relevant hazard information identified through systematic review, EPA may update the list of potential hazard effects to be analyzed in the risk evaluation.

### 2.5 Potentially Exposed or Susceptible Subpopulations

TSCA § 6(b)(4) requires EPA to determine whether a chemical substance presents an unreasonable risk to "a potentially exposed or susceptible subpopulation identified as relevant to the risk evaluation." TSCA § 3(12) states that "the term 'potentially exposed or susceptible subpopulation' means a group of individuals within the general population identified by the Administrator who, due to either greater susceptibility or greater exposure, may be at greater risk than the general population of adverse health effects from exposure to a chemical substance or mixture, such as infants, children, pregnant women, workers, or the elderly." General population is "the total of individuals inhabiting an area or making up a whole group" and refers here to the U.S. general population (U.S. EPA, 2011a).

EPA identified the following PESS based on CDR information and studies reporting developmental and reproductive effects: children, women of reproductive age (*e.g.*, pregnant women), workers, including ONUs and users, and consumers, including users and bystanders (U.S. EPA, 2019d). EPA plans to include these PESS in the risk evaluation. Following further evaluation of the reasonably available information, EPA may evaluate PESS in the general population as they relate to fence line communities.

In developing exposure scenarios, EPA plans to analyze available data to ascertain whether some human receptor groups may be exposed via exposure pathways that may be distinct to a particular subpopulation or life stage and whether some human receptor groups may have higher exposure via identified pathways of exposure due to unique characteristics (*e.g.*, activities, duration or location of exposure) when compared with the general population (U.S. EPA, 2006b). Likewise, EPA plans to evaluate available human health hazard information to ascertain whether some human receptor groups may have greater susceptibility than the general population to the chemical's hazard(s). Based on these analyses, EPA may update the list of PESS in the risk evaluation.

### 2.6 Conceptual Models


In this section, EPA presents the conceptual models describing the identified exposures (pathways and routes), receptors and hazards associated with the conditions of use of 1,1,2-trichloroethane. Pathways and routes of exposure associated with workers and ONU 's are described in Section 2.6.1, and pathways and routes of exposure associated with consumers are described in Section 2.6.2. Pathways and routes of exposure associated with environmental releases and wastes, including those pathways that are under the jurisdiction of other EPA-administered laws, are discussed and depicted in the conceptual model shown in Section 2.6.3.1. Pathways that are under the jurisdiction of other EPA-administered laws, are presented in the conceptual model shown in Section 2.6.3.2.

#### 2.6.1 Conceptual Model for Industrial and Commercial Activities and Uses

Figure 2-13 illustrates the conceptual model for the pathways of exposure from industrial and commercial activities and uses of 1,1,2-trichloroethane that EPA plans to include in the risk evaluation. There is potential for exposure to workers and ONUs via inhalation routes and exposure to workers via dermal routes. It is expected that inhalation exposure to vapors is the most likely exposure route. In addition, workers at waste management facilities may be exposed via inhalation or dermal routes from wastewater treatment, incineration or via other disposal methods. EPA plans to evaluate activities resulting in exposures associated with distribution in commerce (*e.g.*, loading, unloading) throughout the

various lifecycle stages and conditions of use (e.g., manufacturing, processing, industrial use, commercial use, and disposal) rather than a single distribution scenario.

For each condition of use identified in Table 2-2, a determination was made as to whether or not EPA plans to evaluate each combination of exposure pathway, route, and receptor in the risk evaluation. The results of that analysis along with the supporting rationale are presented in Appendix F.




### Figure 2-13. 1,1,2-Trichloroethane Conceptual Model for Industrial and Commercial Activities and Uses: Worker and Occupational Non-User Exposures and Hazards

The conceptual model presents the exposure pathways, exposure routes, and hazards to human receptors from industrial and commercial activities and uses of 1,1,2-trichloroethane.

#### 2.6.2 Conceptual Model for Consumer Activities and Uses

The conceptual model in Figure 2-14 presents the exposure pathways, exposure routes and hazards to human receptors from consumer activities and uses of 1,1,2-trichloroethane. EPA expects inhalation to be the primary route of exposure and plans to analyze inhalation exposures to 1,1,2-trichloroethane vapor for consumers and bystanders. There is potential for dermal exposures to 1,1,2-trichloroethane via direct contact with liquid during consumer uses. Bystanders are not expected to have direct dermal contact to 1,1,2-trichloroethane. EPA plans to analyze direct dermal contact with liquid 1,1,2-trichloroethane. EPA plans to analyze direct dermal contact with liquid 1,1,2-trichloroethane for consumers using adhesives and sealants products. Known adhesive products are supplied in tubes and similar packaging. Because the product will not be spray-applied, mist generation for consumers and bystanders is expected to be negligible. The supporting rationale for consumer pathways considered for 1,1,2-trichloroethane are included in Appendix G.



#### Figure 2-14. 1,1,2-Trichloroethane Conceptual Model for Consumer Activities and Uses: Consumer Exposures and Hazards


The conceptual model presents the exposure pathways, exposure routes, and hazards to human receptors from consumer activities and uses of 1,1,2-trichloroethane.

a) Receptors include PESS (see Section 2.5).

## 2.6.3 Conceptual Model for Environmental Releases and Wastes: Potential Exposures and Hazards (Regulatory Overlay)

In this section, EPA presents the conceptual models describing the identified exposures (pathways and routes from environmental releases and wastes) and hazards to general population and environmental receptors associated with the conditions of use of 1,1,2-trichloroethane within the scope of the risk evaluation. It also discusses those pathways that may be addressed pursuant to other Federal laws.

The conceptual model in Figure 2-15 presents the potential exposure pathways, exposure routes and hazards to general population and environmental receptors from releases and waste streams associated with industrial, commercial, and consumer uses of 1,1,2-trichloroethane. The conceptual model shows the overlays, labeled and shaded to depict the regulatory programs under EPA-administered statutes and associated pathways that EPA considered for the scope of the risk evaluation. The regulatory programs that cover these environmental release and waste pathways are further described in Section 2.6.3.1.



# Figure 2-15. 1,1,2-Trichloroethane Conceptual Model for Environmental Releases and Wastes: Environmental and General Population Exposures and Hazards (Regulatory Overlay)

The conceptual model presents the exposure pathways, exposure routes and hazards to human and environmental receptors from releases and wastes from industrial, commercial, and consumer uses of 1,1,2-trichloroethane including the environmental statutes covering those pathways.

- a) Industrial wastewater or liquid wastes may be treated on-site and then released to surface water (direct discharge), or pre-treated and released to Publicly Owned Treatment Works (POTW) (indirect discharge). For consumer uses, such wastes may be released directly to POTW. Drinking water will undergo further treatment in drinking water treatment plant. Ground water may also be a source of drinking water. Inhalation from drinking water may occur via showering.
- b) Receptors include PESS (see Section 2.5).

#### 2.6.3.1 Exposure Pathways and Risks Addressed by Other EPA Administered Statutes

In its TSCA Section 6(b) risk evaluations, EPA is coordinating action on certain exposure pathways and risks falling under the jurisdiction of other EPA-administered statutes or regulatory programs. More specifically, EPA is exercising its TSCA authorities to tailor the scope of its risk evaluations, rather than focusing on environmental exposure pathways addressed under other EPA-administered statutes or regulatory programs or risks that could be eliminated or reduced to a sufficient extent by actions taken under other EPA-administered laws. EPA considers this approach to be a reasonable exercise of the Agency's TSCA authorities, which include:

- TSCA Section 6(b)(4)(D): "The Administrator shall, not later than 6 months after the initiation of a risk evaluation, publish the scope of the risk evaluation to be conducted, including the hazards, exposures, conditions of use, and the potentially exposed or susceptible subpopulations the Administrator expects to consider..."
- TSCA Section 9(b)(1): "The Administrator shall coordinate actions taken under this chapter with actions taken under other Federal laws administered in whole or in part by the Administrator. If the Administrator determines that a risk to health or the environment associated with a chemical substance or mixture could be eliminated or reduced to a sufficient extent by actions taken under the authorities contained in such other Federal laws, the Administrator shall use such authorities to protect against such risk unless the Administrator determines, in the Administrator's discretion, that it is in the public interest to protect against such risk by actions taken under this chapter."
- TSCA Section 9(e): "...[I]f the Administrator obtains information related to exposures or releases of a chemical substance or mixture that may be prevented or reduced under another Federal law, including a law not administered by the Administrator, the Administrator shall make such information available to the relevant Federal agency or office of the Environmental Protection Agency."
- TSCA Section 2(c): "It is the intent of Congress that the Administrator shall carry out this chapter in a reasonable and prudent manner, and that the Administrator shall consider the environmental, economic, and social impact of any action the Administrator takes or proposes as provided under this chapter."
- TSCA section 18(d)(1): "Nothing in this chapter, nor any amendment made by the Frank R. Lautenberg Chemical Safety for the 21st Century Act, nor any rule, standard of performance, risk evaluation, or scientific assessment implemented pursuant to this chapter, shall affect the right of a State or a political subdivision of a State to adopt or enforce any rule, standard of performance, risk evaluation, scientific assessment, or any other protection for public health or the environment that— (i) is adopted or authorized under the authority of any other Federal law or adopted to satisfy or obtain authorization or approval under any other Federal law..."

These TSCA authorities supporting tailored risk evaluations and intra-agency referrals are described in more detail below:

#### TSCA Section 6(b)(4)(D)

TSCA Section 6(b)(4)(D) requires EPA, in developing the scope of a risk evaluation, to identify the hazards, exposures, conditions of use, and PESS the Agency "expects to consider" in a risk evaluation. This language suggests that EPA is not required to consider all conditions of use, hazards, or exposure

pathways in risk evaluations. As EPA explained in the "*Procedures for Chemical Risk Evaluation Under the Amended Toxic Substances Control Act*" (82 FR 33726, July 20, 2017) ("Risk Evaluation Rule"), "EPA may, on a case-by-case basis, exclude certain activities that EPA has determined to be conditions of use in order to focus its analytical efforts on those exposures that are likely to present the greatest concern, and consequently merit an unreasonable risk determination." 82 FR 33726, 33729 (July 20, 2017).

In the problem formulation documents for many of the first 10 chemicals undergoing risk evaluation, EPA applied the same authority and rationale to certain exposure pathways, explaining that "EPA is planning to exercise its discretion under TSCA 6(b)(4)(D) to focus its analytical efforts on exposures that are likely to present the greatest concern and consequently merit a risk evaluation under TSCA, by excluding, on a case-by-case basis, certain exposure pathways that fall under the jurisdiction of other EPA-administered statutes." The approach discussed in the Risk Evaluation Rule and applied in the problem formulation documents is informed by the legislative history of the amended TSCA, which supports the Agency's exercise of discretion to focus the risk evaluation on conditions of use that raise the greatest potential for risk, especially given that some conditions of use pose greater potential for exposure than others and the risks from many conditions of use are deemed negligible or already well controlled. See June 7, 2016 Cong. Rec., S3519-S3520. Consistent with the approach articulated in the problem formulation documents, and as described in more detail below, EPA is exercising its authority under TSCA to tailor the scope of exposures evaluated in TSCA risk evaluations, rather than focusing on environmental exposure pathways addressed under other EPA-administered, media-specific statutes and regulatory programs.

#### TSCA Section 9(b)(1)

In addition to TSCA Section 6(b)(4)(D), the Agency also has discretionary authority under the first sentence of TSCA Section 9(b)(1) to "coordinate actions taken under [TSCA] with actions taken under other Federal laws administered in whole or in part by the Administrator." This broad, freestanding authority provides for intra-agency coordination and cooperation on a range of "actions." In EPA's view, the phrase "actions taken under [TSCA]" in the first sentence of Section 9(b)(1) is reasonably read to encompass more than just risk management actions, and to include actions taken during risk evaluation as well. More specifically, the authority to coordinate intra-agency actions exists regardless of whether the Administrator has first made a definitive finding of risk, formally determined that such risk could be eliminated or reduced to a sufficient extent by actions taken under authorities in other EPA-administered Federal laws, and/or made any associated finding as to whether it is in the public interest to protect against such risk by actions taken under TSCA. TSCA Section 9(b)(1) therefore provides EPA authority to coordinate actions with other EPA offices without ever making a risk finding, or following an identification of risk. This includes coordination on tailoring the scope of TSCA risk evaluations to focus on areas of greatest concern rather than exposure pathways addressed by other EPA-administered statutes and regulatory programs, which does not involve a risk determination or public interest finding under TSCA Section 9(b)(2).

In a narrower application of the broad authority provided by the first sentence of TSCA Section 9(b)(1), the remaining provisions of Section 9(b)(1) provide EPA authority to identify risks and refer certain of those risks for action by other EPA offices. Under the second sentence of Section 9(b)(1), "[i]f the Administrator determines that a risk to health or the environment associated with a chemical substance or mixture could be eliminated or reduced to a sufficient extent by actions taken under the authorities contained in such other Federal laws, the Administrator shall use such authorities to protect against such

risk unless the Administrator determines, in the Administrator's discretion, that it is in the public interest to protect against such risk by actions taken under [TSCA]." Coordination of intra-agency action on risks under TSCA Section 9(b)(1) therefore entails both an identification of risk, and a referral of any risk that could be eliminated or reduced to a sufficient extent under other EPA-administered laws to the EPA office(s) responsible for implementing those laws (absent a finding that it is in the public interest to protect against the risk by actions taken under TSCA).

Risk may be identified by OPPT or another EPA office, and the form of the identification may vary. For instance, OPPT may find that one or more conditions of use for a chemical substance present(s) a risk to human or ecological receptors through specific exposure routes and/or pathways. This could involve a quantitative or qualitative assessment of risk based on reasonably available information (which might include, *e.g.*, findings or statements by other EPA offices or other federal agencies). Alternatively, risk could be identified by another EPA office. For example, another EPA office administering non-TSCA authorities may have sufficient monitoring or modeling data to indicate that a particular condition of use presents risk to certain human or ecological receptors, based on expected hazards and exposures. This risk finding could be informed by information made available to the relevant office under TSCA Section 9(e), which supports cooperative actions through coordinated information-sharing.

Following an identification of risk, EPA would determine if that risk could be eliminated or reduced to a sufficient extent by actions taken under authorities in other EPA-administered laws. If so, TSCA requires EPA to "use such authorities to protect against such risk," unless EPA determines that it is in the public interest to protect against that risk by actions taken under TSCA. In some instances, EPA may find that a risk could be sufficiently reduced or eliminated by future action taken under non-TSCA authority. This might include, *e.g.*, action taken under the authority of the Safe Drinking Water Act (SDWA) to address risk to the general population from a chemical substance in drinking water, particularly if the Office of Water has taken preliminary steps such as listing the subject chemical substance on the Contaminant Candidate List (CCL). This sort of risk finding and referral could occur during the risk evaluation process, thereby enabling EPA to use more a relevant and appropriate authority administered by another EPA office to protect against hazards or exposures to affected receptors.

Legislative history on TSCA Section 9(b)(1) supports both broad coordination on current intra-agency actions, and narrower coordination when risk is identified and referred to another EPA office for action. A Conference Report from the time of TSCA's passage explained that Section 9 is intended "to assure that overlapping or duplicative regulation is avoided while attempting to provide for the greatest possible measure of protection to health and the environment." S. Rep. No. 94-1302 at 84. See also H. Rep. No. 114-176 at 28 (stating that the 2016 TSCA amendments "reinforce TSCA's original purpose of filling gaps in Federal law," and citing new language in Section 9(b)(2) intended "to focus the Administrator's exercise of discretion regarding which statute to apply and to encourage decisions that avoid confusion, complication, and duplication"). Exercising TSCA Section 9(b)(1) authority to coordinate on tailoring TSCA risk evaluations is consistent with this expression of Congressional intent.

Legislative history also supports a reading of Section 9(b)(1) under which EPA coordinates intra-agency action, including information-sharing under TSCA Section 9(e), and the appropriately positioned EPA office is responsible for the identification of risk and actions to protect against such risks. See, *e.g.*, Senate Report 114-67, 2016 Cong. Rec. S3522 (under TSCA Section 9, "if the Administrator finds that disposal of a chemical substance may pose risks that could be prevented or reduced under the Solid Waste Disposal Act, the Administrator should ensure that the relevant office of EPA receives that

information"); H. Rep. No. 114-176 at 28, 2016 Cong. Rec. S3522 (under Section 9, "if the Administrator determines that a risk to health or the environment associated with disposal of a chemical substance could be eliminated or reduced to a sufficient extent under the Solid Waste Disposal Act, the Administrator should use those authorities to protect against the risk"). Legislative history on Section 9(b)(1) therefore supports coordination with and referral of action to other EPA offices, especially when statutes and associated regulatory programs administered by those offices could address exposure pathways or risks associated with conditions of use, hazards, and/or exposure pathways that may otherwise be within the scope of TSCA risk evaluations.

#### TSCA Sections 2(c) and 18(d)

Finally, TSCA Section 2(c) supports coordinated action on exposure pathways and risks addressed by other EPA-administered statutes and regulatory programs. Section 2(c) directs EPA to carry out TSCA in a "reasonable and prudent manner" and to consider "the environmental, economic, and social impact" of its actions under TSCA. Legislative history from around the time of TSCA's passage indicates that Congress intended EPA to consider the context and take into account the impacts of each action under TSCA. S. Rep. No. 94-698 at 14 ("the intent of Congress as stated in this subsection should guide each action the Administrator takes under other sections of the bill").

Section 18(d)(1) specifies that state actions adopted or authorized under any Federal law are not preempted by an order of no unreasonable risk issued pursuant to TSCA section 6(i)(1) or a rule to address unreasonable risk issued under TSCA section 6(a). Thus, even if a risk evaluation were to address exposures or risks that are otherwise addressed by other federal laws and, for example, implemented by states, the state laws implementing those federal requirements would not be preempted. In such a case, both the other federal and state laws, as well as any TSCA section 6(i)(1) order or TSCA section 6(a) rule, would apply to the same issue area. See also TSCA section 18(d)(1)(A)(iii). In legislative history on amended TSCA pertaining to section 18(d), Congress opined that "[t]his approach is appropriate for the considerable body of law regulating chemical releases to the environment, such as air and water quality, where the states have traditionally had a significant regulatory role and often have a uniquely local concern." Sen. Rep. 114-67 at 26.

EPA's careful consideration of whether other EPA-administered authorities are available, and more appropriate, for addressing certain exposures and risks is consistent with Congress' intent to maintain existing federal requirements and the state actions adopted to locally and more specifically implement those federal requirements, and to carry out TSCA in a reasonable and prudent manner. EPA believes it is both reasonable and prudent to tailor TSCA risk evaluations when other EPA offices have expertise and experience to address specific environmental media, rather than attempt to evaluate and regulate potential exposures and risks from those media under TSCA. This approach furthers Congressional direction and EPA aims to efficiently use Agency resources, avoid duplicating efforts taken pursuant to other Agency programs, and meet the statutory deadline for completing risk evaluations.

EPA-administered statutes and regulatory programs that address specific exposure pathways and/or risks are listed as follows:

#### Ambient Air Pathway

The Clean Air Act (CAA) contains a list of hazardous air pollutants (HAP) and provides EPA with the authority to add to that list pollutants that present, or may present, a threat of adverse human health effects or adverse environmental effects. For stationary source categories emitting HAP, the CAA

requires issuance of technology-based standards and, if necessary, additions or revisions to address developments in practices, processes, and control technologies, and to ensure the standards adequately protect public health and the environment. The CAA thereby provides EPA with comprehensive authority to regulate emissions to ambient air of any hazardous air pollutant. 1,1,2-Trichloroethane is a HAP. See 42 U.S.C. 7412. EPA has issued a number of technology-based standards for source categories that emit 1,1,2-trichloroethane to ambient air and, as appropriate, has reviewed, or is in the process of reviewing remaining risks. See 40 CFR part 63.

Emission pathways to ambient air from commercial and industrial stationary sources and associated inhalation exposure of the general population or terrestrial species in this TSCA evaluation from stationary source releases of 1,1,2-trichlorethane to ambient air are covered under the jurisdiction of the CAA. EPA's Office of Air and Radiation and Office of Pollution Prevention and Toxics will continue to work together to provide an understanding and analysis of the CAA regulatory analytical processes and to exchange information related to toxicity and occurrence data on chemicals undergoing risk evaluation under TSCA. As such, EPA does not plan to evaluate exposures to the general population from ambient air in the risk evaluation under TSCA. This regulatory coverage is represented by the red shading in Figure 2-15.

#### Drinking Water Pathway

EPA has regular analytical processes to identify and evaluate unregulated drinking water contaminants of potential regulatory concern for public water systems under the SDWA. In addition, the SDWA requires EPA to review and revise "as appropriate" existing drinking water regulations every 6 years.

EPA has promulgated National Primary Drinking Water Regulations (NPDWRs) under the Safe Drinking Water Act for 1,1,2-trichloroethane. EPA has set an enforceable Maximum Contaminant Level (MCL) as close as feasible to a health based, non-enforceable Maximum Contaminant Level Goal (MCLG). Public water systems are required to monitor for the regulated chemical based on a standardized monitoring schedule to ensure compliance with the MCL. The MCL for 1,1,2-trichloroethane in water is 0.005 mg/liter and the MCLG is 0.003 mg/liter (U.S. EPA, 1995).

The drinking water exposure pathway for 1,1,2-trichloroethane is currently addressed in the SDWA regulatory analytical process for public water systems. As such, EPA does not plan to evaluate exposures to the general population from drinking water exposure in the risk evaluation. This regulatory coverage is represented by the dark blue shading in Figure 2-15. EPA's Office of Water and Office of Pollution Prevention and Toxics will continue to work together to provide an understanding and analysis of the SDWA regulatory analytical processes and to exchange information related to toxicity and occurrence data on chemicals undergoing risk evaluation under TSCA.

#### Ambient Water Pathway

EPA has developed Clean Water Act (CWA) Section 304(a) recommended human health criteria for 122 chemicals and aquatic life criteria for 47 chemicals. A subset of these chemicals is identified as "priority pollutants" (103 human health and 27 aquatic life), including for 1,1,2-trichloroethane. The CWA requires that states adopt numeric criteria for priority pollutants for which EPA has published recommended criteria under Section 304(a), the discharge or presence of which in the affected waters could reasonably be expected to interfere with designated uses adopted the state. For pollutants with recommended human health criteria, EPA regulations require that state criteria contain sufficient parameters and constituents to protect designated uses. Once states adopt criteria as water quality standards, the CWA requires that National Pollutant Discharge Elimination System (NPDES) discharge

permits include effluent limits as stringent as necessary to meet standards CWA Section 301(b)(1)(C). This permit issuance process accounts for risk in accordance with the applicable ambient water exposure pathway (human health or aquatic life as applicable) for the designated water use.

EPA develops recommended water quality criteria under Section 304(a) of the CWA for pollutants in surface water that are protective of aquatic life or human health designated uses. EPA has developed recommended water quality criteria for protection of human health for 1,1,2-trichloroethane which are available for possible adoption into state water quality standards and are available for possible use by NPDES permitting authorities in deriving effluent limits to meet state narrative and/or numeric criteria. See, *e.g.*, 40 CFR part 423, Appendix A; 40 CFR 131.11(b)(1); 40 CFR 122.44(d)(1)(vi). As such, EPA does not plan to evaluate exposures to the general population from surface water in the risk evaluation under TSCA. This regulatory coverage is represented by the light blue shading in Figure 2-15. EPA's OW and OPPT will continue to work together to exchange information related to toxicity of chemicals undergoing risk evaluation under TSCA. EPA may update its CWA Section 304(a) water quality criteria for 1,1,2-trichloroethane in the future under the CWA.

EPA has not developed CWA Section 304(a) recommended water quality criteria for the protection of aquatic life for 1,1,2-trichloroethane, so there are no national recommended criteria for this use available for adoption into state water quality standards and available for use in NPDES permits. As such, EPA is evaluating exposures to aquatic species from surface water in the risk evaluation under TSCA. The Office of Water may issue CWA Section 304(a) aquatic life criteria for 1,1,2-trichloroethane in the future.

#### Onsite Releases to Land Pathway

The Comprehensive Environmental Response, Compensation, and Liability Act, otherwise known as CERCLA, provides broad authority under the statute (generally referred to as Superfund) to clean up uncontrolled or abandoned hazardous-waste sites as well as accidents, spills, and other releases of hazardous substances, pollutants and contaminants into the environment. Through CERCLA, EPA was given authority to seek out those parties potentially responsible for the release of hazardous substances and either have them clean up the release or compensate the Federal government for undertaking the response action.

CERCLA Section 101(14) defines "hazardous substance" by referencing other environmental statutes, including toxic pollutants listed under CWA Section 307(a); hazardous substances designated pursuant to CWA Section 311(b)(2)(A); hazardous air pollutants listed under CAA Section 112; imminently hazardous substances with respect to which EPA has taken action pursuant to TSCA Section 7; and hazardous wastes having characteristics identified under or listed pursuant to RCRA Section 3001. See 40 CFR 302.4. CERCLA Section 102(a) also authorizes EPA to promulgate regulations designating as hazardous substances those substances which, when released into the environment, may present substantial danger to the public health or welfare or the environment. EPA must also promulgate regulations establishing the quantity of any hazardous substance the release of which must be reported under Section 103. Section 103 requires persons in charge of vessels or facilities to report to the National Response Center if they have knowledge of a release of a hazardous substance above the reportable quantity threshold.

1,1,2-Trichloroethane is a hazardous substance under CERCLA. Releases of 1,1,2-trichloroethane in excess of 100 pounds within a 24-hour period must be reported (40 CFR 302.4, 302.6). The scope of this

EPA TSCA risk evaluation does not include on-site releases to the environment of 1,1,2-trichloroethane at Superfund sites and subsequent exposure of the general population or non-human species.

#### **Disposal and Soil Pathways**

1,1,2-Trichloroethane is included on the list of hazardous wastes pursuant to the Resource Conservation and Recovery Act (RCRA) 3001 (40 CFR §§ 261.33) as a listed waste on the U, F, and K lists (U227, F002, F024, F025, K019, K020, K095, K096). The general standard in Section RCRA 3004(a) for the technical criteria that govern the management (treatment, storage, and disposal) of hazardous waste are those "necessary to protect human health and the environment," RCRA 3004(a). The regulatory criteria for identifying "characteristic" hazardous wastes and for "listing" a waste as hazardous also relate solely to the potential risks to human health or the environment (40 CFR §§ 261.11, 261.21-261.24). RCRA statutory criteria for identifying hazardous wastes require EPA to "tak[e] into account toxicity, persistence, and degradability in nature, potential for accumulation in tissue, and other related factors such as flammability, corrosiveness, and other hazardous characteristics." Subtitle C controls cover not only hazardous wastes that are landfilled, but also hazardous wastes that are incinerated (subject to joint control under RCRA Subtitle C and the Clean Air Act (CAA) hazardous waste combustion Maximum Achievable Control Technology (MACT)) or injected into Underground Injection Control (UIC) Class I hazardous waste wells (subject to joint control under Subtitle C and the SDWA)<sup>8</sup>.

Emissions to ambient air from municipal and industrial waste incineration and energy recovery units that form combustion by-products from incineration treatment of 1,1,2-trichloroethane wastes may be subject to regulations, as would 1,1,2-trichloroethane burned for energy recovery.

TRI reporting in 2018 indicated 96 pounds released to underground injection to Class I hazardous waste wells. Environmental disposal of 1,1,2-trichloroethane injected into Class I hazardous well types fall under the jurisdiction of RCRA and SDWA and disposal of 1,1,2-trichloroethane via underground injection is not likely to result in environmental and general population exposures.

EPA has identified releases to land that go to RCRA Subtitle C hazardous waste landfills. Based on 2018 reporting, the majority of TRI land disposal includes Subtitle C landfills (18,434 pounds) with a much smaller amount transferred to "all other land disposal" both on-site and off-site (5 pounds reported in 2018). Design standards for Subtitle C landfills require double liner, double leachate collection and removal systems, leak detection system, run on, runoff, and wind dispersal controls, and a construction quality assurance program. They are also subject to closure and post-closure care requirements including installing and maintaining a final cover, continuing operation of the leachate collection and removal system until leachate is no longer detected, maintaining and monitoring the leak detection and groundwater monitoring system. Bulk liquids may not be disposed in Subtitle C landfills. Subtitle C landfill operators are required to implement an analysis and testing program to ensure adequate knowledge of waste being managed, and to train personnel on routine and emergency operations at the facility. Hazardous waste being disposed in Subtitle C landfills must also meet RCRA waste treatment standards before disposal. Given these controls, general population exposure in groundwater from Subtitle C landfill leachate is not expected to be a significant pathway. See 40 CFR part 264. As a result, EPA does not plan to evaluate on-site releases to land from RCRA Subtitle C hazardous waste landfills or exposures of the general population or terrestrial species from such releases in the TSCA evaluation. This regulatory coverage is represented by the pink shading in Figure 2-15.

<sup>&</sup>lt;sup>8</sup> This is not an exclusive list of Subtitle C authority, as it also covers, for example, disposal to surface impoundments, waste piles, and land treatment.


1,1,2-Trichloroethane is present in commercial and consumer products that may be disposed of in Municipal Solid Waste (MSW) landfills. On-site releases RCRA Subtitle D municipal solid waste landfills leading to exposures of the general population (including susceptible populations) or terrestrial species from such releases are expected to be minimal based on current TRI releases (*i.e.*, 5 lb in 2018) for 1,1,2- trichloroethane. While permitted and managed by the individual states, municipal solid waste (MSW) landfills are required by federal regulations to implement some of the same requirements as Subtitle C landfills. MSW landfills generally must have a liner system with leachate collection and conduct groundwater monitoring and corrective action when releases are detected. MSW landfills are also subject to closure and post-closure care requirements and must have financial assurance for funding of any needed corrective actions. MSW landfills have also been designed to allow for the small amounts of hazardous waste generated by households and very small quantity waste generators (less than 220 lb per month). Bulk liquids, such as free solvent, may not be disposed of at MSW landfills. See 40 CFR part 258. As a result, EPA does not plan to evaluate on-site releases to land from RCRA Subtitle D municipal solid waste (MSW) landfills or exposures of the general population or terrestrial species from such releases in the TSCA evaluation. This regulatory coverage is represented by the pink shading in Figure 2-15.

On-site releases to land from industrial non-hazardous and construction/demolition waste landfills may occur for 1,1,2-trichloroethane. Industrial non-hazardous and construction/demolition waste landfills are primarily regulated under authorized state regulatory programs. States must also implement limited federal regulatory requirements for siting, groundwater monitoring, and corrective action, and a prohibition on open dumping and disposal of bulk liquids. States may also establish additional requirements such as for liners, post-closure and financial assurance, but are not required to do so. See e.g., RCRA Section 3004(c), 4007; 40 CFR part 257. As a result, EPA is not evaluating on-site releases to land from industrial non-hazardous waste and construction/demolition waste landfills or associated exposures to the general population. As a result, EPA does not plan to evaluate on-site releases to land from industrial non-hazardous waste and construction/demolition waste landfills or associated exposures to the general population. This regulatory coverage is represented by the pink shading in Figure 2-15.

# 2.6.3.2 Conceptual Model for Environmental Releases and Wastes: Potential Exposures and Hazards

As described in Section 2.6.3.1, some pathways in the conceptual models are covered under the jurisdiction of other environmental statutes administered by EPA. The conceptual model depicted in Figure 2-16 presents the exposure pathways, exposure routes and hazards to environmental receptors from releases and wastes from industrial, commercial, and consumer uses of 1,1,2-trichloroethane that EPA plans to evaluate.

The diagram shown in Figure 2-16 includes releases from industrial, commercial and/or consumer uses to water/sediment; biosolids and soil, via direct and indirect discharges to water, that may lead to exposure to aquatic receptors. The supporting basis for environmental pathways considered for 1,1,2-trichloroethane are included in Appendix H.



### Figure 2-16. 1,1,2-Trichloroethane Conceptual Model for Environmental Releases and Wastes: Environmental Exposures and Hazards

The conceptual model presents the exposure pathways, exposure routes and hazards to environmental receptors from releases and wastes from industrial, commercial, and consumer uses of 1,1,2-trichloroethane that EPA plans to consider in the risk evaluation.

- a) Industrial wastewater or liquid wastes may be treated on-site and then released to surface water (direct discharge), or pre-treated and released to POTW (indirect discharge). For consumer uses, such wastes may be released directly to POTW.
- b) Receptors include PESS (see Section 2.5).

### 2.7 Analysis Plan

The analysis plan is based on EPA's knowledge of 1,1,2-trichloroethane resulting from the full-text screening of reasonably available information as described in Section 2.1. EPA encourages submission of additional existing data, such as full study reports or workplace monitoring from industry sources, that may be relevant to EPA's evaluation of conditions of use, exposures, hazards and PESS during risk evaluation. As discussed in the *Application of Systematic Review in TSCA Risk Evaluations* document (U.S. EPA, 2018), targeted supplemental searches during the analysis phase may be necessary to identify additional information (*e.g.*, commercial mixtures) for the risk evaluation of 1,1,2-trichloroethane. For any additional data needs identified during the risk evaluation, EPA may use the Agency's TSCA authorities under Sections 4, 8 or 11, as appropriate.

#### 2.7.1 Physical and Chemical Properties and Environmental Fate

EPA plans to analyze the physical and chemical properties and environmental fate and transport of 1,1,2-trichloroethane as follows:

- Review reasonably available measured or estimated physical and chemical properties and environmental fate endpoint data collected using systematic review procedures and, where reasonably available, environmental assessments conducted by other regulatory agencies. EPA plans to evaluate data and information collected through the systematic review methods and public comments about the physical and chemical properties (Appendix B) and fate endpoints (Appendix C), some of which appeared in the *Proposed Designation of 1,1,2-Trichloroethane* (*CASRN 79-00-5*) as a High-Priority Substance for Risk Evaluation (U.S. EPA, 2019d). All sources cited in EPA's analysis will be evaluated according to the procedures and metrics described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018). Where the systematic review process does not identify experimentally measured chemical property values of sufficiently high quality, testing will be requested under the TSCA Section 4 authority, or values will be estimated using chemical parameter estimation models as appropriate. Model-estimated fate properties will be reviewed for applicability and quality.
- 2) Using measured data and/or modeling, determine the influence of physical and chemical properties and environmental fate endpoints (*e.g.*, persistence, bioaccumulation, partitioning, transport) on exposure pathways and routes of exposure to human and environmental receptors.

Measured data and, where necessary, model predictions of physical and chemical properties and environmental fate endpoints will be used to characterize the persistence and movement of 1,1,2trichloroethane within and across environmental media. The fate endpoints of interest include volatilization, sorption to organic matter in sediments, water solubility, aqueous and atmospheric photolysis rates, aerobic and anaerobic biodegradation rates, and potential bioconcentration and bioaccumulation. These endpoints will be used in exposure calculations. **3)** Conduct a weight of the scientific evidence evaluation of physical and chemical properties and environmental fate data, including qualitative and quantitative sources of information. During risk evaluation, EPA plans to evaluate and integrate the physical and chemical properties and environmental fate evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018).

#### 2.7.2 Exposure

EPA plans to analyze exposure levels for indoor air, ambient air, surface water, sediment, and aquatic biota associated to exposure to 1,1,2-trichloroethane. Based on its physical and chemical properties, expected sources, and transport and transformation within the outdoor and indoor environment, 1,1,2-trichloroethane is more likely to be present in some of these media and less likely to be present in others. EPA has not yet determined the exposure levels in these media. Exposure level(s) can be characterized through a combination of reasonably available monitoring data and estimated exposure levels from modeling approaches. Exposure scenarios are combinations of sources (uses), exposure pathways, and exposed receptors. Draft exposure scenarios corresponding to various conditions of use for 1,1,2-trichloroethane are presented in Appendix F, Appendix G and Appendix H. EPA plans to analyze scenario-specific exposures.

#### 2.7.2.1 Environmental Releases

EPA plans to analyze releases to environmental media as follows:

1) Review reasonably available published literature and other reasonably available information on processes and activities associated with the conditions of use to analyze the types of releases and wastes generated.

EPA has reviewed some sources containing information on processes and activities resulting in releases, and the information found is described in Appendix E. EPA plans to continue to review data sources identified. Potential sources of environmental release data are summarized in Table 2-6 below:

Tuble 2 of Subgories and Sources of Entit officiental Release Data
U.S. EPA TRI Data
U.S. EPA Generic Scenarios
OECD Emission Scenario Documents
Discharge Monitoring Report (DMR) surface water discharge data for 1,1,2-trichloroethane from
NPDES-permitted facilities

2) Review reasonably available chemical-specific release data, including measured or estimated release data (e.g., data from risk assessments by other environmental agencies). EPA has reviewed key release data sources including the TRI, and the data from this source is summarized in Section 2.3.3. EPA will continue to consider additional reasonably available information and will evaluate it during development of the risk evaluation. EPA plans to match identified data to applicable conditions of use and identify data gaps where no data are found for particular conditions of use. EPA plans to attempt to address data gaps identified as described in steps 3 and 4 below by considering potential surrogate data and models.

Additionally, for conditions of use where no measured data on releases are reasonably available, EPA may use a variety of methods including release estimation approaches and assumptions in

the Chemical Screening Tool for Exposures and Environmental Releases (ChemSTEER) (U.S. EPA, 2015a).

**3**) Review reasonably available measured or estimated release data for surrogate chemicals that have similar uses and physical properties.

EPA plans to review literature sources identified and if surrogate data are found, these data will be matched to applicable conditions of use for potentially filling data gaps.

4) Review reasonably available data that may be used in developing, adapting or applying exposure models to the particular risk evaluation.

This item will be performed after completion of #2 and #3 above. EPA plans to evaluate relevant data to determine whether the data can be used to develop, adapt or apply models for specific conditions of use (and corresponding release scenarios). EPA has identified information from various EPA statutes and sources (including, for example, regulatory limits, reporting thresholds or disposal requirements) that may be relevant to consider for release estimation and environmental exposures. EPA plans to consider relevant regulatory requirements in estimating releases during risk evaluation.

5) Review and determine applicability of OECD Emission Scenario Documents (ESDs) and EPA Generic Scenarios to estimation of environmental releases.

EPA has identified potentially relevant OECD Emission Scenario Documents (ESDs) and EPA Generic Scenarios (GS) that correspond to some conditions of use; for example, the <u>September</u> 2011 ESD on the Chemical Industry (OECD, 2011) may be useful in estimating environmental releases. EPA plans to need to critically review these generic scenarios and ESDs to determine their applicability to the conditions of use assessed.

EPA Generic Scenarios are available at the following: <u>https://www.epa.gov/tsca-screening-tools/chemsteer-chemical-screening-tool-exposures-and-environmental-releases.</u>

OECD Emission Scenario Documents are available at the following: http://www.oecd.org/chemicalsafety/risk-assessment/emissionscenariodocuments.htm

If ESDs and GSs are not available, other methods may be considered. Additionally EPA may also perform supplemental targeted searches of peer-reviewed or gray literature for applicable models and associated parameters that EPA may use to estimate releases for certain conditions of use., for conditions of use where no measured data on releases are available, EPA may use a variety of methods including the application of default assumptions such as standard loss fractions associated with drum cleaning (3%) or single process vessel cleanout (1%).

#### 6) Map or group each condition of use to a release assessment scenario(s).

EPA has identified release scenarios and mapped (*i.e.*, grouped) them to relevant conditions of use as shown in Appendix F. EPA plans to refine the mapping/grouping of release scenarios based on factors (*e.g.*, process equipment and handling, magnitude of production volume used, and exposure/release sources) corresponding to conditions of use using reasonably available information identified. EPA may perform supplemental targeted searches of peer-reviewed or gray literature to better understand certain conditions of use to further develop release scenarios.

#### 7) Evaluate the weight of the scientific evidence of environmental release data.

During risk evaluation, EPA plans to evaluate and integrate the exposure evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018). EPA plans to integrate the data using systematic review methods to assemble the relevant data, evaluate the data for quality and relevance, including strengths and limitations, followed by synthesis and integration of the evidence.

#### 2.7.2.2 Environmental Exposures

EPA plans to analyze the following in developing its environmental exposure assessment of 1,1,2-trichloroethane:

1) Review available environmental and biological monitoring data for all media relevant to environmental exposure.

For 1,1,2-trichloroethane, environmental media which EPA plans to analyze are sediment and surface water.

2) Review reasonably available information on releases to determine how modeled estimates of concentrations near industrial point sources compare with reasonably available monitoring data.

EPA plans to analyze and consider reasonably available environmental exposure models that meet the scientific standards under TSCA Section 26(h) and that estimate surface water, sediment concentrations will be analyzed and considered alongside reasonably available surface water, and sediment monitoring data to characterize environmental exposures. Modeling approaches to estimate surface water concentrations, sediment concentrations may generally include the following inputs: direct release into surface water, or sediment, and indirect release into surface water, sediment, fate and transport (partitioning within media) and characteristics of the environment (*e.g.*, river flow, volume of lake, meteorological data).

3) Determine applicability of existing additional contextualizing information for any monitored data or modeled estimates during risk evaluation.

Monitoring data or modeled estimates will be reviewed to determine how use patterns have changed over recent years and will determine how representative environmental concentrations are of applicable use patterns. Studies which relate levels of 1,1,2-trichloroethane in the environment or biota with specific sources or groups of sources will also be evaluated.

4) Group each condition(s) of use to environmental assessment scenario(s).

EPA plans to refine and finalize exposure scenarios for environmental receptors by considering combinations of sources (use descriptors), exposure pathways including routes, and populations exposed. For 1,1,2-trichloroethane, the following are noteworthy considerations in constructing exposure scenarios for environmental receptors:

- Estimates of surface water concentrations, sediment concentrations near industrial point sources based on reasonably available monitoring data.
- Generally, consider the following modeling inputs: release into the media of interest, fate and transport and characteristics of the environment.
- Reasonably available biomonitoring data. Monitoring data could be used to compare with species or taxa-specific toxicological benchmarks.

- Applicability of existing additional contextualizing information for any monitored data or modeled estimates during risk evaluation. Review and characterize the spatial and temporal variability, to the extent that data are reasonably available, and characterize exposed aquatic populations.
- Weight of the scientific evidence of environmental occurrence data and modeled estimates
- 5) Evaluate the weight of the scientific evidence of environmental occurrence data and modeled estimates.

During risk evaluation, EPA plans to evaluate and integrate the exposure evidence identified in the literature inventory using the methods described in the using the methods described in the Application of Systematic Review in TSCA Risk Evaluation (U.S. EPA, 2018).

#### 2.7.2.3 Occupational Exposures

1,1,2-trichloroethane.

EPA plans to analyze both worker and occupational non-user exposures as follows:

1) Review reasonably available exposure monitoring data for specific condition(s) of use. EPA plans to review exposure data including workplace monitoring data collected by government agencies such as the OSHA and the NIOSH, and monitoring data found in published literature. These workplace monitoring data include personal exposure monitoring data (direct exposures) and area monitoring data (indirect exposures).

EPA plans to consider the influence of applicable regulatory limits and recommended exposure guidelines on occupational exposures in the occupational exposure assessment. The following are some data sources identified thus far:

#### Table 2-7. Potential Sources of Occupational Exposure Data

2019 ATSDR Toxicological Profile for 1,1,2-Trichloroethane
U.S. OSHA Chemical Exposure Health Data (CEHD) program data
U.S. NIOSH Health Hazard Evaluation (HHE) Program reports

- 2) Review reasonably available exposure data for surrogate chemicals that have uses, volatility and physical and chemical properties similar to 1,1,2-tricloroethane. EPA plans to review literature sources identified and if surrogate data are found, these data will be matched to applicable conditions of use for potentially filling data gaps. For several conditions of use, EPA believes data for other chlorinated solvents may serve as surrogates for
- 3) For conditions of use where data are limited or not reasonably available, review existing exposure models that may be applicable in estimating exposure levels.

EPA has identified potentially relevant OECD emission scenario documents (ESDs) and EPA generic scenarios (GSs) corresponding to some conditions of use. For example, the <u>April 2015</u> <u>ESD on the Use of Adhesives</u> (OECD, 2015) may be used to estimate occupational exposures. EPA plans to critically review these generic scenarios and ESDs to determine their applicability to the conditions of use assessed. EPA was not able to identify ESDs or GSs corresponding to several conditions of use, including the use of 1,1,2-trichloroethane as a laboratory chemical. EPA may conduct industry outreach or perform supplemental targeted searches of peer-reviewed

or gray literature to understand those conditions of use, which may inform identification of exposure scenarios. EPA may also need to perform targeted supplemental searches to identify applicable models that EPA may use to estimate exposures for certain conditions of use.

## 4) Review reasonably available data that may be used in developing, adapting or applying exposure models to a particular risk evaluation scenario.

This will be performed after #2 and #3 are completed, and based on information developed from #2 and #3, EPA plans to evaluate relevant data to determine whether the data can be used to develop, adapt, or apply models for specific conditions of use (and corresponding exposure scenarios). EPA may utilize existing, peer-reviewed exposure models developed by EPA, other government agencies, or reasonably available in the scientific literature, or EPA may elect to develop additional models to assess specific condition(s) of use. Inhalation exposure models may be simple box models or two-zone (near-field/far-field) models. In two-zone models, the near-field exposure represents potential inhalation exposures to Workers, and the far-field exposure represents potential inhalation exposures to ONUs.

#### 5) Consider and incorporate applicable EC and/or PPE into exposure scenarios.

EPA plans to review potentially relevant data sources on EC and PPE to determine their applicability and incorporation into exposure scenarios during risk evaluation. OSHA recommends employers utilize the hierarchy of controls to address hazardous exposures in the workplace. The hierarchy of controls strategy outlines, in descending order of priority, the use of elimination, substitution, engineering controls, administrative controls, and lastly personal protective equipment (PPE). EPA plans to assess worker exposure pre- and post-implementation of EC, using reasonably available information on control technologies and control effectiveness. For example, EPA may assess worker exposure in industrial use scenarios before and after implementation of local exhaust ventilation.

#### 6) Map or group each condition of use to occupational exposure assessment scenario(s). EPA has identified occupational exposure scenarios and mapped them to relevant conditions of use (see Appendix F). As presented in the fourth column in Table\_Apx F-1, EPA has completed an initial mapping of exposure scenarios to conditions of use. EPA plans to refine mapping or grouping of occupational exposure scenarios based on factors (*e.g.*, process equipment and handling, magnitude of production volume used, and exposure/release sources) corresponding to conditions of use as additional information is reviewed during risk evaluation. EPA may perform supplemental targeted searches of peer-reviewed or gray literature to better understand certain conditions of use to further develop exposure scenarios.

# 7) Evaluate the weight of the scientific evidence of occupational exposure data, which may include qualitative and quantitative sources of information.

During risk evaluation, EPA plans to evaluate and integrate the exposure evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018). EPA plans to rely on the weight of the scientific evidence when evaluating and integrating occupational data. EPA plans to integrate the data using systematic review methods to assemble the relevant data, evaluate the data for quality and relevance, including strengths and limitations, followed by synthesis and integration of the evidence.

#### **2.7.2.4 Consumer Exposures**

EPA plans to analyze both consumers using a consumer product and bystanders associated with the consumer using the product as follows:

#### 1) Group each condition of use to consumer exposure assessment scenario(s).

Refine and finalize exposure scenarios for consumers by considering sources, exposure pathways including routes, and exposed populations.

For 1,1,2-trichloroethane, the following are noteworthy considerations in constructing consumer exposure scenarios:

- Conditions of use and type of consumer product
- Duration, frequency and magnitude of exposure
- Weight fraction of chemical in products
- Amount of chemical used

### 2) Evaluate the relative potential of indoor exposure pathways based on reasonably available data.

Based on the physical and chemical properties of 1,1,2-trichloroethane and the consumer uses identified, inhalation of vapors and dermal absorption of liquid are expected to be important indoor exposure pathways for consumers. EPA plans to review all reasonably available information in developing the consumer exposure scenarios and evaluating the exposure pathways in indoor environments.

# 3) Review existing indoor exposure models that may be applicable in estimating indoor air exposures.

Indoor exposure models that estimate emissions from use of consumer products are available. These models generally consider physical and chemical properties (e.g., vapor pressure, molecular weight), product specific properties (e.g., weight fraction of the chemical in the product), use patterns (e.g., duration and frequency of use), user environment (e.g., room of use, ventilation rates), and receptor characteristics (e.g., exposure factors, activity patterns). The OPPT's Consumer Exposure Model (CEM) and other similar models can be used to estimate indoor air exposures from consumer products.

4) Review reasonably available empirical data that may be used in developing, adapting or applying exposure models to a particular risk evaluation scenario. For example, existing models developed for a chemical assessment may be applicable to another chemical assessment if model parameter data are reasonably available.

To the extent other organizations have already modeled a 1,1,2-trichloroethane consumer exposure scenario that is relevant to the OPPT's assessment, EPA plans to evaluate those modeled estimates. In addition, if other chemicals similar to 1,1,2-trichloroethane have been modeled for similar uses, those modeled estimates will also be evaluated. The underlying parameters and assumptions of the models will also be evaluated.

5) Review reasonably available consumer product-specific sources to determine how those exposure estimates compare with each other and with indoor monitoring data reporting 1,1,2-trichloroethane in specific media (*e.g.*, indoor air).

The availability of 1,1,2-trichloroethane concentration for various conditions of use will be evaluated. This data provides the source term for any subsequent indoor modeling.

# 6) Review reasonably available population- or subpopulation-specific exposure factors and activity patterns to determine if PESS need to be further refined.

For 1,1,2-trichloroethane, EPA plans to evaluate exposure scenarios that involve PESS and plans to consider age-specific behaviors, activity patterns and exposure factors unique to those subpopulations. For some exposure scenarios related to consumer uses, EPA plans to consider whether exposures for adults may different from those of children due to different activities (*e.g.*, children may mouth certain products) or exposure factors (*e.g.*, inhalation rates).

7) Evaluate the weight of the scientific evidence of consumer exposure estimates based on different approaches.

EPA plans to rely on the weight of the scientific evidence when evaluating and integrating data related to consumer exposure. The weight of the scientific evidence may include qualitative and quantitative sources of information. EPA plans to integrate the data using systematic review methods to assemble the relevant data, evaluate the data for quality and relevance, including strengths and limitations, followed by synthesis and integration of the evidence.

#### 2.7.2.5 General Population

EPA does not plan to analyze general population exposures, based on a review of exposure pathways as described in Section 2.6.3. EPA does not plan to include in the risk evaluation pathways under programs of other environmental statutes, administered by EPA, for which long-standing regulatory and analytical processes already exist.

#### 2.7.3 Hazards (Effects)

#### 2.7.3.1 Environmental Hazards

EPA plans to conduct an environmental hazard assessment of 1,1,2-trichloroethane as follows:

1) Review reasonably available environmental hazard data, including data from alternative test methods (*e.g.*, computational toxicology and bioinformatics; high-throughput screening methods; data on categories and read-across; *in vitro* studies).

EPA plans to analyze the hazards of 1,1,2-trichloroethane to aquatic organisms, including plants, invertebrates (*e.g.*, insects, arachnids, mollusks, crustaceans), and vertebrates (*e.g.*, mammals, birds, amphibians, fish, reptiles) across exposure durations and conditions if potential environmental hazards are identified through systematic review results and public comments. Additional types of environmental hazard information will also be considered (*e.g.*, analogue and read-across data) when characterizing the potential hazards of 1,1,2-trichloroethane to aquatic organisms.

EPA plans to evaluate environmental hazard data using the evaluation strategies laid out in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018). The study evaluation results will be documented in the risk evaluation phase and data from acceptable studies will be extracted and integrated in the risk evaluation process.

Mechanistic data may include analyses of alternative test data such as novel *in vitro* test methods and high throughput screening. The association between acute and chronic exposure scenarios to the agent and each health outcome will also be integrated. Study results will be extracted and presented in evidence tables or another appropriate format by organ/system.

#### 2) Derive hazard thresholds for aquatic organisms.

Depending on the robustness of the evaluated data for a particular organism or taxa (*e.g.*, aquatic invertebrates), environmental hazard values (*e.g.*, EC<sub>x</sub>. LC<sub>x</sub>, NOEC, LOEC) may be derived and used to further understand the hazard characteristics of 1,1,2-trichloroethane to aquatic species. Identified environmental hazard thresholds may be used to derive concentrations of concern (COC), based on endpoints that may affect populations of organisms or taxa analyzed.

#### 3) Evaluate the weight of the scientific evidence of environmental hazard data.

During risk evaluation, EPA plans to evaluate and integrate the environmental hazard evidence identified in the literature inventory using the methods described in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018).

4) Consider the route(s) of exposure, based on reasonably available monitoring and modeling data and other available approaches to integrate exposure and hazard assessments. EPA plans to consider aquatic (*e.g.*, water and sediment exposures) pathways in the 1,1,2-trichloroethane conceptual model. These organisms may be exposed to 1,1,2-trichloroethane via a number of environmental pathways (*e.g.*, surface water, sediment, diet).

#### 5) Consider a persistent, bioaccumulative, and toxic (PBT) assessment of 1,1,2trichloroethane.

EPA plans to consider the persistence, bioaccumulation, and toxic (PBT) potential of 1,1,2trichloroethane after reviewing relevant physical and chemical properties and exposure pathways. EPA plans to assess the reasonably available studies collected from the systematic review process relating to bioaccumulation and bioconcentration (*e.g.*, BAF, BCF) of 1,1,2trichloroethane. In addition, EPA plans to integrate traditional environmental hazard endpoint values (*e.g.*, LC<sub>50</sub>, LOEC) and exposure concentrations (*e.g.*, surface water concentrations, tissue concentrations) for 1,1,2-trichloroethane with the fate parameters (*e.g.*, BAF, BCF, BMF, TMF).

6) Conduct an environmental risk estimation and characterization of 1,1,2-trichloroethane. EPA plans to conduct a risk estimation and characterization of 1,1,2-trichloroethane to identify if there are risks to the aquatic environments from the measured and/or predicted concentrations of 1,1,2-trichloroethane in environmental media (*e.g.*, water, sediment). Risk quotients (RQs) may be derived by the application of hazard and exposure benchmarks to characterize environmental risk (U.S. EPA, 1998; Barnthouse et al., 1982). Analysis of risk for characterization includes a confidence statement in risk estimation which qualitative judgment describing the certainty of the risk estimate considering the strength the evidence scores for hazard and exposure and the limitations, and relevance.

#### 2.7.3.2 Human Health Hazards

EPA plans to analyze human health hazards as follows:

Review reasonably available human health hazard data, including data from alternative test methods (*e.g.*, computational toxicology and bioinformatics; high-throughput screening methods; data on categories and read-across; *in vitro* studies; systems biology).
 EPA plans to evaluate human health studies using the evaluation strategies laid out in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018) and updates to the epidemiological data quality criteria released with the first ten risk evaluations. The study evaluation results will be documented in the risk evaluation phase and data from acceptable studies will be extracted and integrated in the risk evaluation process.

Mechanistic data may include analyses of alternative test data such as novel *in vitro* test methods and high throughput screening. The association between acute and chronic exposure scenarios to the agent and each health outcome will also be integrated. Study results will be extracted and presented in evidence tables or another appropriate format by organ/system.

2) In evaluating reasonably available data, determine whether particular human receptor groups may have greater susceptibility to the chemical's hazard(s) than the general population.

Reasonably available human health hazard data will be evaluated to ascertain whether some human receptor groups may have greater susceptibility than the general population to 1,1,2-trichloroethane hazard(s). Susceptibility of particular populations or subpopulations to 1,1,2-trichloroethane will be determined by evaluating information on factors that influence susceptibility.

EPA has reviewed some sources containing hazard information associated with susceptible populations and lifestages such as pregnant women and infants. Pregnancy (*i.e.*, gestation) and childhood are potential susceptible lifestages for 1,1,2-trichloroethane exposure. EPA may quantify these differences in the risk evaluation following further evaluation of the reasonably available data and information.

# 3) Conduct hazard identification (the qualitative process of identifying non-cancer and cancer endpoints) and dose-response assessment (the quantitative relationship between hazard and exposure) for identified human health hazard endpoints.

Human health hazards from acute and chronic exposures will be identified by evaluating the human and animal data that meet the systematic review data quality criteria described in the *Application of Systematic Review in TSCA Risk Evaluation* (U.S. EPA, 2018). Hazards identified by studies meeting data quality criteria will be grouped by routes of exposure relevant to humans (*e.g.*, oral, dermal, inhalation) and by cancer and noncancer endpoints identified in Section 2.4.2.

Dose-response assessment will be performed in accordance with EPA guidance (U.S. EPA, 2012a, 2011b, 1994) developing points of departure (POD) for either margins of exposure (MOEs), cancer slope factors (CSFs), oral slope factors (OSFs), and/or inhalation unit risks (IURs). Dose-response analyses may be used if the data meet data quality criteria and if additional information on the identified hazard endpoints are not reasonably available or would not alter the analysis.

The cancer mode of action (MOA) analyses determine the relevancy of animal data to human risk and how data can be quantitatively evaluated. If cancer hazard is determined to be applicable

to 1,1,2-trichloroethane, EPA plans to evaluate information on genotoxicity and the MOA for all cancer endpoints to determine the appropriate approach for quantitative cancer assessment in accordance with the U.S. EPA Guidelines for Carcinogen Risk Assessment (U.S. EPA, 2005a). In accordance with EPA's Supplemental Guidance for Assessing Susceptibility from Early-life Exposures to Carcinogens (U.S. EPA, 2005b), EPA plans to determine whether age-dependent adjustment factors (ADAFs) are appropriate for 1,1,2-trichloroethane for specific conditions of use based upon potential exposures to children.

4) Derive points of departure (PODs) where appropriate; conduct benchmark dose modeling depending on the reasonably available data. Adjust the PODs as appropriate to conform (*e.g.*, adjust for duration of exposure) to the specific exposure scenarios evaluated. Hazard data will be evaluated to determine the type of dose-response modeling that is applicable. Where modeling is feasible, a set of dose-response models that are consistent with a variety of potentially underlying biological processes will be applied to empirically model the dose-response relationships in the range of the observed data consistent with EPA's *Benchmark Dose Technical Guidance Document* (U.S. EPA, 2012a). Where dose-response modeling is not feasible, NOAELs or LOAELs will be identified. Non-quantitative data will also be evaluated for contribution to weight of the scientific evidence or for evaluation of qualitative endpoints that are not appropriate for dose-response assessment.

EPA plans to evaluate whether the reasonably available PBPK and empirical kinetic models are adequate for route-to-route and interspecies extrapolation of the POD, or for extrapolation of the POD to standard exposure durations (*e.g.*, lifetime continuous exposure). If application of the PBPK model is not possible, oral PODs may be adjusted by  $BW^{3/4}$  scaling in accordance with U.S. EPA (2011b), and inhalation PODs may be adjusted by exposure duration and chemical properties in accordance with U.S. EPA (1994).

#### 5) Evaluate the weight of the scientific evidence of human health hazard data.

During risk evaluation, EPA plans to evaluate and integrate the human health hazard evidence identified in the literature inventory under acute and chronic exposure conditions using the methods described in the *Application of Systematic Review in TSCA Risk Evaluations* (U.S. EPA, 2018).

6) Consider the route(s) of exposure (*e.g.*, oral, inhalation, dermal), reasonably available route-to-route extrapolation approaches; biomonitoring data; and approaches to correlate internal and external exposures to integrate exposure and hazard assessment. At this stage of review, EPA believes there will be sufficient reasonably available data to conduct a dose-response analysis and/or benchmark dose modeling for the oral route of exposure. EPA plans to also evaluate any potential human health hazards following dermal and inhalation exposure to 1,1,2-trichloroethane, which could be important for worker, consumer and general population risk analysis. Reasonably available data will be assessed to determine whether or not a point of departure can be identified for the dermal and inhalation routes.

If sufficient reasonably available toxicity studies are not identified through the systematic review process to assess risks from inhalation or dermal exposure, then a route-to-route extrapolation may be needed. The preferred approach is to use a PBPK model (U.S. EPA, 2006a). Without an adequate PBPK model, considerations regarding the adequacy of data for route-to-route

extrapolation are described in *Methods for Derivation of Inhalation Reference Concentrations* and Application of Inhalation Dosimetry (U.S. EPA, 1994). EPA may use these considerations when determining whether to extrapolate from the oral to the inhalation route of exposure. Similar approaches for oral-to-dermal route extrapolation are described in EPA guidance document *Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part E, Supplemental Guidance for Dermal Risk Assessment)* (U.S. EPA, 2004).

If there are acceptable inhalation data after completion of systematic review, EPA may also consider extrapolating from the inhalation to the dermal route if first-pass metabolism through the liver via the oral route is expected because in that case, use of data from the oral route is not recommended (U.S. EPA, 1994). EPA may also consider inhalation-to-dermal route extrapolation if an inhalation toxicity study with a sensitive hazard endpoint is used to evaluate risks. Based on these considerations, EPA extrapolated from the inhalation to the dermal route for several of the first ten risk evaluations under amended TSCA, including methylene chloride (U.S. EPA, 2020d) and carbon tetrachloride (U.S. EPA, 2020b).

7) Conduct a human health risk estimation and characterization of 1,1,2-trichloroethane. Analysis of risk for characterization includes a confidence statement in risk estimation. This confidence statement is based on qualitative judgment describing the certainty of the risk estimate considering the strength of the evidence scores for hazard and exposure along with their limitations and relevance. The lowest confidence evaluation for either hazard or exposure will drive the overall confidence estimate.

#### 2.7.4 Summary of Risk Approaches for Characterization

Risk characterization is an integral component of the risk assessment process for both environmental and human health risks. EPA plans to derive the risk characterization in accordance with the EPA's *Risk Characterization Handbook* (U.S. EPA, 2000). As defined in the EPA's Risk Characterization Policy, "*the risk characterization integrates information from the preceding components of the risk evaluation and synthesizes an overall conclusion about risk that is complete, informative and useful for decision makers*" (U.S. EPA, 2000). Risk characterization is considered to be a conscious and deliberate process to bring all important considerations about risk, not only the likelihood of the risk but also the strengths and limitations of the assessment, and a description of how others have assessed the risk into an integrated picture.

The level of information contained in each risk characterization varies according to the type of assessment for which the characterization is written. Regardless of the level of complexity or information, the risk characterization for TSCA risk evaluations will be prepared in a manner that is transparent, clear, consistent, and reasonable (U.S. EPA, 2000) and consistent with the requirements of the *Procedures for Chemical Risk Evaluation Under the Amended Toxic Substances Control Act* (82 FR 33726, July 20, 2017). As discussed in 40 CFR 702.43, risk characterization has a number of considerations. This is the step where EPA integrates the hazard and exposure assessments into risk estimates for the identified populations (including any PESS) and ecological characterization does not consider costs or other nonrisk factors, and takes into account, "where relevant, the likely duration, intensity, frequency, and number of exposures under the condition(s) of use ....." The risk characterization also summarizes the following considerations: (1) uncertainty and variability in each step of the risk evaluation; (2) data quality, and any applicable assumptions used; (3) alternative

interpretations of data and analyses, where appropriate; and (4) any considerations for environmental risk evaluations, if necessary (*e.g.*, related to nature and magnitude of effects).

EPA plans to also be guided by EPA's Information Quality Guidelines (U.S. EPA, 2002) as it provides guidance for presenting risk information. Consistent with those guidelines, in the risk characterization, EPA plans to also identify: (1) Each population addressed by an estimate of applicable risk effects; (2) the expected risk or central estimate of risk for the PESS affected; (3) each appropriate upper-bound or lower bound estimate of risk; (4) each significant uncertainty identified in the process of the assessment of risk effects and the studies that would assist in resolving the uncertainty; and (5) peer reviewed studies known to the Agency that support, are directly relevant to, or fail to support any estimate of risk effects and the methodology used to reconcile inconsistencies in the scientific information.

### 2.8 Peer Review

Peer review will be conducted in accordance with EPA's regulatory procedures for chemical risk evaluations, including using EPA's Peer Review Handbook (U.S. EPA, 2015b) and other methods consistent with Section 26 of TSCA (see 40 CFR 702.45). As explained in the Risk Evaluation Rule, the purpose of peer review is for the independent review of the science underlying the risk assessment. Peer review will therefore address aspects of the underlying science as outlined in the charge to the peer review panel such as hazard assessment, assessment of dose-response, exposure assessment, and risk characterization. The draft risk evaluation for 1,1,2-trichloroethane will be peer reviewed.

### REFERENCES

- ATSDR (Agency for Toxic Substances and Disease Registry). (1989). Toxicological profile for 1,1,2trichloroethane. Atlanta, GA: U.S. Department of Health and Human Services, U.S. Public Health Service <u>https://www.atsdr.cdc.gov/ToxProfiles/tp148.pdf</u>
- ATSDR (Agency for Toxic Substances and Disease Registry). (2010). Addendum to the toxicological profile for 1,1,2-trichloroethane. Atlanta, GA: U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry, Division of Toxicology and Environmental Medicine.

https://www.atsdr.cdc.gov/toxprofiles/1\_1\_2\_trichloroethane\_addendum.pdf?id=1166&tid=155

- Barnthouse, LW; DeAngelis, DL; Gardner, RH; O'Neill, RV; Suter, GW; Vaughan, DS. (1982). Methodology for environmental risk analysis. (ORNL/TM-8167). Oak Ridge, TN: Oak Ridge National Laboratory.
- <u>CalEPA</u> (California Environmental Protection Agency). (2006). Public health goals for chemicals in drinking water: 1,1,2-Trichloroethane. California: California Environmental Protection Agency, Office of Environmental Health Hazard Assessment, Pesticide and Environmental Toxicology Branch. <u>https://oehha.ca.gov/media/downloads/water/chemicals/phg/phg112tca030306.pdf</u>
- <u>CDC</u> (Centers for Diseases Control and Prevention). (2013). National Health and Nutrition Examination Survey Data (NHANES) [Database]. Atlanta, GA: CDC National Center for Health Statistics. <u>https://www.cdc.gov/nchs/nhanes/index.htm</u>
- <u>Cherrie, JW; Semple, S; Christopher, Y; Saleem, A; Hughson, GW; Philips, A.</u> (2006). How important is inadvertent ingestion of hazardous substances at work? Ann Occup Hyg 50: 693-704. <u>http://dx.doi.org/10.1093/annhyg/mel035</u>
- ECHA (European Chemicals Agency). (2019). Registration dossier: 1,1,2-Trichloroethane (CASRN 79-00-5): Phototransformation in air: 001 Supporting | Experimental result. <u>https://echa.europa.eu/da/registration-dossier/-/registered-</u> <u>dossier/1878/5/2/2/?documentUUID=0bce661d-13eb-40a0-8643-afa0c3e17484</u>
- Elsevier. (2019). Reaxys: physical-chemical property data for 1,1,2-trichloroethane. CAS Registry Number: 79-00-5. Available online
- Gan, D; Dupont, R. (1989). Hazard waste hazard materials (pp. 363-383).
- Gorilla Glue Company. (2018). Safety data sheet-Gorilla Weld-Resin.

https://www.gorillatough.com/wp-content/uploads/Gorilla-Weld-Resin-v1.2.pdf

- Gorilla Glue Company. (2020). GorillaWeld. https://www.gorillatough.com/product/gorillaweld/
- <u>IARC</u> (International Agency for Research on Cancer). (1991). IARC monographs on the evaluation of carcinogenic risks to humans: Occupational exposures in insecticide application, and some pesticides [IARC Monograph]. In IARC monographs on the evaluation of carcinogenic risks to humans: Occupational exposures in insecticide application, and some pesticides. Lyon, France: World Health Organization.
- <u>NIOSH</u> (National Institute for Occupational Safety and Health). (2005). NIOSH pocket guide to chemical hazards. Index of chemical abstracts service registry numbers (CAS No.). <u>https://www.cdc.gov/niosh/npg/npgdcas.html</u>
- <u>NITE</u> (National Institute of Technology and Evaluation). (2010). Japan CHEmicals Collaborative Knowledge database (J-CHECK). CASRN: 79-00-5. Available online at <u>https://www.nite.go.jp/chem/jcheck/detail.action?cno=79-00-5&mno=2-</u>0055&request\_locale=en

- NLM (National Institutes of Health, National Library of Medicine). (2018). PubChem: Hazardous Substance Data Bank: 1,1,2-Trichloroethane, 79-00-5. Available online at https://pubchem.ncbi.nlm.nih.gov/compound/6574#source=HSDB
- O'Neil, MJ; Heckelman, PE; Dobbelaar, PH; Roman, KJ; Kenney, CM; Karaffa, LS. (2013). The Merck Index: An Encyclopedia of Chemicals, Drugs, and Biologicals (15th ed.). Cambridge, U.K.: The Royal Society of Chemistry.
- OECD (Organisation for Economic Co-operation and Development). (2000). SIDS Initial Assessment Report: 1,1,2 Trichloroethane. Tokyo, Japan. http://www.inchem.org/documents/sids/sids/79005.pdf
- OECD (Organisation for Economic Co-operation and Development). (2011). Emission scenario document on the chemical industry. (JT03307750). Paris, France. http://www.oecd.org/env/ehs/risk-assessment/48774702.pdf
- OECD (Organisation for Economic Co-operation and Development). (2015). Emission scenario document on the use of adhesives. (JT03373626). Paris, France. <u>http://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=ENV/JM/MONO(2015</u> )4&doclanguage=en
- <u>OECD</u> (Organisation for Economic Co-operation and Development). (2018). OECD Monitoring Database [Database].
- OSHA (Occupational Safety & Health Administration). (2009). Occupational Safety and Health Administration (OSHA) permissible exposure limits (PELs). https://www.osha.gov/dsg/annotated-pels/tablez-1.html
- Rathbun, RE. (1998). Transport, behavior, and fate of volatile organic compounds in streams (pp. 1-151). (U.S. Geological Survey Professional Paper 1589). U.S. Geological Survey.
- Reed, NR; Reed, W; Beltran, L; Babapour, R; Hsieh, DPH. (1988). Health Risk Assessment of 1,1,2-Trichloroethane (1,1,2-TCA) in California Drinking Water.
- <u>RSC</u> (Royal Society of Chemistry). (2019). ChemSpider: 1,1,2-Trichloroethane. Available online at <u>http://www.chemspider.com/Chemical-Structure.6326.html?rid=2a2c9061-8bb6-4138-93e0-67641f978a2c</u>
- Rumble, JR. (2018). 1,1,2-Trichloroethane [Type of Work]. Boca Raton, FL: CRC Press. Taylor & Francis Group.
- Sathyanarayana, B; Jyostna, TS; Satyanarayana, N. (2006). Acoustic studies of binary mixtures of Nmethylacetamide with some chloroethanes and chloroethenes at 308.15 K. Indian J Pure Appl Phys 44: 587-591.
- <u>SYKE</u> (The Finnish Environment Institute). (2018). Data bank of environmental properties of chemicals 1,1,2-Trichloroethane, CAS RN: 79-00-5. The Finnish Environment Institute. <u>http://wwwp.ymparisto.fi/scripts/Kemrek/Kemrek\_uk.asp?Method=MAKECHEMdetailsform&t</u> xtChemId=298
- Tabak, HH; Quave, SA; Mashni, CI; Barth, EF. (1981). Biodegradability studies with organic priority pollutant compounds. 53: 1503-1518.
- <u>ThermoFisher Scientific.</u> (2018). Safety data sheet: 1,1,2-Trichloroethane. Revision no. 3. Revision date 1/18/2018.

https://www.fishersci.com/store/msds?partNumber=AC139430010&productDescription=1%2C1 %2C2-

 $\underline{\text{TRICHLOROETHANE\%2C+9+1LT\&vendorId=VN00032119\&countryCode=US\&language=e}\underline{n}$ 

- <u>Thomas, R; Byrne, M; Gilbert, D; Goyer, M; Wood, M.</u> (1982). An exposure and risk assessment for trichloroethanes. (EPA-440/4-85-018; PB85-220598). Washington, DC: Office of Water Regulations and Standards, US Environmental Protection Agency.
- Tomer, A; Kane, J. (2015). The great port mismatch. U.S. goods trade and international transportation. The Global Cities Initiative. A joint project of Brookings and JPMorgan Chase. https://www.brookings.edu/wp-content/uploads/2015/06/brgkssrvygcifreightnetworks.pdf
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (1990). EPA Ambient Monitoring Technology Information Center (AMTIC): Air toxics data [Database]. U.S. EPA (U.S. Environmental Protection Agency). <u>https://www3.epa.gov/ttnamti1/toxdat.html</u>
- U.S. EPA (U.S. Environmental Protection Agency). (1994). Methods for derivation of inhalation reference concentrations and application of inhalation dosimetry [EPA Report]. (EPA/600/8-90/066F). Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office.

https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=71993&CFID=51174829&CFTOKEN=2 5006317

U.S. EPA (U.S. Environmental Protection Agency). (1995). National Primary Drinking Water Regulations: Contaminant specific fact sheets, inorganic chemicals, consumer version. (811F95002C).

https://nepis.epa.gov/Exe/ZyNET.exe/20001RYK.TXT?ZyActionD=ZyDocument&Client=EPA &Index=1995+Thru+1999&Docs=&Query=&Time=&EndTime=&SearchMethod=1&TocRestri ct=n&Toc=&TocEntry=&QField=&QFieldYear=&QFieldMonth=&QFieldDay=&IntQFieldOp =0&ExtQFieldOp=0&XmlQuery=&File=D%3A%5Czyfiles%5CIndex%20Data%5C95thru99% 5CTxt%5C00000001%5C20001RYK.txt&User=ANONYMOUS&Password=anonymous&Sort Method=h%7C-

<u>&MaximumDocuments=1&FuzzyDegree=0&ImageQuality=r75g8/r75g8/r150y150g16/i425&D</u> isplay=hpfr&DefSeekPage=x&SearchBack=ZyActionL&Back=ZyActionS&BackDesc=Results %20page&MaximumPages=1&ZyEntry=1&SeekPage=x&ZyPURL

- U.S. EPA (U.S. Environmental Protection Agency). (1998). Guidelines for ecological risk assessment [EPA Report]. (EPA/630/R-95/002F). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. https://www.epa.gov/risk/guidelines-ecological-risk-assessment
- U.S. EPA (U.S. Environmental Protection Agency). (2000). Science policy council handbook: Risk characterization handbook. (EPA/100/B-00/002). Washington, D.C.: U.S. Environmental Protection Agency, Science Policy Council. <u>https://www.epa.gov/risk/risk-characterization-handbook</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2002). Guidelines for ensuring and maximizing the quality, objectivity, utility, and integrity of information disseminated by the Environmental Protection Agency. (EPA/260/R-02/008). Washington, DC: U.S. Environmental Protection Agency, Office of Environmental Information. <u>https://www.epa.gov/sites/production/files/2017-03/documents/epa-info-quality-guidelines.pdf</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2004). Risk Assessment Guidance for Superfund (RAGS), Volume I: Human health evaluation manual, (Part E: Supplemental guidance for dermal risk assessment). (EPA/540/R/99/005). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <u>http://www.epa.gov/oswer/riskassessment/ragse/index.htm</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2005a). Guidelines for carcinogen risk assessment [EPA Report]. (EPA/630/P-03/001B). Washington, DC: U.S. Environmental Protection Agency,

Risk Assessment Forum. <u>https://www.epa.gov/sites/production/files/2013-09/documents/cancer\_guidelines\_final\_3-25-05.pdf</u>

- U.S. EPA. (U.S. Environmental Protection Agency). (2005b). Supplemental guidance for assessing susceptibility from early-life exposure to carcinogens [EPA Report]. (EPA/630/R-03/003F). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. https://www3.epa.gov/airtoxics/childrens\_supplement\_final.pdf
- U.S. EPA. (U.S. Environmental Protection Agency). (2006a). Approaches for the application of physiologically based pharmacokinetic (PBPK) models and supporting data in risk assessment (Final Report) [EPA Report] (pp. 1-123). (EPA/600/R-05/043F). Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. <u>http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=157668</u>
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2006b). A framework for assessing health risk of environmental exposures to children (pp. 1-145). (EPA/600/R-05/093F). Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. <u>http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=158363</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2011a). Exposure factors handbook: 2011 edition [EPA Report]. (EPA/600/R-090/052F). Washington, DC. http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=236252
- U.S. EPA (U.S. Environmental Protection Agency). (2011b). Recommended use of body weight 3/4 as the default method in derivation of the oral reference dose (pp. 1-50). (EPA/100/R-11/0001). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum, Office of the Science Advisor. <u>https://www.epa.gov/sites/production/files/2013-09/documents/recommended-use-of-bw34.pdf</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2012a). Benchmark dose technical guidance. (EPA/100/R-12/001). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <u>https://www.epa.gov/risk/benchmark-dose-technical-guidance</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2012b). Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.11. Washington, DC. Retreived from <u>https://www.epa.gov/tsca-screening-tools/epi-suitetm-estimation-program-interface</u>
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2015a). ChemSTEER user guide Chemical screening tool for exposures and environmental releases. Washington, DC. https://www.epa.gov/sites/production/files/2015-05/documents/user\_guide.pdf
- U.S. EPA (U.S. Environmental Protection Agency). (2015b). Peer review handbook [EPA Report] (4th ed.). (EPA/100/B-15/001). Washington, DC: U.S. Environmental Protection Agency, Science Policy Council. <u>https://www.epa.gov/osa/peer-review-handbook-4th-edition-2015</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2015c). Technical support document, EPA's 2011 National-scale Air Toxics Assessment, 2011 NATA TSD. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air and Radiation, Office of Air Quality Planning and Standards. <u>https://www.epa.gov/sites/production/files/2015-12/documents/2011-nata-tsd.pdf</u>
- <u>U.S. EPA</u> (U.S. Environmental Protection Agency). (2017). Toxics Release Inventory (TRI), reporting year 2015. <u>https://www.epa.gov/toxics-release-inventory-tri-program/tri-data-and-tools</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2018). Application of systematic review in TSCA risk evaluations. (740-P1-8001). Washington, DC: U.S. Environmental Protection Agency, Office of Chemical Safety and Pollution Prevention. <u>https://www.epa.gov/sites/production/files/2018-</u>06/documents/final\_application\_of\_sr\_in\_tsca\_05-31-18.pdf

- U.S. EPA (U.S. Environmental Protection Agency). (2019a). Chemical data reporting (2012 and 2016 CBI CDR database). U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics.
- U.S. EPA (U.S. Environmental Protection Agency). (2019b). Chemistry Dashboard Information for 1,1,2-Trichloroethane. 79-00-5. Available online at

https://comptox.epa.gov/dashboard/dsstoxdb/results?search=DTXSID5021380

- U.S. EPA (U.S. Environmental Protection Agency). (2019c). CPCat: Chemical and product categories. https://actor.epa.gov/cpcat/faces/home.xhtml
- U.S. EPA (U.S. Environmental Protection Agency). (2019d). Proposed designation of 1,1,2trichloroethane (casrn 79-00-5) as a high-priority substance for risk evaluation. <u>https://www.epa.gov/sites/production/files/2019-08/documents/112-trichloroethane\_79-00-5\_high-priority\_proposeddesignation\_082319.pdf</u>
- U.S. EPA (U.S. Environmental Protection Agency). (2019e). TRI Explorer (2018 dataset released November 2019). Washington, DC: U.S. Environmental Protection Agency. https://enviro.epa.gov/triexplorer/
- U.S. EPA (U.S. Environmental Protection Agency). (2020a). Chemical data reporting (2012 and 2016 Public CDR database). Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics. ChemView: July 2020. <u>https://chemview.epa.gov/chemview</u>
- U.S. EPA. (U.S. Environmental Protection Agency). (2020b). Draft risk evaluation for carbon tetrachloride (methane, tetrachloro-); CASRN: 56-23-5 (pp. 1-301). (EPA-740-R1-8014). Office of Chemical Safety and Pollution Prevention, U.S. Environmental Protection Agency. https://nepis.epa.gov/Exe/ZyPDF.cgi/P100YHUW.PDF?Dockey=P100YHUW.PDF
- U.S. EPA. (U.S. Environmental Protection Agency). (2020c). Draft Scope of the risk evaluation for 1,1,2-Trichloroethane CASRN 79-00-5 [EPA Report]. (EPA-740-D-20-003). Washington, DC. <u>https://www.epa.gov/sites/production/files/2020-04/documents/casrn-79-00-5\_112-</u> <u>trichloroethane\_draft\_scope.pdf</u>
- U.S. EPA. (U.S. Environmental Protection Agency). (2020d). Risk evaluation for methylene chloride (dichloromethane, dcm); CASRN: 75-09-2 (pp. 1-753). (EPA-740-R1-8010). Office of Chemical Safety and Pollution Prevention, U.S. Environmental Protection Agency. <u>https://www.epa.gov/sites/production/files/2020-</u>06/documents/1\_mecl\_risk\_evaluation\_final.pdf
- <u>USGS</u> (U.S. Geological Survey). (1991a). USGS Monitoring Data: National Water Quality Monitoring Council [Database]. <u>http://www.waterqualitydata.us/portal/</u>
- USGS (U.S. Geological Survey). (1991b). USGS Monitoring Data: National Water Quality Monitoring Council - Air. https://www.waterquality.data.us/portal/#siteType=Aggregate%20groundwater%20use&sample

https://www.waterqualitydata.us/portal/#siteType=Aggregate%20groundwater%20use&sample Media=Water&mimeType=csv&dataProfile=activityAll

<u>USGS</u> (U.S. Geological Survey). (1991c). USGS Monitoring Data: National Water Quality Monitoring Council - Groundwater. https://www.waterqualitydata.us/portal/#siteType=Aggregate%20groundwater%20use&sample

<u>Media=Water&mimeType=csv&dataProfile=activityAll</u>

<u>USGS</u> (U.S. Geological Survey). (1991d). USGS Monitoring Data: National Water Quality Monitoring Council - Sediment.

https://www.waterqualitydata.us/portal/#sampleMedia=Sediment&mimeType=csv

<u>USGS</u> (U.S. Geological Survey). (1991e). USGS Monitoring Data: National Water Quality Monitoring Council - Soil. <u>https://www.waterqualitydata.us/portal/#sampleMedia=Soil&mimeType=csv</u>

- <u>USGS</u> (U.S. Geological Survey). (1991f). USGS Monitoring Data: National Water Quality Monitoring Council - Surface Water. <u>https://www.waterqualitydata.us/portal/#siteType=Aggregate%20surface-water-</u> use&sampleMedia=Water&mimeType=csv
- <u>USGS</u> (U.S. Geological Survey). (1991g). USGS Monitoring Data: National Water Quality Monitoring Council - Tissue.

https://www.waterqualitydata.us/portal/#sampleMedia=Tissue&mimeType=csv

<u>Verschueren, K.</u> (2001). Handbook of Environmental Data on Organic Chemicals (4th ed.). New York, NY: John Wiley & Sons, Incorporated.

## APPENDICES

# Appendix A ABBREVIATED METHODS FOR SEARCHING AND SCREENING

## A.1 Literature Search of Publicly Available Databases

### A.1.1 Search Term Genesis and Chemical Verification

To develop the chemical terms for the subsequent literature search for 1,1,2-trichloroethane, several online sources were queried.

- California Department of Pesticide Regulation: <u>https://www.cdpr.ca.gov/docs/chemical/monster2.htm</u>
- USEPA Chemistry Dashboard: https://comptox.epa.gov/dashboard
- University of Hertfordshire PPDB: Pesticide Properties DataBase: <u>https://sitem.herts.ac.uk/aeru/ppdb/en/search.htm</u>
- USEPA Reregistration Eligibility Decision (RED) documents: <u>https://archive.epa.gov/pesticides/reregistration/web/html/status.html</u>
- Office of Pesticide Programs Pesticide Chemical Search: https://ofmpub.epa.gov/apex/pesticides/f?p=CHEMICALSEARCH:1
- Food and Agriculture Organization of the United Nations: <u>http://www.fao.org/home/en/</u>
- PAN Pesticides Database: <u>http://www.pesticideinfo.org/Search\_Chemicals.jsp</u>

Prior to inclusion in the search term string, all forms of chemical names were subjected to verification from several potential sources (*e.g.*, US EPA Chemistry Dashboard, STN International-CAS; see complete list of sources for chemical verification in Table\_Apx A-1). From these sources, all chemical names, synonyms, CAS number(s), trade names, etc. were documented and used to generate terms for database searches.

CHEMICAL SOURCE	CONTENTS	DOCUMENT LOCATION
Chemistry Dashboard (https://comptox.epa.gov/dashboard)	CAS Numbers, Synonyms, Structures, Properties, Environmental Fate and Transport.	Online
Dictionary of Chemical Names and Synonyms	Wide assortment of chemical compounds by chemical name and synonym, has CAS index and some structure data	ECOTOX
Farm Chemicals Handbook-1992	Pesticide information, CAS numbers and synonyms, some structure data ***Sometimes CAS number presented for a compound is for the main constituent only	ECOTOX
OPPT SMILES Verification Source	Structure Data	Electronic verification
RTECS (Registry of Toxic Effects of chemical substance, 1983-84 ed., 2 vols)	Chemical names, synonyms and CAS numbers	ECOTOX
Sigma – Aldrich website58784 http://www.sigma-aldrich.com	Organic and inorganic Compounds by chemical name, has CAS index and some structure and Physical Property data	Online

#### Table\_Apx A-1. Sources of Verification for Chemical Names and Structures

CHEMICAL SOURCE	CONTENTS	DOCUMENT LOCATION
STN International (CAS) 1994	***Most complete source of chemical name, synonym	Online
	and structure information, no physical properties	
The Pesticide Manual 10th edition, 1994	Pesticide Compounds by chemical name, synonym,	ECOTOX
	product code, has CAS index and some structure and	
	Physical Property data	
TSCA (Toxic Substances Control Act	Chemical names, synonyms and CAS numbers	ECOTOX
Chemical Substance Inventory, 1985 ed., 5		
vols)		
World Wide Web (misc. web sources) A copy	Chemical names, synonyms and CAS numbers	Online
of the verification page is saved to the		
Attachments tab of the chemical entry. This		
includes company MSDS sheets or Chemical		
Labels.		0.1
California Department of Pesticide Regulation	Multiple databases containing chemicals, pesticides,	Online
(http://www.cdpr.ca.gov/dprdatabase.htm) PAN Pesticide Database	companies, products, etc.	Online
	Pesticides searchable by name or CAS #. Includes CAS	Online
( <u>http://www.pesticideinfo.org/Search_Chemic</u>	#, Name, synonyms, targets, toxicity data, related	
als.jsp)	chemicals and regulatory information.	0.1
US EPA Office of Pesticide Programs	Multiple databases containing chemicals, pesticides,	Online
Pesticide Fate Database – No web access	companies, products, etc.	
available. An electronic copy of the data file is		
located at the Contractor site:		
PFATE_37_Tables.mdb.		

### A.1.2 Publicly Available Database Searches

The databases listed below were searched for literature containing the chemical search terms. Database searching occurred during April and May of 2019 by an information specialist and the results were stored in the Health and Environmental Research Online (HERO) database and assigned a HERO reference identification number.<sup>9</sup> The present literature search focused only on the chemical name (including synonyms and trade names) with no additional limits. Full details of the search strategy for each database are presented in Appendix A.1.2.1.

After initial deduplication in HERO<sup>10</sup>, these studies were imported into <u>SWIFT Review</u> software to identify those references most likely to be applicable to each discipline area (*i.e.*, consumer, environmental, and general population exposure, occupational exposure and environmental releases, environmental hazards, human health hazards, and fate and physical chemistry).

### A.1.2.1 Query Strings for the Publicly Available Database Searches on 1,1,2-Trichlorobenzene

Table\_Apx A-2 presents a list of the data sources, the search dates and number of peer-reviewed references resulting from the searches for 1,1,2-trichloroethane. The sources are found as online

<sup>&</sup>lt;sup>9</sup>EPA's HERO database provides access to the scientific literature behind EPA science assessments. The database includes more than 600,000 scientific references and data from the peer-reviewed literature used by EPA to develop its regulations.

<sup>&</sup>lt;sup>10</sup> Deduplication in HERO involves first determining whether a matching unique ID exists (*e.g.*, PMID, WOSid, or DOI). If one matches one that already exists in HERO, HERO will tag the existing reference instead of adding the reference again. Second, HERO checks if the same journal, volume, issue and page number are already in HERO. Third, HERO matches on the title, year, and first author. Title comparisons ignore punctuation and case.

databases and the resulting references were gathered and uploaded into the EPA Health and Environmental Research Online (HERO) database for literature screening.

Table_Apx A-2. Summary of Data Sources, Search Dates and Number of Peer-Reviewed
Literature Search Results for 1,1,2-Trichloroethane

Source	Date of Search	Number of References
Current Contents	05/09/2019	1576
Web of Science	09/10/2019	2809
ProQuest CSA	05/09/2019	3874
Dissertation Abstracts	05/10/2019	39
Science Direct	05/09/2019	2156
Agricola	05/10/2019	509
TOXNET	05/09/2019	3635
PubMed	07/02/2019	1489
UNIFY	05/09/2019	183
Totals:		16,270

#### General:

General search terms were compiled and used in the search strategies for each of the databases/sources listed below. Based upon the online search manuals for the respective databases/sources, it was necessary to construct searches as noted for each of the sources. The search terms are listed below in full for each source and noted if the general search terms or other search terms were used.

."1,1,1-Trichloraethan" OR "1,1,1-Trichlorethan" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichlorethane" OR "1,1,2-Trichloroethane" OR "Setteme TT" OR "alpha-Trichloroethane" OR "Baltana" OR "beta-trichloroethane" OR "Chlorothene TT" OR "alpha-Trichloroethane" OR "Chlorothene" OR "Chlorothene NU" OR "Chlorothene SM" OR "Chlorothene VG" OR "Chlorothene" OR "Chlorothene NU" OR "Chlorothene SM" OR "Chlorothene VG" OR "Chlorothene OR "Chlorothene SM" OR "Chlorothene SM" OR "HCC 140a" OR "ICI-CF 2" OR "Inhibisol" OR "Methyl chloroform" OR "Methylchloroform" OR "Methyltrichloromethane" OR "NCI-C04579" OR "NCI-C04626" OR "NSC 405074" OR "NSC 9367" OR "Solvent 111" OR "Tafclean" OR "Three One A" OR "Trichloroethane" OR "Trichloroethane" OR "Trichloroethane" OR "UNII-28E9ERN9WU" OR "Used"

."Vinyl trichloride" OR "Vinyltrichloride"

Based upon the online search manuals for the respective databases below, it was necessary to construct searches as follows:

<u>**CURRENT CONTENTS CONNECT</u>**: (access.webofknowledge.com) General Search Terms applied to the search strategy for Current Contents.</u>

Date Searched: 05/09/2019

Date Range of Search: 1970 to Present N = 1.576

TS=("1,1,1-Trichloraethan" OR "1,1,1-Trichlorethan" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichloroethane" OR "1,1,2-TRICHLORAETHAN" OR "1,1,2-Trichlorethan" OR "1,1,2-trichlorethane" OR "1,1,2-Trichloroethane" OR "1,2,2-Trichloroethane" OR "111trichloroethane" OR "112trichloroethane" OR "Aerothene TT" OR "alpha-Trichloroethane" OR "Baltana" OR "beta-trichloroethane" OR "Caswell No. 875" OR "Caswell No. 875A" OR "Chlorotene" OR "Chlorothene" OR "Chlorothene NU" OR "Chlorothene SM" OR "Chlorothene VG" OR "Chlorten" OR "Cleanite" OR "Dowclene LS" OR "Ethana NU" OR "Genklene LB" OR "HCC 140a" OR "ICI-CF 2" OR "Inhibisol" OR "Methyl chloroform" OR "Methylchloroform" OR "Methyltrichloromethane" OR "NCI-C04579" OR "NCI-C04626" OR "NSC 405074" OR "NSC 9367" OR "Solvent 111" OR "Tafclean" OR "Three One A" OR "Three One S" OR "trichloro-1,1,1-ethane" OR "Trichloroethane" OR "Trichloromethylmethane" OR "Tricloroethane" OR "UN 2831" OR "UNII-113C650IR1" OR "UNII-28E9ERN9WU") N = 1.576

TS=("Vinyl trichloride" OR "Vinyltrichloride") N = 0

## **WOS Core Collection**:

Web of Science Core Collection may be accessed through EPA Desktop Library (https://intranet.epa.gov/desktop/databases.htm) by clicking on the Web of Science Link or copying and pasting (https://apps.webofknowledge.com).

Date Searched: 09/10/2019 Date Range of Search: 1970 to Present N = 2809

TS=("1,1,1-Trichloraethan" OR "1,1,1-Trichlorethan" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichloroethane" OR "1,1,2-TRICHLORAETHAN" OR "1,1,2-Trichlorethan" OR "1,1,2-trichlorethane" OR "1,1,2-Trichloroethane" OR "1,2,2-Trichloroethane" OR "111trichloroethane" OR "112trichloroethane" OR "Aerothene TT" OR "alpha-Trichloroethane" OR "Baltana" OR "beta-trichloroethane" OR "Caswell No. 875" OR "Caswell No. 875A" OR "Chlorotene" OR "Chlorothene" OR "Chlorothene NU" OR "Chlorothene SM" OR "Chlorothene VG" OR "Chlorten" OR "Cleanite" OR "Dowclene LS" OR "Ethana NU" OR "Genklene LB" OR "HCC 140a" OR "ICI-CF 2" OR "Inhibisol" OR "Methyl chloroform" OR "Methylchloroform" OR "Methyltrichloromethane" OR "NCI-C04579" OR "NCI-C04626" OR "NSC 405074" OR "NSC 9367" OR "Solvent 111" OR "Tafclean" OR "Three One A" OR "Three One S" OR "trichloro-1,1,1-ethane" OR "Trichloroethane" OR "Trichloromethylmethane" OR "Tricloroethane" OR "UN 2831" OR "UNII-113C650IR1" OR "UNII-28E9ERN9WU")

N = 2809

TS=("Vinyl trichloride" OR "Vinyltrichloride") N = 0

PROQUEST Agricultural and Scientific Database: (www.csa.com)

General Search Terms applied to the search strategy for ProQuest Agricultural and Scientific Database.

Date Searched: 05/09/2019Date Range of Search: 1900 to Present N = 3,874

ALL("1,1,1-Trichloraethan" OR "1,1,1-Trichlorethan" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichloroethane" OR "1,1,2-TRICHLORAETHAN" OR "1,1,2-Trichlorethan" OR "1,1,2-trichlorethane" OR "1,1,2-Trichloroethane" OR "1,2,2-Trichloroethane" OR "111trichloroethane" OR "112trichloroethane" OR "Aerothene TT" OR "alpha-Trichloroethane" OR "Baltana" OR "beta-trichloroethane" OR "Caswell No. 875" OR "Caswell No. 875A" OR "Chlorotene" OR "Chlorothene" OR "Chlorothene NU" OR "Chlorothene SM" OR "Chlorothene VG" OR "Chlorotene" OR "Chlorothene" OR "Dowclene LS" OR "Ethana NU" OR "Genklene LB" OR "HCC 140a" OR "ICI-CF 2" OR "Inhibisol" OR "Methyl chloroform" OR "Methylchloroform" OR "Methyltrichloromethane" OR "NCI-C04579" OR "NCI-C04626" OR "NSC 405074" OR "NSC 9367" OR "Solvent 111" OR "Tafclean" OR "Three One A" OR "Three One S" OR "trichloro-1,1,1-ethane" OR "Trichloroethane" OR "Trichloromethylmethane" OR "Tricloroethane" OR "UN 2831" OR "UNII-113C650IR1" OR "UNII-28E9ERN9WU") AND STYPE("Scholarly Journals" OR Reports OR Thesis OR "Government Documents") AND LA(ENG) N = 3,874

ALL("Vinyl trichloride" OR "Vinyltrichloride") AND STYPE("Scholarly Journals" OR Reports OR Thesis OR "Government Documents") AND LA(ENG) N = 0

PROQUEST Dissertations and Theses: (search.proquest.com)

General Search Terms applied to the search strategy for ProQuest Dissertations and Theses.

Date Searched: 05/10/2019Date Range of Search: 1900 to Present N = 39

ALL("1,1,1-Trichloraethan" OR "1,1,1-Trichlorethan" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichloroethane" OR "1,1,2-TRICHLORAETHAN" OR "1,1,2-Trichlorethan" OR "1,1,2-trichloroethane" OR "1,1,2-Trichloroethane" OR "1,2,2-Trichloroethane" OR "111trichloroethane" OR "112trichloroethane" OR "Aerothene TT" OR "alpha-Trichloroethane" OR "Baltana" OR "beta-trichloroethane" OR "Caswell No. 875" OR "Caswell No. 875A" OR "Chlorotene" OR "Chlorothene" OR "Chlorothene NU" OR "Chlorothene SM" OR "Chlorothene VG" OR "Chlorotene" OR "Chlorothene" OR "Dowclene LS" OR "Ethana NU" OR "Genklene LB" OR "HCC 140a" OR "ICI-CF 2" OR "Inhibisol" OR "Methyl chloroform" OR "Methylchloroform" OR "Methyltrichloromethane" OR "NCI-C04579" OR "NCI-C04626" OR "NSC 405074" OR "NSC 9367" OR "Solvent 111" OR "Tafclean" OR "Three One A" OR "Tricloroethane" OR "UN 2831" OR "UNII-113C650IR1" OR "UNII-28E9ERN9WU") AND LA(ENG) N = 31

ALL("Vinyl trichloride" OR "Vinyltrichloride") AND LA(ENG)

N = 8

**<u>SCIENCE DIRECT</u>**: (www.sciencedirect.com) General Search Terms applied to the search strategy for Science Direct

Date Searched: 05/09/2019Date Range of Search: 1823 to Present N = 2,156

Science Direct 01: "1,1,1-Trichloraethan" OR "1,1,1-Trichlorethan" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichlorethane" OR "1,1,2-Trichlorethane" OR "1,1,2-Trichlorethane" OR "1,1,2-Trichlorethane" OR "1,1,2-Trichlorethane" N = 663 (7 downloads)

Science Direct 02: "1,2,2-Trichloroethane" OR "111trichloroethane" OR "112trichloroethane" OR "Aerothene TT" OR "alpha-Trichloroethane" OR "Baltana" OR "beta-trichloroethane" OR "Caswell No. 875" OR "Caswell No. 875A" N = 9 (1 download)

Science Direct 03: "Chlorothene" OR "Chlorothene NU" OR "Chlorothene SM" OR "Chlorothene VG" OR "Chlorten" OR "Cleanite" OR "Dowclene LS" OR "Ethana NU" N = 0

Science Direct 04: "Genklene LB" OR "HCC 140a" OR "ICI-CF 2" OR "Inhibisol" OR "Methyl chloroform" OR "Methylchloroform" OR "Methyltrichloromethane" OR "NCI-C04579" OR "NCI-C04626" N = 97 (1 download)

Science Direct 05: "NSC 405074" OR "NSC 9367" OR "Solvent 111" OR "Tafclean" OR "Three One A" OR "Three One S" OR "trichloro-1,1,1-ethane" OR "Trichloroethane" OR "Trichloromethylmethane" N = 1,386 (14 downloads)

```
Science Direct 06:

"Tricloroethane" OR "UN 2831" OR "UNII-113C650IR1" OR "UNII-28E9ERN9WU"

N = 1

Science Direct 07:

"Vinyl trichloride" OR "Vinyltrichloride"

N = 0
```

#### AGRICOLA: (www.nal.usda.gov)

General Search Terms applied to the search strategy for Agricola. The Agricola database contains a significant amount of gray literature including proceedings, symposia, and progress reports from

government and educational institutions. Agricola is not used when conducting a search for the Office of Water.

Date Searched: 05/10/2019Date Range of Search: 15th century to the Present N = 509

Agricola 01: 1,1,1-Trichloraethan 1,1,1-Trichlorethan 1,1,1-Trichlorethane 1,1,1-Trichlorethane 1,1,1-Trichloroethane 1,1,2-TRICHLORAETHAN 1,1,2-Trichlorethan 1,1,2-trichlorethane 1,1,2-Trichloroethane 1,2,2-Trichloroethane Agricola 02: 111trichloroethane 112trichloroethane Aerothene TT alpha-Trichloroethane Baltana beta-trichloroethane Caswell No. 875 Caswell No. 875A Chlorotene Chlorothene Agricola 03: Chlorothene NU Chlorothene SM Chlorothene VG Chlorten Cleanite Dowclene LS Ethana NU Genklene LB HCC 140a ICI-CF 2 Agricola 04: Inhibisol Methyl chloroform Methylchloroform

Methyltrichloromethane NCI-C04579 NCI-C04626 NSC 405074 NSC 9367 Solvent 111 Tafclean Agricola 05: Three One A Three One S trichloro-1,1,1-ethane Trichloroethane Trichloromethylmethane Tricloroethane UN 2831 UNII-113C650IR1 UNII-28E9ERN9WU

Agricola 06: Vinyl trichloride Vinyltrichloride

**<u>TOXNET</u>**: (toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?TOXLINE) General Search Terms applied to the search strategy for TOXNET.

Date Searched: 05/09/2019Date Range of Search: 1900 to Present N = 3635

TOXNET 01: 79-00-5 OR 71-55-6 OR 25323-89-1 OR 1299-89-4 OR 74552-83-3 N = 3,635

#### PubMed:

PubMed may be accessed through the EPA Desktop Library (https://www.ncbi.nlm.nih.gov/pubmed/)

Date Searched: 07/02/2019Date Range of Search: 1900 to present N = 1489

."1,1,1-Trichloraethan" OR "1,1,1-Trichlorethan" OR "1,1,1-Trichlorethane" OR "1,1,1-Trichlorethane" OR "1,1,2-Trichlorethane" OR "1,1,2-Trichlorethane" OR "1,1,2-Trichlorethane" OR "1,1,2-Trichloroethane" OR "1,2,2-Trichloroethane" OR "1,1,2-Trichloroethane" OR "1,1,2-Trichloroethane" OR "1,2,2-Trichloroethane" OR "1,1,2-Trichloroethane" OR "1,2,2-Trichloroethane" OR "Baltana" OR "beta-trichloroethane" OR "Caswell No. 875" OR "Caswell No. 875A" OR "Chlorothene" OR "Chlorothene" OR "Chlorothene SM" OR "Chlorothene VG" OR "Chlorten" OR "Cleanite" OR

"Dowclene LS" OR "Ethana NU" OR "Genklene LB" OR "HCC 140a" OR "ICI-CF 2" OR "Inhibisol" OR "Methyl chloroform" OR "Methylchloroform" OR "Methyltrichloromethane" OR "NCI-C04579" OR "NCI-C04626" OR "NSC 405074" OR "NSC 9367" OR "Solvent 111" OR "Tafclean" OR "Three One A" OR "Three One S" OR "trichloro-1,1,1-ethane" OR "Trichloroethane" OR "UNII-113C650IR1" OR "UNII-28E9ERN9WU" OR "Used" N = 1489

."Vinyl trichloride" OR "Vinyltrichloride" N = 0

### ECOTOX UNIFY:

This is an internal EPA database that is not accessible to the public. Results from the ECOTOX Unify search strategy.

Date Searched: 05/09/2019Date Range of Search: all years N = 183

## A.1.2.2 Data Prioritization for Environmental Hazard, Human Health Hazard, Fate and Physical Chemistry

In brief, SWIFT Review has pre-set literature search strategies ("filters") developed by information specialists that can be applied to identify studies that are more likely to be useful for identifying human health and ecotoxicity content from those that likely do not (*e.g.*, analytical methods). The filters function like a typical search strategy where studies are tagged as belonging to a certain filter if the terms in the filter literature search strategy appear in title, abstract, keyword or medical subject headings (MeSH) fields content. The applied SWIFT Review filters focused on lines of evidence: human, animal models for human health, ecological taxa (which includes ecotoxicological animal models, plants, and other taxa), and *in vitro* studies. The details of the search strategies that underlie the filters are available online. Studies not retrieved using these filters were not considered further. Studies that included one or more of the search terms in the title, abstract, keyword, or MeSH fields were exported as a RIS file for screening in Swift-ActiveScreener or DistillerSR<sup>11</sup>.

#### A.1.2.3 Data Prioritization for Occupational Exposures and Environmental Releases and Gen Pop, Consumer and Environmental Exposures

To prioritize references related to occupational exposure, environmental release, general population exposure, consumer exposure, and environmental exposure, EPA used positive and negative seed studies to build a classification model in SWIFT Review. The positive seeds were identified using relevant literature pool for the first ten TSCA risk evaluations, while the negative seeds were identified from a subset of literature for the current high-priority substances. The model was then applied to the unclassified literature to generate a classification score for each reference. Scores above a certain threshold value were then prioritized for further review in SWIFT-ActiveScreener.

<sup>&</sup>lt;sup>11</sup><u>DistillerSR</u> is a web-based systematic review software used to screen studies available at https://www.evidencepartners.com/products/distillersr-systematic-review-software.

## A.2 Peer-Reviewed Screening Process

The studies identified from publicly available database searches and SWIFT-Review filtering/prioritization were housed in HERO system and imported into SWIFT-ActiveScreener or DistillerSR for title/abstract and full-text screening. Both title/abstract and full-text screening were conducted by two independent reviewers. Screening is initiated with a pilot phase of screening (between 10 and 50) studies to identify areas where clarification in screening criteria might be needed or chemical-specific supplemental material tags might be identified. Records that met PECO (or equivalent criteria (Appendix A.2.1) during title and abstract screening were considered for full-text screening. At both the title/abstract and full-text review levels, screening conflicts were resolved by topic-specific experts and/or discussion among the primary screeners. For citations with no abstract, the articles are initially screened based on all or some of the following: title relevance (titles that suggest a record is not relevant can be excluded rather than marked as unclear), and page numbers (articles two pages in length or less were assumed to be conference reports, editorials, or letters). During title/abstract or full-text level screening in DistillerSR, studies that did not meet the PECO criteria, but which could provide supporting information were categorized (or "tagged") as supplemental information.

It is important to emphasize that being tagged as supplemental material does not mean the study would necessarily be excluded from consideration in an assessment. The initial screening level distinctions between a study meeting the PECO criteria and a supplemental study are often made for practical reasons and the tagging structures (as seen in the literature inventory trees and heat maps in Section 2.1. of this document) are designed to ensure the supplemental studies are categorized for easy retrieval if needed while conducting the assessment. The impact on the assessment conclusions of individual studies tagged as supporting material is often difficult to assess during the screening phase of the assessment. These studies may emerge as being critically important to the assessment and need to be evaluated and summarized at the individual study level (e.g., cancer MOA mechanistic or non-English-language studies), or be helpful to provide context (e.g., summarize current levels of exposure, provide hazard evidence from routes or durations of exposure not pertinent to the PECO), or not be cited at all in the assessment (e.g., individual studies that contribute to a well-established scientific conclusion). Studies maybe be tagged as supplemental material during either title and abstract or full-text screening. When tagged as supplemental material during title and abstract screening, it may not be completely clear whether the chemical of interest is reported in the study (*i.e.*, abstracts may not describe all chemicals investigated). In these cases, studies are still tagged with the expectation that if full-text retrieval is pursued, then additional screening would be needed to clarify if the study is pertinent.

## A.2.1 Inclusion/Exclusion Criteria

A PECO statement is typically used to focus the research question(s), search terms, and inclusion/exclusion criteria in a systematic review. PECO criteria were developed *a priori* to screening and modified to fit the various discipline areas supporting the TSCA risk evaluations. Variations include the RESO (receptor, exposure, scenario/setting, and outcome) used for the occupational exposure and environmental releases discipline, and PESO (pathways/processes, exposures, setting/scenario, and outcomes) used by the fate and transport discipline. All PECOs and PECO-equivalent criteria can be found in the following sections.

## A.2.1.1 PECO for Environmental and Human Health Hazards

The PECO used in this evidence map to identify literature pertinent to 1,1,2-trichloroethane effects on human health and environmental hazard is presented in Table\_Apx A-3. In addition to the PECO

criteria, studies containing potentially relevant supplemental material were tracked and categorized during the literature screening process as outlined in Table\_Apx A-4.

PECO	Evidence
Element	
Р	<ul> <li>Human: Any population and life stage (<i>e.g.</i>, occupational or general population, including children and other sensitive populations).</li> <li>Animal: Aquatic and terrestrial species (live, whole organism) from any life stage (<i>e.g.</i>, preconception, in utero, lactation, peripubertal, and adult stages). Tests of the single toxicants in <i>in vitro</i> systems or on live, whole, taxonomically verifiable organisms (<i>e.g.</i>, gametes, embryos, or plant or fungal sections capable of forming whole, new organisms) that are not bacteria, humans, monkeys, viruses, or yeast. In most cases, transgenic animal models will get screened as "yes" or "unclear" at the Title and Abstract (TIAB) screening level. Although certain non-mammalian model systems are increasing used to identify potential human health hazards (<i>e.g.</i>, <i>Xenopus</i> and zebrafish), for simplicity animal models will be further inventoried according to the categorization below:</li> <li><u>Human health models</u>: rat, mouse, rabbit, dog, hamster, guinea pig, cat, non-human primate, pig, hen (neurotoxicity only)</li> <li><u>Environmental models</u>: invertebrates (<i>e.g.</i>, insects, spiders, crustaceans, mollusks, and worms) and vertebrates (<i>e.g.</i>, mammals and all amphibians, birds, fish, and reptiles). All hen studies (including</li> </ul>
	<ul> <li>neurotoxicity studies) will be included for ecotoxicological models.</li> <li>Plants: All aquatic and terrestrial species (live), including algal, moss, lichen and fungi species.</li> </ul>
	<ul> <li>Screener note:</li> <li>To identify human health and environmental hazards, other organisms not listed above in their respective categories can also be used. Non-mammalian model systems are increasingly used to identify potential human health hazards (<i>e.g.</i>, Xenopus, zebrafish), and traditional human health models (<i>e.g.</i>, rodents) can be used to identify potential environmental hazard. Neurotoxicity studies performed in hens (<i>e.g.</i>, OECD 418 and 419) are considered relevant to both human and eco hazard.</li> <li>PECO considerations should be directed toward effects on target species only and not on the indirect effects expressed in taxa as a result of chemical treatment (<i>e.g.</i>, substance is lethal to a targeted pest species leading to positive effects on plant growth due to diminished presence of the targeted pest species).</li> </ul>
	Relevant forms and isomers         1,1,2-Trichloroethane (CASRN 79-00-5)         Related isomers:         1,1,1-Trichloroethane (CASRN 71-55-6)         Trichloroethane (CASRN 25323-89-1)
Ε	<ul> <li>For synonyms see the EPA Chemistry Dashboard.</li> <li>Human: Any exposure to 1,1,2-Trichloroethane.</li> <li>Animal: Any exposure to 1,1,2-Trichloroethane, including via water, soil or sediment, injection (<i>i.e.</i>, oral or topical), gavage, diet, dermal, and inhalation.</li> <li>Plants: Exposure to 1,1,2-Trichloroethane via water and/or soil, with reported concentration and duration. Studies involving exposures to mixtures will be included only if they also include exposure to one of these solvents alone.</li> </ul>
	Screener note:

Table\_Apx A-3. Hazards Title and Abstract and Full-text PECO Criteria for 1,1,2-Trichloroethane

PECO Element	Evidence
	Field studies with media concentrations (surface water, interstitial water, soil) and/or body/tissue concentrations of animals or plants are to be identified as <u>Supplemental</u> if any biological effects are reported.
	<ul> <li>Human: A comparison or referent population exposed to lower levels (or no exposure/exposure below detection limits) of 1,1,2-Trichloroethane, or exposure to one of these solvents for shorter periods of time. case-crossover, case-referent, case-only, case-specular, case-cohort, case-parent, nested case-control study designs are all included.</li> <li>Animal and Plants: A concurrent control group exposed to vehicle-only treatment and/or untreated control (control could be a baseline measurement).</li> </ul>
C	<ul> <li>Screener note:</li> <li>If no control group is explicitly stated or implied (<i>e.g.</i>, by mention of statistical results that could only be obtained if a control group was present), the study will be marked as <u>unclear</u> during Title/Abstract Screening.</li> <li>All case reports and case studies/series describing findings in a sample size of less than 20 people in any setting (<i>e.g.</i>, occupation, general population) will be tracked as "potentially relevant supplemental information".</li> </ul>
	<ul> <li>Human: All health outcomes (cancer and noncancer).</li> <li>Animal and Plants: All biological effects (including bioaccumulation from laboratory studies with concurrently measured water and tissue concentrations).</li> </ul>
0	<ul> <li>Screener note:</li> <li>Measurable biological effects relevant for humans, animals and plants may include but are not limited to: mortality, behavioral, population, cellular, physiological, growth, reproduction of an acceptable organism to a chemical toxicant.</li> </ul>

# Table\_Apx A-4. Major Categories of Potentially Relevant Supplemental Materials for 1,1,2 Trichloroethane

Category	Evidence	
Mechanistic studies	Studies reporting measurements related to a health outcome that inform the biological or chemical events associated with phenotypic effects, in both mammalian and non-mammalian model systems, including <i>in vitro</i> , <i>in vivo</i> (by various non-inhalation routes of exposure), <i>ex vivo</i> , and <i>in silico</i> studies.	
ADME, PBPK, and toxicokinetic	Studies designed to capture information regarding absorption, distribution, metabolism, and excretion (ADME), toxicokinetic studies, or physiologically based pharmacokinetic (PBPK) models.	
Susceptible populations (no health outcome)	Studies that identify potentially susceptible subgroups; for example, studies that focus on a specific demographic, life stage, or genotype.	
Mixture studies	Mixture studies that are not considered PECO-relevant because they do not contain an exposure or treatment group assessing only the chemical of interest.	
Case reports or case series	Case reports ( $n \le 3$ cases) and case series/studies (<20 cases) will be tracked as potentially relevant supplemental information.	
Records with no original data	Records that do not contain original data, such as other agency assessments, informative scientific literature reviews, editorials or commentaries.	
Conference abstracts	Records that do not contain sufficient documentation to support study evaluation and data extraction.	
Field Studies	Field studies where there are accompanying body/tissue concentrations of animals without any biological effects reported	

A.2.1.2 PECO for Consumer, Environmental, and General Population Exposures Table\_Apx A-5. General PECO for Consumer, Environmental, and General Population Exposures

PECO Element	Evidence
<b>P</b> opulation	Human: General population; consumers; bystanders in the home; near-facility populations (includes industrial and commercial facilities manufacturing, processing, or using the chemical substance); children; susceptible populations (life stages, preexisting conditions, genetic factors), pregnant women; lactating women, women of child bearing age. Many human population groups may be exposed. No chemical-specific exclusions are suggested at this time.Environmental: studies only)
	Expected Primary Exposure Sources, Pathways, Routes:
<u>E</u> xposure	<u>Pathways:</u> indoor air/vapor/mist; indoor dust; particles; outdoor/ambient air; surface water; biosolids; sediment; breastmilk; food items containing 1,1,2- trichloroethane including fish; consumer product uses in the home (including consumer product containing chemical);
	Routes of Exposure: Inhalation, Oral, Dermal
Comparator	<b>Human</b> : Consider media-specific background exposure scenarios and use/source specific exposure scenarios as well as which receptors are and are not reasonably exposed across the projected exposure scenarios.
(Scenario)	<b>Environmental</b> Consider media-specific background exposure scenarios and use/source specific exposure scenarios as well as which receptors are and are not reasonably exposed across the projected exposure scenarios.
<u>O</u> utcomes for Exposure Concentration or	<b>Human:</b> Acute, subchronic, and/or indoor air and water concentration estimates (mg/m <sup>3</sup> or mg/L). Both external potential dose and internal dose based on biomonitoring and reverse dosimetry mg/kg/day will be considered. Characteristics of consumer products or articles (weight fraction, emission rates, etc) containing 1,1,2- trichloroethane
Dose	<b>Environmental:</b> A wide range of ecological receptors will be considered (range depending on available ecotoxicity data) using surface water concentrations, sediment concentrations.

## Table\_Apx A-6. Pathways Identified as Supplemental for 1,1,2-Trichloroethane<sup>a</sup>

Chemical	Drinking Water	Ambient Air	Air Disposal	Land Disposal	Underground Disposal	Ground Water
1,1,2-Trichloroethane	Х	Х	Х	Х	Х	Х

<sup>a</sup> "Supplemental pathways" refer to pathways addressed by other EPA administered statutes (see Section 2.6.3.1). Studies tagged under these pathways provide media information that is not prioritized in the screening process.

## A.2.1.3 RESO for Occupational Exposure and Environmental Releases

EPA developed a generic RESO statement to guide the screening of engineering and occupational exposure data or information sources for the TSCA risk evaluations. Data or information sources that comply with the inclusion criteria specified in the RESO statement are eligible for inclusion, considered

for evaluation, and possibly included in the environmental release and occupational exposure assessments. On the other hand, data or information sources that fail to meet the criteria in the RESO statement are excluded from further consideration.

Assessors seek information on various chemical-specific engineering and occupational exposure data needs as part of the process of developing the exposure assessment for each risk evaluation. EPA uses the RESO statement (Table\_Apx A-7) along with the information in Table\_Apx A-8 when screening environmental release and occupational exposure data and information.

Table_Apx A-7. Inclusion Criteria for Data Sources Reporting Environmental Release and
Occupational Exposure Data

<b>RESO Element</b>	Evidence
<u>R</u> eceptors	<ul> <li><u>Humans</u>: Workers, including occupational non-users</li> <li><u>Environment</u>: All environmental receptors (relevant release estimates input to Exposure)</li> </ul>
	Please refer to the conceptual models for more information about the environmental and human receptors included in the TSCA risk evaluation.
<u>E</u> xposure	<ul> <li>Worker exposure to and relevant environmental releases of the chemical substance from occupational scenarios:         <ul> <li>Dermal and inhalation exposure routes (as indicated in the conceptual model)</li> <li>Oral route (as indicated in the conceptual model)</li> </ul> </li> </ul>
	Please refer to the conceptual models for more information about the routes and media/pathways included in the TSCA risk evaluation.
<u>S</u> etting or <u>Scenario</u>	• Any occupational setting or scenario resulting in worker exposure and relevant environmental releases (includes all manufacturing, processing, use, disposal.
<u>O</u> utcomes	<ul> <li>Quantitative estimates* of worker exposures and of relevant environmental releases from occupational settings</li> <li>General information and data related and relevant to the occupational estimates*</li> </ul>

\* Metrics (*e.g.*, mg/kg/day or mg/m<sup>3</sup> for worker exposures, kg/site/day for releases) are determined by toxicologists for worker exposures and by exposure assessors for releases; also, the Engineering, Release and Occupational Exposure Data Needs (Table\_Apx A-8)) provides a list of related and relevant general information. TSCA=Toxic Substances Control Act

Develop the Environmental Release and Occupational Exposure Assessments           Objective				
Determined	Type of Data <sup>a</sup>			
during Scoping				
General Engineering Assessment (may apply to Occupational Exposures and / or Environmental Releases)	<ul> <li>Description of the life cycle of the chemical(s) of interest, from manufacture to end-of-life (<i>e.g.</i>, each manufacturing, processing, or use step), and material flow between the industrial and commercial life cycle stages.</li> <li>The total annual U.S. volume (lb/yr or kg/yr) of the chemical(s) of interest manufactured, imported, processed, and used; and the share of total annual manufacturing and import volume that is processed or used in each life cycle step.</li> <li>Description of processes, equipment, and unit operations during each industrial/ commercial life cycle step.</li> <li>Material flows, use rates, and frequencies (lb/site-day or kg/site-day and days/yr; lb/site-batch and batches/yr) of the chemical(s) of interest during each industrial/ commercial life cycle step. Note: if available, include weight fractions of the chemicals (s) of interest and material flows of all associated primary chemicals (especially water).</li> <li>Number of sites that manufacture, process, or use the chemical(s) of interest for each industrial/ commercial life cycle step and site locations.</li> </ul>			
Occupational Exposures	<ul> <li>Concentration of the chemical of interest</li> <li>Description of worker activities with exposure potential during the manufacture, processing, or use of the chemical(s) of interest in each industrial/commercial life cycle stage.</li> <li>Potential routes of exposure (<i>e.g.</i>, inhalation, dermal).</li> <li>Physical form of the chemical(s) of interest for each exposure route (<i>e.g.</i>, liquid, vapor, mist) and activity.</li> <li>Breathing zone (personal sample) measurements of occupational exposures to the chemical(s) of interest, measured as time-weighted averages (TWAs), short-term exposures, or peak exposures in each occupational life cycle stage (or in a workplace scenario similar to an occupational life cycle stage).</li> <li>Area or stationary measurements of airborne concentrations of the chemical(s) of interest in each occupational setting and life cycle stage (or in a workplace scenario similar to the life cycle stage of interest).</li> <li>For solids, bulk and dust particle size characterization data.</li> <li>Dermal exposure data.</li> <li>Exposure frequency (days/yr).</li> <li>Number of workers who potentially handle or have exposure to the chemical(s) of interest in each occupational life cycle stage.</li> <li>Personal protective equipment (PPE) types employed by the industries within scope.</li> <li>Engineering controls employed to reduce occupational exposures in each occupational life cycle stage (or in a workplace scenario sime ach occupational life cycle stage of interest in each occupational life cycle stage.</li> </ul>			
Environmental Releases (to relevant environmental media)	<ul> <li>Description of sources of potential environmental releases, including cleaning of residues from process equipment and transport containers, involved during the manufacture, processing, or use of the chemical(s) of interest in each life cycle stage.</li> <li>Estimated mass (lb or kg) of the chemical(s) of interest released from industrial and commercial sites to each environmental medium (water) and treatment and disposal methods (POTW), including releases per site and aggregated over all sites (annual release rates, daily release rates)</li> <li>Release or emission factors.</li> <li>Number of release days per year.</li> </ul>			

## Table\_Apx A-8. Engineering, Environmental Release and Occupational Data Necessary to Develop the Environmental Release and Occupational Exposure Assessments

Objective						
Determined	Type of Data <sup>a</sup>					
during Scoping						
	Waste treatment methods and pollution control devices employed by the industries within scope and					
	associated data on release/emission reductions.					
<sup>a</sup> These are the tage	s included in the full-text screening form. The screener makes a selection from these specific tags, which					
describe more spec	ific types of data or information.					
In addition to the d	ata types listed above, EPA may identify additional data needs for mathematical modeling. These data					
needs will be deter	mined on a case-by-case basis.					
Abbreviations:						
hr=Hour						
kg=Kilogram(s)						
lb=Pound(s)						
yr=Year	yr=Year					
PV=Particle volum	PV=Particle volume					
POTW=Publicly owned treatment works						
PPE=Personal protection equipment						
PSD=Particle size	PSD=Particle size distribution					
TWA=Time-weigh	ited average					

### A.2.1.4 PESO for Fate and Transport

EPA developed a generic PESO statement to guide the screening of environmental fate data or information sources for the TSCA risk evaluations. Data or information sources that comply with the inclusion criteria in the PESO statement are eligible for inclusion, considered for evaluation, and possibly included in the environmental fate assessment. On the other hand, data or information sources that fail to meet the criteria in the PESO statement are excluded from further consideration.

Assessors seek information on various chemical-specific fate endpoints and associated fate processes, environmental media and exposure pathways as part of the process of developing the environmental fate assessment for each risk evaluation. EPA uses the PESO statement (Table\_Apx A-9) along with the information in Table\_Apx A-10 when screening the fate data or information sources to ensure complete coverage of the processes, pathways and data or information relevant to the environmental fate and transport of the chemical substance undergoing risk evaluation.

#### PESO Evidence Element Environmental fate, transport, partitioning and degradation behavior across environmental media to inform exposure pathways of the chemical substance of interest Exposure pathways included in the conceptual models: air, surface water, groundwater, wastewater, soil, sediment and biosolids. Pathways and Processes associated with the target exposure pathways Processes Bioconcentration and bioaccumulation Destruction and removal by incineration Please refer to the conceptual models for more information about the exposure pathways included in each TSCA risk evaluation. Environmental exposure of environmental receptors (*i.e.*, aquatic and terrestrial organisms) to the chemical substance of interest, mixtures including the chemical substance, and/or its degradation products and metabolites Environmental exposure of human receptors, including any PESS to the chemical substance of **E**xposure interest, mixtures including the chemical substance, and/or its degradation products and metabolites Please refer to the conceptual models for more information about the environmental and human receptors included in each TSCA risk evaluation. Any setting or scenario resulting in releases of the chemical substance of interest into the natural or built environment (e.g., buildings including homes or workplaces, or wastewater treatment Setting or Scenario facilities) that would expose environmental (i.e., aquatic and terrestrial organisms) or human receptors (i.e., general population, and PESS) Fate properties which allow assessments of exposure pathways: Abiotic and biotic degradation rates, mechanisms, pathways, and products Outcomes Bioaccumulation magnitude and metabolism rates Partitioning within and between environmental media (see Pathways and Processes)

## Table\_Apx A-9. Inclusion Criteria for Data or Information Sources Reporting Environmental Fate and Transport Data

## Table\_Apx A-10. Fate Endpoints and Associated Processes, Media and Exposure Pathways Considered in the Development of the Environmental Fate Assessment

Considered in the Develo		Associated Media/Exposure Pathways					
Fate Data Endpoint	Associated Process(es)	Surface Water, Wastewater, Sediment	Soil, Biosolids	Groundwater	Air		
<b>Required Environmental Fa</b>	te Data	Γ		1			
Abiotic reduction rates or half-lives	Abiotic reduction, Abiotic dehalogenation	Х					
Aerobic biodegradation rates or half-lives	Aerobic biodegradation	Х	Х				
Anaerobic biodegradation rates or half-lives	Anaerobic biodegradation	Х	Х	Х			
Aqueous photolysis (direct and indirect) rates or half- lives	Aqueous photolysis (direct and indirect)	Х					
Atmospheric photolysis (direct and indirect) rates or half-lives	Atmospheric photolysis (direct and indirect)				Х		
Bioconcentration factor (BCF), Bioaccumulation factor (BAF)	Bioconcentration, Bioaccumulation	Х	Х		Х		
Biomagnification and related information	Trophic magnification	Х					
Desorption information	Sorption, Mobility	Х	Х	Х			
Destruction and removal by incineration	Incineration				Х		
Hydrolysis rates or half-lives	Hydrolysis	Х	Х	Х			
K <sub>OC</sub> and other sorption information	Sorption, Mobility	Х	Х	Х			
Wastewater treatment removal information	Wastewater treatment	Х	Х				
Supplemental (or Optional)	Environmental Fate Data						
Abiotic transformation products	Hydrolysis, Photolysis, Incineration	Х			Х		
Aerobic biotransformation products	Aerobic biodegradation	Х	X				
Anaerobic biotransformation products	Anaerobic biodegradation	X	X	Х			
Atmospheric deposition information	Atmospheric deposition				Х		
Coagulation information	Coagulation, Mobility	Х		Х			
Incineration removal information	Incineration				Х		

## A.2.1.5 Generation of Hazard Heat Maps

As stated in Appendix A.1.2.2, SWIFT Review has pre-set literature search strategies ("filters") developed by information specialists that can be applied to identify studies that are more likely to be useful for identifying human health and ecotoxicity content. The filters function like a typical search strategy where studies are tagged as belonging to a certain filter if the terms in the filter literature search strategy appear in title, abstract, keyword or MeSH fields content.

After the completion of full-text screening for hazard data, all references tagged as included (or "PECOrelevant) were uploaded to the SWIFT Review tool for further filtering. The SWIFT Review filters applied at this phase focused on types of health outcomes included: "ADME", "PBPK", "cancer", "cardiovascular", "developmental", "endocrine", "gastrointestinal", "hematological and immune", "hepatic", "mortality", "musculoskeletal", "neurological", "nutritional and metabolic", "ocular and sensory", "renal", "reproductive", "respiratory", and "skin and connective tissue". The details of these health outcome search strategies that underlie the filters are available <u>online</u>. Studies that included one or more of the search terms in the title, abstract, keyword, or MeSH fields were exported and used to populate the Hazard Heat Map (Figure 2-10). Studies that were not retrieved using these filters were tagged as "No Tag". The evidence type listed in the heat map (*e.g.*, human, animal-human health model, animal- environmental model, and plant) was manually assigned to each reference by screeners during the full-text screening.

The health outcome tags were originally designed for vertebrate systems, and as such, did not conform well to plant evidence. Therefore, any plant studies tagged for: "cancer", "cardiovascular", "gastrointestinal", "hematological and immune", "hepatic", "musculoskeletal", "neurological", "ocular and sensory" and "renal and respiratory" were manually reviewed and re-tagged to more appropriate health outcomes.


## A.3 Gray Literature Search and Screening Strategies

EPA conducted a gray literature search for available information to support the TSCA risk evaluations for the next twenty TSCA risk evaluations. Gray literature is defined as the broad category of data/information sources not found in standard, peer-reviewed literature databases (*e.g.*, PubMed and Web of Science). Gray literature includes data/information sources such as white papers, conference proceedings, technical reports, reference books, dissertations, information on various stakeholder websites, and other databases. Given the nature of how gray literature is searched and collected, results may not come with a bibliographic citation or abstract and were therefore processed using a decision tree logic described in Appendix A.3.1 for potential relevance prior to entering full text screening where a discipline-specific PECO is applied.

Search terms were variable dependent on source and based on knowledge of a given source to provide discipline-specific information. A summary of sources is provided in Appendix A.3.3. The criteria for determining the potential relevance of documents identified from gray literature sources is described in the following sections for each discipline.

## A.3.1 Screening of Gray Literature

To reduce the overall burden of processing gray literature results, EPA developed a screening process to determine the potential relevance of gray literature. This step was introduced prior to collecting the resulting documents. Figure\_Apx A-1 describes the decision logic used to screen gray literature results.





#### A.3.2 Initial Screening of Sources using Decision Logic Tree

The purpose of the inclusion/exclusion decision logic tree in Figure\_Apx A-1 is to provide a broad, general screening technique to determine whether each gray literature source should be included and further screened or excluded with no additional screening necessary. The diamonds in the decision tree require analysis by the screener, whereas the rectangular boxes are used to classify the type of source. All the questions used in the decision process are provided in Table\_Apx A-11.

Step	Metric	Questions to consider
1	Potential Relevance	Does the result have information (qualitative or quantitative) related to TSCA risk evaluations? *Apply Discipline relevancy metric
2.1.1	Complete / Available	Is it a secondary data source (assessment, robust summary, TSCA submission databases, etc.)?
2.1.2		Is the document from a peer reviewed/published journal?

Table	Apx	A-11.	Decision	Logic	Tree	Overview
I GOIC_	P		Decision	LUGIC		

Step	Metric	Questions to consider
2.2		Is there an established procedure for data collection, communication, peer review, and/or reporting?
2.2.1	]	Has the data been provided by a US governmental/state source?
2.2.2		Has the data been provided by an international governmental source?
2.3	]	Are these data publicly available/accessible?
2.3.1		Is the source TSCA CBI, proprietary, TSCA or NGO stakeholder submission?
3	Duplicate	Does the result contain any duplicative information found in other sources?

Results of the gray literature search and decision tree process are included in Appendix A.3.4.

## A.3.3 TSCA Submission Searching and Title Screening

EPA screens information submitted under TSCA Sections 4, 5, 8(e), and 8(d), as well as for your information (FYI) submissions. In the gray literature process defined in Appendix A.3.2, EPA considers the databases that contain TSCA submissions to be secondary sources (Step 1.1) because the metadata in the databases are secondary. These databases then advance to Step 2.3.1 and then to Process C. The Process C steps are described here.

EPA first screens the titles using two screeners per title. EPA conducts this step primarily to reduce the number of full studies to be obtained because some studies are available only on microfiche or in long-term storage. Screening is done using the inclusion and exclusion criteria within the relevant PECOs, PESOs or RESOs for each topic area (Appendix A.2.1). EPA excludes interim reports (*e.g.*, interim sacrifices for toxicity studies) and only final reports are further considered. If the title is not clear regarding the document's contents, EPA obtains the full text and advances to the next steps.

After full texts are obtained, EPA reviewed some sources (prior to full-text screening) based on whether they have several factors; primary data, an established procedure for peer review, data collection, communication and/or reporting and are publicly available. Sources that have these factors will move on to full text screening. Other sources will go straight to full text screening using PECO-type criteria without going through this extra step.

EPA may decide to initiate a backwards search on sources that are deemed to have secondary data. In situations where parameters such as procedures for peer review and data collection are unclear, EPA may reach out to the authors to retrieve information to gauge whether the source should be included or excluded. Studies that are not publicly available (such as proprietary or CBI sources) may undergo additional screening steps.

During the full-text screening step, two individuals screen each source according to the PECOs, PESOs and RESOs (Appendix A.2.1).

Results of the TSCA submission search and decision tree process are included in Appendix A.3.4.

## A.3.4 Gray Literature Search Results for 1,1,2-Trichloroethane

Table\_Apx A-12 provides a list of gray literature sources that yielded results for 1,1,2-trichloroethane.

Source Agency	Source Name	Source Type	Source Category	Source Website
ATSDR	ATSDR Tox Profile Updates and Addendums	Other US Agency Resources	Assessment or Related Document	https://www.atsdr.cdc.gov/toxpr ofiles/profilesaddenda.asp
ATSDR	ATSDR Toxicological Profiles (original publication)	Other US Agency Resources	Assessment or Related Document	https://www.atsdr.cdc.gov/toxpr ofiles/index.asp
Australian Government, Department of Health	NICNAS Assessments (human health, Tier I, II or III)	International Resources	Assessment or Related Document	https://www.industrialchemicals .gov.au/chemical- information/search-assessments
CAL EPA	Technical Support Documents for regulations: Cancer Potency Information	Other US Agency Resources	Assessment or Related Document	https://oehha.ca.gov/chemicals
CAL EPA	Technical Support Documents for regulations: Drinking Water Public Health Goals	Other US Agency Resources	Assessment or Related Document	https://oehha.ca.gov/chemicals
CAL EPA	Technical Support Documents for regulations: Proposition 65, Cancer	Other US Agency Resources	Assessment or Related Document	https://oehha.ca.gov/chemicals
CDC	CDC Biomonitoring Tables	Other US Agency Resources	Database	https://www.cdc.gov/exposurere port/index.html/
ECHA	ECHA Documents	International Resources	Assessment or Related Document	https://echa.europa.eu/informati on-on-chemicals
ECHA	Annex XVII To REACH - Conditions of Use	International Resources	Assessment or Related Document	https://echa.europa.eu/substance s-restricted-under-reach
Env Canada	Chemicals at a Glance (fact sheets)	International Resources	Assessment or Related Document	https://www.canada.ca/en/health -canada/services/chemical- substances/fact- sheets/chemicals-glance.html
EPA	OPPT: TSCATS database maintained at SRC (TSCA submissions)	US EPA Resources	Database	

 Table\_Apx A-12. List of Gray Literature Sources for 1,1,2-Trichloroethane

Source Agency	Source Name	Source Type	Source Category	Source Website
EPA	OPPT: Chemview (TSCA submissions - chemical test rule data and substantial risk reports)	US EPA Resources	Database	https://chemview.epa.gov/chem view
EPA	OPPT: CIS (CBI LAN) (TSCA submissions)	US EPA Resources	Database	
EPA	2011 NATA	US EPA Resources	Assessment or Related Document	https://www.epa.gov/aegl/access -acute-exposure-guideline- levels-aegls-values#chemicals
EPA	Office of Air: AQS, Annual	US EPA Resources	Database	https://aqs.epa.gov/aqsweb/airda ta/download_files.html#Annual
EPA	Office of Air: National Emissions Inventory (NEI) - National Emissions Inventory (NEI) Data (2014, 2011, 2008)	US EPA Resources	Database	https://www.epa.gov/air- emissions-inventories/2014- national-emissions-inventory- nei-data
EPA	Office of Water: STORET and WQX	US EPA Resources	Database	https://www.waterqualitydata.us /portal/
EPA	PPRTV Derivation Support Document	US EPA Resources	Assessment or Related Document	https://hhpprtv.ornl.gov/quickvi ew/pprtv_papers.php
EPA	EPA Office of Water: Ambient Water Quality Criteria documents	US EPA Resources	Assessment or Related Document	https://www.epa.gov/wqc
EPA	IRIS Summary	US EPA Resources	Assessment or Related Document	https://cfpub.epa.gov/ncea/iris_d rafts/atoz.cfm?list_type=alpha
EPA	Office of Air: TRI	US EPA Resources	Database	https://www.epa.gov/toxics- release-inventory-tri- program/tri-data-and-tools
EPA	Other EPA: Misc sources	US EPA Resources	General Search	https://www.epa.gov/
EPA	EPA: AP-42	US EPA Resources	Regulatory Document or List	https://www.epa.gov/air- emissions-factors-and- quantification/ap-42- compilation-air-emissions- factors
EPA	Office of Water: Drinking Water Standards Health Effects Support Documents	US EPA Resources	Regulatory Document or List	https://www.epa.gov/eg
EPA	Office of Water: CFRs	US EPA Resources	Regulatory Document or List	https://www.epa.gov/eg

Source Agency	Source Name	Source Type	Source Category	Source Website
EPA	Office of Air: CFRs and Dockets	US EPA Resources	Regulatory Document or List	https://www.epa.gov/stationary- sources-air-pollution
EPA	EPA: Generic Scenario	US EPA Resources	Assessment or Related Document	https://www.epa.gov/tsca- screening-tools/chemsteer- chemical-screening-tool- exposures-and-environmental- releases#genericscenarios
IARC	IARC Monograph	International Resources	Assessment or Related Document	http://monographs.iarc.fr/ENG/ Monographs/PDFs/index.php
ILO	International Chemical Safety Cards (ICSCs)	International Resources	Database	https://www.ilo.org/safework/i nfo/publications/WCMS_1131 34/langen/index.htm
Japan	Japanese Ministry of the Environment Assessments - Environmental Risk Assessments (Class I Designated Chemical Substances Summary Table)	International Resources	Regulatory Document or List	https://www.env.go.jp/en/chemi/ prtr/substances/
KOECT	Kirk-Othmer Encyclopedia of Chemical Technology Journal Article	Other Resource	Encyclopedi a	https://onlinelibrary.wiley.com/ doi/book/10.1002/0471238961
NIOSH	CDC NIOSH - Occupational Health Guideline Documents	Other US Agency Resources	Assessment or Related Document	https://www.cdc.gov/niosh/docs /81-123/default.html
NIOSH	CDC NIOSH - Pocket Guide	Other US Agency Resources	Database	https://www.cdc.gov/niosh/npg/ default.html
NIOSH	CDC NIOSH - Health Hazard Evaluations (HHEs)	Other US Agency Resources	Assessment or Related Document	https://www2a.cdc.gov/hhe/sear ch.asp
NIOSH	CDC NIOSH - Publications and Products	Other US Agency Resources	Assessment or Related Document	https://www2a.cdc.gov/nioshtic- 2/
NTP	Technical Reports	Other US Agency Resources	Assessment or Related Document	https://ntp.niehs.nih.gov/publica tions/reports/index.html?type=T echnical+Report
OECD	OECD SIDS	International Resources	Assessment or Related Document	https://hpvchemicals.oecd.org/ui /Publications.aspx
OECD	OECD: General Site	International Resources	General Search	https://www.oecd.org/

Source Agency	Source Name	Source Type	Source Category	Source Website
OECD	OECD Emission Scenario Documents	International Resources	Assessment or Related Document	http://www.oecd.org/document/ 46/0,2340,en_2649_201185_24 12462_1_1_1_00.html
OSHA	OSHA Chemical Exposure Health Data	Other US Agency Resources	Database	https://www.osha.gov/opengov/ healthsamples.html

## Appendix B PHYSICAL AND CHEMICAL PROPERTIES OF 1,1,2-TRICHLOROETHANE

Table\_Apx B-1 summarizes statistics for the physical and chemical property values identified through systematic review as of June 2020. The "N" column indicates the number of unique primary sources of data for that endpoint. That is, if multiple sources presented equivalent values and cited the same primary source, only one of those was included in these statistics and included in the statistical calculations. All physical and chemical properties that were extracted and evaluated as of June 2020 are presented in the supplemental file *Data Extraction and Data Evaluation Tables for Physical and Chemical Property Studies* (EPA-HQ-OPPT-2018-0421).

Property or Endpoint	N	Unit	Mean	Standard Deviation	Min	Max
Molecular formula	-	-	NA	NA	NA	NA
Molecular weight	-	g/mol	NA	NA	NA	NA
Physical state	5	-	NA	NA	NA	NA
Physical properties	4	-	NA	NA	NA	NA
Melting point	14	°C	-36.2	0.91	-37.4	-35.0
Boiling point	14	°C	113.4	1.22	110	115
Density	8	g/cm <sup>3</sup>	1.44	0.014	1.43	1.47
Vapor pressure	3	mm Hg	21.75	2.38	19	23.25
Vapor density	1		4.21		4.21	4.21
Water solubility	9	mg/L	4735	736.6	3697	6200
Octanol/water partition coefficient (log Kow)	5	-	1.96	0.35	1.43	2.38
Henry's Law constant	2	$atm \cdot m^3/mol$	$8.66  imes 10^{-4}$	$5.94  imes 10^{-5}$	$8.24  imes 10^{-4}$	$9.08  imes 10^{-4}$
Flash point	1	°C	114	-	114	114
Auto flammability	0	°C	-	-	-	-
Viscosity	4	cP	1.34	0.30	1.076	1.69

Table\_Apx B-1. Summary Statistics for Reviewed Physical Properties

Property or Endpoint	N	Unit	Mean	Standard Deviation	Min	Max
Refractive index	5	-	1.46776	0.011878	1.447	1.4773
Dielectric constant	1	-	7.29	-	7.29	7.29

NA = Not applicable

## Appendix C ENVIRONMENTAL FATE AND TRANSORT PROPERTIES OF 1,1,2-TRICHLOROETHANE

Table\_Apx C-1 provides the environmental fate characteristics that EPA identified and considered in developing the scope for 1,1,2-trichloroethane. This information was presented in the *Proposed Designation of 1,1,2-trichloroethane (CASRN 79-00-5) as a High-Priority Substance for Risk Evaluation* (U.S. EPA, 2019d) and may be updated as EPA collects additional information through systematic review methods.

Property or Endpoint	Value <sup>a</sup>	Reference
Direct Photodegradation	Not expected to be susceptible to direct photolysis by sunlight because the chemical structure of 1,1,2-trichloroethane does not contain chromophores that absorb at wavelengths >290 nm	<u>NLM (2018)</u>
Indirect Photodegradation	$\begin{array}{l} t_{1/2} = 54.6 \text{ days (from } \cdot \text{OH rate} \\ \text{constant of } 1.96 \times 10^{-13} \\ \text{cm}^3/\text{molecule-sec and } 12\text{-hour} \\ \text{day with } 1.5 \times 10^6 \cdot \text{OH/cm}^3) \end{array}$	<u>U.S. EPA (2012b)</u>
	$t_{1/2} = 87.2$ days at 295 K, $5 \times 10^5$ ·OH-radical molecules/cm <sup>3</sup>	<u>ECHA (2019)</u>
	$t_{1/2} = 50.2$ days at 339 K, $5 \times 10^5$ ·OH-radical molecules/cm <sup>3</sup>	<u>ECHA (2019)</u>
Hydrolysis	Stable at pH 7; $t_{1/2} = 85$ days pH 9 and 25 °C	<u>OECD (2000)</u>
Biodegradation (Aerobic)	Water: 5% after 28 days measured by GC (Japanese MITI test)	<u>NITE (2010)</u>
	Water: Not readily biodegradable (OECD TG 301C)	<u>OECD (2000)</u>
	Water: 0% after a 24-day modified shake-flask test; 0% after a 25-day river die-away test	<u>NLM (2018)</u> citing <u>Tabak et al.</u> ( <u>1981)</u>
	Soil: $t_{1/2} = 6$ months to 1 year	<u>NLM (2018)</u> citing <u>Rathbun</u> (1998)
Biodegradation (Anaerobic)	Water: 6 days in an unpolluted aquifer (1% organic carbon content);16 years in an unpolluted aquifer (0.001% organic carbon content)	NLM (2018) citing <u>Verschueren</u> (2001)

Table_Apx C-1. Environmental Fate and Trans	sport Properties of 1.1.2-Trichloroethane
Tuble_TipA C Ti Environmental T ate and Tran	sport r roper des or 1,1,2 rriemor occhune

Property or Endpoint	Value <sup>a</sup>	Reference
Wastewater Treatment	27.43% total removal (0.08% by biodegradation, 1.64% by sludge adsorption, and 25.72% by volatilization to air; estimated) <sup>b</sup>	<u>U.S. EPA (2012b)</u>
Bioconcentration Factor	<10 to 17	<u>ATSDR (1989)</u>
	OECD TG 305C (carp): 0.7–2.6 at 0.3 mg/L; 2.9–4.0 at 0.03 mg/L	<u>OECD (2000)</u>
	Carp ( <i>Cyprinus carpio</i> ): 0.7–2.6 at 0.3 ppm; 2.7–6.7 at 0.03 ppm (MITI)	<u>SYKE (2018)</u>
Bioaccumulation Factor	6.9 (estimated) <sup>b</sup>	<u>U.S. EPA (2012b)</u>
Soil Organic Carbon:Water Partition Coefficient (Log K <sub>OC</sub> )	1.9–2.05 ( $K_{OC} = 83-111$ ) in silty clay soil; 2.2–2.32 ( $K_{OC} = 174-209$ ) in sandy loam soil	<u>NLM (2018)</u> citing <u>Gan and</u> <u>Dupont (1989)</u>

<sup>a</sup> Measured unless otherwise noted; <sup>b</sup>EPI Suite<sup>TM</sup> physical property inputs: Log  $K_{OW} = 1.89$ , BP = 113.80 °C, MP = -36.60 °C, VP = 23 mm Hg, WS = 4,590 mg/L, HLC = 0.000824 ·OH = hydroxyl radical; OECD = Organization for Economic Cooperation and Development; TG = test guideline; GC = gas chromatography; MITI = Ministry of International Trade and Industry; BOD = biochemical oxygen demand; HPLC = high performance liquid chromatography

## Appendix D REGULATORY HISTORY

The chemical substance, 1,1,2-trichloroethane, is subject to federal and state laws and regulations in the United States (Table\_Apx D-1 and Table\_Apx D-2). Regulatory actions by other governments, tribes and international agreements applicable to 1,1,2-trichloroethane are listed in Table\_Apx D-3.

## **D.1** Federal Laws and Regulations

Statutes/Regulations	Description of Authority/Regulation	Description of Regulation	
EPA Statutes/Regulation	EPA Statutes/Regulations		
Toxic Substances Control Act (TSCA) – Section 6(b)	EPA is directed to identify high-priority chemical substances for risk evaluation; and conduct risk evaluations on at least 20 high priority substances no later than three and one-half years after the date of enactment of the Frank R. Lautenberg Chemical Safety for the 21st Century Act.	1,1,2-Trichloroethane is one of the 20 chemicals EPA designated as a High-Priority Substance for risk evaluation under TSCA ( <u>84 FR 71924</u> , December 30, 2019). Designation of 1,1,2- trichloroethane as a high- priority substance constitutes the initiation of the risk evaluation on the chemical.	
Toxic Substances Control Act (TSCA) – Section 8(a)	The TSCA Section 8(a) CDR Rule requires manufacturers (including importers) to give EPA basic exposure-related information on the types, quantities and uses of chemical substances produced domestically and imported into the United States.	1,1,2-Trichloroethane manufacturing (including importing), processing and use information is reported under the CDR rule (85 FR 20122, <u>April 2, 2020</u> ).	
Toxic Substances Control Act (TSCA) – Section 8(b)	EPA must compile, keep current and publish a list (the TSCA Inventory) of each chemical substance manufactured (including imported) or processed in the United States.	1,1,2-Trichloroethane was on the initial TSCA Inventory and therefore was not subject to EPA's new chemicals review process under TSCA Section 5 ( <u>60 FR 16309</u> , March 29, 1995).	
Toxic Substances Control Act (TSCA) – Section 8(e)	Manufacturers (including importers), processors, and distributors must immediately notify EPA if they obtain information that supports the conclusion that a chemical substance or mixture presents a substantial risk of injury to health or the environment.	One risk report was received for 1,1,2-trichloroethane (1992) ( <u>U.S. EPA, ChemView.</u> Accessed April 10, 2019).	

Table_Apx D-1. Federal I	Laws and Regulations
--------------------------	----------------------

Statutes/Regulations	Description of Authority/Regulation	<b>Description of Regulation</b>
Toxic Substances Control Act (TSCA) – Section 4	Provides EPA with authority to issue rules, enforceable consent agreements and orders requiring manufacturers (including importers) and processors to test chemical substances and mixtures.	14 chemical data submissions from test rules or enforceable consent agreements were received for 1,1,2- trichloroethane: acute inhalation toxicity with histopathology, reproduction and fertility effects, prenatal developmental toxicity, immunotoxicity, pharmacokinetics, carcinogenicity, and neurotoxicity (1985-2006) ( <u>U.S. EPA, ChemView.</u> Accessed April 10, 2019).
Emergency Planning and Community Right- To-Know Act (EPCRA) – Section 313	Requires annual reporting from facilities in specific industry sectors that employ 10 or more full-time equivalent employees and that manufacture, process or otherwise use a TRI-listed chemical in quantities above threshold levels. A facility that meets reporting requirements must submit a reporting form for each chemical for which it triggered reporting, providing data across a variety of categories, including activities and uses of the chemical, releases and other waste management ( <i>e.g.</i> , quantities recycled, treated, combusted) and pollution prevention activities (under Section 6607 of the Pollution Prevention Act). These data include on- and off-site data as well as multimedia data ( <i>i.e.</i> , air, land and water).	1,1,2-Trichloroethane is a listed substance subject to reporting requirements under <u>40 CFR 372.65</u> effective as of January 01, 1987.
Clean Air Act (CAA) – Section 111(b)	Requires EPA to establish new source performance standards (NSPS) for any category of new or modified stationary sources that EPA determines causes, or contributes significantly to, air pollution, which may reasonably be anticipated to endanger public health or welfare. The standards are based on the degree of emission limitation achievable through the application of the best system of emission reduction (BSER) which (taking into account the cost of achieving reductions and environmental impacts and energy requirements) EPA	1,1,2-Trichloroethane is subject to the NSPS for equipment leaks of volatile organic compounds (VOCs) in the synthetic organic chemicals manufacturing industry for which construction, reconstruction or modification began after January 5, 1981 ( <u>40 CFR Part</u> <u>60, Subparts VV, NNN, and</u> <u>RRR</u> ).

Statutes/Regulations	Description of Authority/Regulation	<b>Description of Regulation</b>
	determines has been adequately demonstrated.	
Clean Air Act (CAA) – Section 112(b)	Defines the original list of 189 hazardous air pollutants (HAPs). Under 112(c) of the CAA, EPA must identify and list source categories that emit HAP and then set emission standards for those listed source categories under CAA Section 112(d). CAA Section 112(b)(3)(A) specifies that any person may petition the Administrator to modify the list of HAP by adding or deleting a substance. Since 1990, EPA has removed two pollutants from the original list leaving 187 at present.	1,1,2-Trichloroethane is listed as a HAP ( <u>42 U.S. Code</u> <u>Section 7412</u> ).
Clean Air Act (CAA) – Section 112(d)	Directs EPA to establish, by rule, NESHAPs for each category or subcategory of listed major sources and area sources of HAPs (listed pursuant to Section 112(c)). For major sources, the standards must require the maximum degree of emission reduction that EPA determines is achievable by each particular source category. This is generally referred to as maximum achievable control technology (MACT). For area sources, the standards must require generally achievable control technology (GACT) though may require MACT.	EPA has established <u>NESHAPs</u> for a number of source categories that emit 1,1,2-trichloroethane to air.
Clean Air Act (CAA) – Section 112(d) and 112(f)	Risk and technology review (RTR) of Section 112(d) national emission standards for hazardous air pollutants (NESHAP). Section 112(f)(2) requires EPA to conduct risk assessments for each source category subject to Section 112(d) NESHAP that require maximum achievable control technology (MACT), and to determine if additional standards are needed to reduce remaining risks. Section 112(d)(6) requires EPA to review and revise the emission standards, as necessary, taking into account developments in practices, processes and control technologies.	EPA has promulgated a number of RTR NESHAP and will do so, as required, for the remaining source categories with NESHAP.
Clean Water Act (CWA) - Section 304(a)(1)	Requires EPA to develop and publish ambient water quality criteria (AWQC) reflecting the latest scientific knowledge on the effects on human health that may be	In 2015, EPA published updated AWQC for 1,1,2- trichloroethane, including a recommendation of 0.55

Statutes/Regulations	Description of Authority/Regulation	<b>Description of Regulation</b>
	expected from the presence of pollutants in any body of water.	$(\mu g/L)$ for "Human Health for the consumption of Water + Organism" and 8.9 ( $\mu g/L$ ) for "Human Health for the consumption of Organism Only" for states and authorized tribes to consider when adopting criteria into their water quality standards ( <u>80 FR</u> <u>36986</u> , June 29, 2015).
Clean Water Act (CWA) – Section 301, 304, 306, 307, and 402	Clean Water Act Section 307(a) establishes a list of toxic pollutants or combination of pollutants under the CWA. The statute specifies a list of families of toxic pollutants also listed in the Code of Federal Regulations at 40 CFR Part 401.15. The "priority pollutants" specified by those families are listed in 40 CFR Part 423 Appendix A. These are pollutants for which best available technology effluent limitations must be established on either a national basis through rules (Sections 301(b), 304(b), 307(b), 306) or on a case-by-case best professional judgement basis in NPDES permits, see Section 402(a)(1)(B). EPA identifies the best available technology that is economically achievable for that industry after considering statutorily prescribed factors and sets regulatory requirements based on the performance of that technology.	Under CWA Section 304, 1,1,2-trichloroethane is included in the list of total toxic organics (TTO) ( <u>40 CFR</u> <u>413.02(i)</u> ).
Safe Drinking Water Act (SDWA) – Section 1412	Requires EPA to publish non-enforceable maximum contaminant level goals (MCLG) for a contaminant for which EPA makes the determination that the contaminant: 1. may have an adverse effect on the health of persons; 2. is known to occur or there is a substantial likelihood that the contaminant will occur in public water systems with a frequency and at levels of public health concern; and 3. in the sole judgement of the Administrator, regulation of the contaminant presents a meaningful opportunity for health risk reductions for persons served by public water systems. When EPA publishes an MCLG, EPA must also promulgate a	1,1,2-Trichloroethane is subject to NPDWR under the SDWA with a MCLG of 0.003 and an enforceable MCL of 0.005 mg/L (Section 1412) (57 FR 31776 July 17, 1992).

Statutes/Regulations	Description of Authority/Regulation	<b>Description of Regulation</b>
	National Primary Drinking Water Regulation (NPDWR) which includes either an enforceable maximum contaminant level (MCL), or a required treatment technique. Public water systems are required to comply with NPDWRs.	
Resource Conservation and Recovery Act (RCRA) – Section 3001	Directs EPA to develop and promulgate criteria for identifying the characteristics of hazardous waste, and for listing hazardous waste, taking into account toxicity, persistence, and degradability in nature, potential for accumulation in tissue and other related factors such as flammability, corrosiveness, and other hazardous characteristics.	1,1,2-Trichloroethane is included on the list of hazardous wastes pursuant to RCRA 3001. RCRA Hazardous Waste Code: U227 (40 CFR 261.33); F002, F024, F025 (40 CFR 261.31)
Comprehensive Environmental Response, Compensation and Liability Act (CERCLA) – Sections 102(a) and 103	Authorizes EPA to promulgate regulations designating as hazardous substances those substances which, when released into the environment, may present substantial danger to the public health or welfare or the environment. EPA must also promulgate regulations establishing the quantity of any hazardous substance the release of which must be reported under Section 103. Section 103 requires persons in charge of vessels or facilities to report to the National Response Center if they have knowledge of a release of a hazardous substance above the reportable quantity threshold.	1,1,2-Trichloroethane is a hazardous substance under CERCLA. Releases of 1,1,2- trichloroethane in excess of 100 pounds must be reported (40 CFR 302.4).
Superfund Amendments and Reauthorization Act (SARA) –	Requires the Agency to revise the hazardous ranking system and update the National Priorities List of hazardous waste sites, increases state and citizen involvement in the superfund program and provides new enforcement authorities and settlement tools.	1,1,2-Trichloroethane is listed on <u>SARA</u> , an amendment to CERCLA and the CERCLA Priority List of Hazardous Substances. This list includes substances most commonly found at facilities on the CERCLA National Priorities List (NPL) that have been deemed to pose the greatest threat to public health.
<b>Other Federal Statutes</b>	Other Federal Statutes/Regulations	

Statutes/Regulations	Description of Authority/Regulation	<b>Description of Regulation</b>
Occupational Safety and Health Act (OSHA)	Requires employers to provide their workers with a place of employment free from recognized hazards to safety and health, such as exposure to toxic chemicals, excessive noise levels, mechanical dangers, heat or cold stress or unsanitary conditions (29 U.S.C Section 651 et seq.). Under the Act, OSHA can issue occupational safety and health standards including such provisions as Permissible Exposure Limits (PELs), exposure monitoring, engineering and administrative control measures, and respiratory protection.	In 1980, OSHA issued occupational safety and health standards for 1,1,2- trichloroethane that included a PEL of 10 ppm or 45 mg/m3 as an 8-hour, TWA (29 CFR 1910.1001).
Federal Food, Drug, and Cosmetic Act (FFDCA)	Provides the FDA with authority to oversee the safety of food, drugs and cosmetics.	The FDA regulates 1,1,2- trichloroethane in bottled water. The maximum permissible level 1,1,2- trichloroethane in bottled water is .005 mg/L ( <u>21 CFR</u> <u>165.110</u> ).
Federal Hazardous Materials Transportation Act (HMTA)	<ul> <li>Section 5103 of the Act directs the Secretary of Transportation to:</li> <li>Designate material (including an explosive, radioactive material, infectious substance, flammable or combustible liquid, solid or gas, toxic, oxidizing or corrosive material, and compressed gas) as hazardous when the Secretary determines that transporting the material in commerce may pose an unreasonable risk to health and safety or property. Issue regulations for the safe transportation, including security, of hazardous material in intrastate, interstate and foreign commerce.</li> </ul>	The Department of Transportation (DOT) has designated 1,1,2- trichloroethane as a hazardous material, and there are special requirements for marking, labeling and transporting it ( <u>49</u> <u>CFR Part 172</u> ).

## **D.2** State Laws and Regulations

State Actions	Description of Action	
State Air Regulations	Allowable Ambient Levels: New Hampshire ( <u>Env-A 1400: Regulated Toxic</u> <u>Air Pollutants</u> ). Rhode Island ( <u>Air Pollution Regulation No. 22</u> )	
State Drinking Water Standards and Guidelines	Arizona (14 Ariz. Admin. <u>Register 2978</u> , August 1, 2008), California (Cal Code Regs. <u>Title 26</u> , § 22-64444), Delaware (Del. Admin. <u>Code Title 16</u> , § 4462), Connecticut (Conn. Agencies Regs. § <u>19-13-B102</u> ), Florida (Fla. <u>Admin. Code R. Chap. 62-550</u> ), Maine ( <u>10 144 Me. Code R. Chap. 231</u> ), Massachusetts ( <u>310 Code Mass. <u>Regs.</u> § 22.00</u> ), Michigan (Mich. Admin. <u>Code r.299.44 and r.299.49</u> , 2017), Minnesota (Minn R. <u>Chap. 4720</u> ), New Jersey (7:10 N.J <u>Admin. Code § 5.2</u> ), Pennsylvania ( <u>25 Pa. Code §</u> <u>109.202</u> ), Rhode Island (Rules and Regulations Pertaining to Public Drinking Water <u>R46-13-DWQ</u> ), Texas ( <u>30 Tex. Admin. <u>Code § 290.104</u>).</u>	
State PELs	California (PEL of 10 ppm) (Cal Code Regs. <u>Title 8, § 5155</u> ) Hawaii PEL: 10 ppm ( <u>Hawaii Administrative Rules Section 12-60-50</u> ).	
State Right-to-Know Acts	New Jersey ( <u>N.J.A.C. 7:1G</u> ) and Pennsylvania (P.L. 734, No. 159 and <u>34</u> <u>Pa. Code § 323</u> ).	
Chemicals of High Concern to Children	Several states have adopted reporting laws for chemicals in children's products containing 1,1,2-trichloroethane, including Maine ( <u>38 MRSA</u> <u>Chapter 16-D</u> ), Minnesota ( <u>Toxic Free Kids Act Minn. Stat. 116.9401 to 116.9407</u> ).	
Other	California listed 1,1,2-trichloroethane on Proposition 65 in October 1, 1990 due to cancer. ( <u>Cal Code Regs. Title 27, § 27001</u> ). 1,1,2-Trichloroethane is listed as a Candidate Chemical under California's Safer Consumer Products Program (Health and Safety <u>Code § 25252 and 25253</u> ). California issued a Health Hazard Alert for 1,1,2-trichloroethane ( <u>Hazard</u>	
	Evaluation System and Information Service, 2016). California lists 1,1,2-trichloroethane as a designated priority chemical for biomonitoring ( <u>California SB 1379</u> ).	
	1,1,2-Trichloroethane is on the MA Toxic Use Reduction Act (TURA) list of 2019 ( <u>301 CMR 41.03</u> ).	

Table\_Apx D-2. State Laws and Regulations

## **D.3** International Laws and Regulations

Country/Tribe/ Organization	Requirements and Restrictions
Canada	<ul> <li>1,1,2-Trichloroethane is on the Domestic Substances List (Government of Canada. Managing substances in the environment. Substances search.</li> <li><u>Database</u> accessed April 17, 2019). And a Notice or Order has been published indicating that the Significant New Activity (SNA) provisions of <u>CEPA</u> apply to this substance.</li> <li>Other regulations include:</li> <li>Canada's National Pollutant Release Inventory (NPRI).</li> </ul>
European Union	In 2010, a restriction of sale and use 1,1,2-Trichloroethane as a substance or constituent of other substances, or in mixtures in concentrations equal to or greater than 0.1% or was added to <u>Annex XVII</u> of regulation (EC) No 552/2009- REACH (Registration, Evaluation, Authorization and Restriction of Chemicals). The restriction included a provision requiring that products be marked 'For use in industrial installations only.' The provision does not apply to medicinal or veterinary productos, or cosmetic products (European Chemicals Agency ( <u>ECHA</u> ) database. Accessed April 15, 2019).
Australia	1,1,2-Trichloroethane was assessed under Human Health II of the Inventory Multi-Tiered Assessment and Prioritisation (IMAP). There were no uses reported in Australia. International uses include adhesives, lacquer and coating formulations; and aerosol paints; as a solvent in the manufacture of organic materials (fats, oils, waxes, resins); as an intermediate in the production of vinylidene chloride (1,1-dichloroethylene) and tetrachloroethanes. ( <u>NICNAS</u> , 2013, Ethane, 1,1,2-trichloro-: Human health tier II assessment. Accessed April 15, 2019).
Japan	<ul> <li>1,1,2-Trichloroethane is regulated in Japan under the following legislation:</li> <li>Act on the Evaluation of Chemical Substances and Regulation of Their Manufacture, etc. (Chemical Substances Control Law; CSCL)</li> <li>Act on Confirmation, etc. of Release Amounts of Specific Chemical Substances in the Environment and Promotion of Improvements to the Management Thereof</li> <li>Industrial Safety and Health Act (ISHA)</li> <li>Air Pollution Control Law</li> <li>Water Pollution Control Law</li> <li>Soil Contamination Countermeasures Act (National Institute of Technology and Evaluation [NITE] Chemical Risk Information Platform [CHRIP], Accessed April 16, 2019).</li> </ul>
Australia, Austria, Belgium, Canada, Denmark, Finland, France, Germany, Japan, New	Occupational exposure limits for 1,1,2-Trichloroethane ( <u>GESTIS</u> International limit values for chemical agents (Occupational exposure limits, OELs) database. Accessed April 17, 2019).

## Table\_Apx D-3. Regulatory Actions by other Governments, Tribes, and International Agreements

Country/Tribe/ Organization	<b>Requirements and Restrictions</b>
Zealand, Poland, Singapore, South Korea, Spain, Switzerland	

# Appendix E PROCESS, RELEASE AND OCCUPATIONAL EXPOSURE INFORMATION

This appendix provides information and data found in preliminary data gathering for 1,1,2-trichloroethane.

## E.1 Process Information

Process-related information potentially relevant to the risk evaluation may include process diagrams, descriptions and equipment. Such information may inform potential release sources and worker exposure activities.

## E.1.1 Manufacture (Including Import)

## E.1.1.1 Manufacture

1,1,2-Trichloroethane is produced in the USA by the chlorination of ethylene (<u>IARC</u>, <u>1991</u>) In a twostage manufacturing process, this initially yields 1,2-dichloroethane (<u>IARC</u>, <u>1991</u>); subsequent chlorination yields 1,1,2-trichloroethane and hydrochloric acid. In an alternative production method, ethylene is combined with hydrochloric acid and oxygen at 280–370°C on a catalyst of cupric chloride and potassium chloride to yield 1,1,2-trichloroethane and other chlorinated ethanes (<u>Reed et al.</u>, <u>1988</u>). 1,1,2-Trichloroethane has also been made by the chlorination of vinyl chloride (<u>IARC</u>, <u>1991</u>) in a liquid phase at 300–320°C (<u>Thomas et al.</u>, <u>1982</u>).

## E.1.1.2 Import

Chemicals such as 1,1,2-trichloroethane may be imported into the United States in bulk via water, air, land, and intermodal shipments (Tomer and Kane, 2015). These shipments take the form of oceangoing chemical tankers, railcars, tank trucks, and intermodal tank containers. Chemicals shipped in bulk containers may be repackaged into smaller containers for resale, such as drums or bottles. Domestically manufactured commodity chemicals may be shipped within the United States in liquid cargo barges, railcars, tank trucks, tank containers, intermediate bulk containers (IBCs)/totes, and drums. Both imported and domestically manufactured commodity chemicals may be repackaged by wholesalers for resale; for example, repackaging bulk packaging into drums or bottles. The type and size of container will vary depending on customer requirement. In some cases, QC samples may be taken at import and repackaging sites for analyses. Some import facilities may only serve as storage and distribution locations, and repackaging/sampling may not occur at all import facilities.

1,1,2-Trichloroethane may be imported neat or as a component in a formulation. In the 2016 CDR, one company reported importing a formulation containing less than 1 percent 1,1,2-Trichloroethane.

## E.1.2 Processing and Distribution

## E.1.2.1 Processing as a Reactant or Intermediate

Processing as a reactant or intermediate is the use of 1,1,2-trichloroethane as a feedstock in the production of another chemical via a chemical reaction in which 1,1,2-trichloroethane is consumed to form the product. In the 2016 CDR, companies reported use of 1,1,2-trichloroethane as an intermediate in the manufacture of plastics, petrochemicals, and other chemical products. EPA has not identified specific process information for the processing of 1,1,2-trichloroethane as a reactant at this time.

### E.1.2.2 Recycling

EPA did not identify 1,1,2-trichloroethane specific information for recycling; however, at least one company reported to CDR that 1,1,2-trichloroethane is recycled (<u>U.S. EPA, 2019a</u>). EPA plans to further investigate this condition of use during the risk evaluation.

#### E.1.3 Industrial and Commercial Uses

### E.1.3.1 Industrial Use: Adhesives and Sealants

According to public comments provided by the Aerospace Industries Association (AIA), 1,1,2-trichloroethane is used in the aerospace sector. Specifically, it is a component of proprietary acrylic and methacrylate adhesive formulations used by civil aviation and Department of Defense customers (EPA-HQ-OPPT-2018-0421-0006).

### E.1.3.2 Commercial Use: Adhesives and Sealants

1,1,2-Trichloroethane is present in two-component adhesive products at a concentration of less than 0.1 percent (EPA-HQ-OPPT-2018-0421-0003) (Gorilla Glue Company, 2018). Product instructions indicate that the two components should be dispensed on to a disposable surface, mixed until uniform, and then applied to substrates for bonding. Full curing occurs in 24 hours (Gorilla Glue Company, 2020). A commenter (EPA-HQ-OPPT-2018-0131-0042) provided descriptions of their use of 1,1,2-trichloroethane as a component of common off the shelf adhesives, further informing EPA's understanding of this condition of use.

### E.1.3.3 Commercial Use: Laboratory Use

A Safety Data Sheet for 1,1,2-trichloroethane (98% percent purity) indicates recommended use as a laboratory chemical. However, specific laboratory use activities are unknown (<u>ThermoFisher Scientific</u>, 2018). A commenter (<u>EPA-HQ-OPPT-2018-0131-0042</u>) provided descriptions of their use of 1,1,2-trichloroethane in analytical standards, research, equipment calibration, and sample preparation applications, including as a reference sample for analysis of terrestrial and extraterrestrial material samples, which the commenter also indicated was a critical use, further informing EPA's understanding of this condition of use.

#### E.1.4 Disposal

Each of the conditions of use of 1,1,2-trichloroethane may generate waste streams of the chemical that are collected and transported to third-party sites for disposal, treatment, or recycling. Industrial sites that treat or dispose onsite wastes that they themselves generate will be assessed in each condition of use assessment, as appropriate. Similarly, point source discharges of 1,1,2-trichloroethane to surface water will be assessed in each condition of use. Wastes of 1,1,2-trichloroethane that are generated during a condition of use and sent to a third-party site for treatment, disposal, or recycling may include wastewater and solid wastes. 1,1,2-Trichloroethane may be contained in wastewater discharged to POTW or other, non-public treatment works for treatment. Industrial wastewater containing 1,1,2-trichloroethane discharged to a POTW may be subject to EPA or authorized NPDES state pretreatment programs. The assessment of wastewater discharges to POTWs and non-public treatment works of 1,1,2-trichloroethane will be evaluated in each of the condition of use.

Solid wastes are defined under RCRA as any material that is discarded by being: abandoned; inherently waste-like; a discarded military munition; or recycled in certain ways (certain instances of the generation and legitimate reclamation of secondary materials are exempted as solid wastes under RCRA). The conceptual model depicted in Figure 2-16 presents the exposure pathways, exposure routes and hazards

to general population and environmental receptors from releases and wastes from industrial, commercial, and consumer uses of 1,1,2-trichloroethane that EPA plans to evaluate.

## E.2 Preliminary Occupational Exposure Data

EPA presents below examples of occupational exposure-related information from the preliminary data gathering. EPA plans to consider this information and data in combination of other data and methods for use in the risk evaluation. Note there are no OSHA Chemical Exposure and Health Data (CEHD) or NIOSH Health Hazard Evaluations for 1,1,2-trichloroethane within the last ten years.

Table_Apx E-1. Summary of Industry Sectors with 1,1,2-Trichloroethane Personal Monitoring
Air Samples Obtained from OSHA Inspections Conducted since 1984

SIC Code	SIC Description	Number of Data Points				
1794	Excavation Work	4				
2284	Thread Mills	1				
2752	Commercial Printing, Lithographic	7				
2869	Industrial Organic Chemicals, Not Elsewhere Classified	22				
3089	Plastics Products, Not Elsewhere Classified	7				
3312	Steel Works, Blast Eurnaces (Including Coke Ovens) and Rolling					
3531	Construction Machinery and Equipment	3				
3559	Special Industry Machinery, Not Elsewhere Classified	2				
3561	Pumps and Pumping Equipment	4				
3672	Printed Circuit Boards	2				
3713	Truck and Bus Bodies	1				
3961	Costume Jewelry	3				
7211	Power Laundries, Family and Commercial	32				
8748	Business Consulting Services, Not Elsewhere Classified	2				
9711	National Security	39				

Source: https://www.osha.gov/opengov/healthsamples.html.

Table\_Apx E-2. Potentially Relevant Data Sources for Exposure Monitoring and Area Monitoring Data from NIOSH Health Hazard Evaluations (<u>https://www.cdc.gov/niosh/hhe/</u>)

Year of Publication	Report Number	Facility Description				
1973	Not available	Appliance manufacturing				
1977	HHE-76-42-407	Production of precision parts for aircraft and missiles				
1977	HHE-77-86-439	Production of friction igniters				
1986 HETA-84-214-1633		Electronic material and flexible printed circuit manufacturing				
1988	HETA-87-181-0000	Production of microwave popcorn bags				

# Appendix F SUPPORTING INFORMATION - CONCEPTUAL MODEL FOR INDUSTRIAL AND COMMERCIAL ACTIVITIES AND USES

Table_Apx F-1. Worker	and Occupational Non-	-User Exposure	<b>Conceptual Mod</b>	el Supporting Table

Life Cycle Stage	Category	Subcategory	Release / Exposure Scenario	Exposure Pathway	Exposure Route	Receptor / Population	Plans to Evaluate?	Rationale					
				Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,1,2-trichloroethane.					
	Domostia		Manufacture of	Vapor	Inhalation	Worker	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.					
	Domestic manufacture Domestic manufacture	Domestic manufacture	1,1,2- trichloroethane	<i>, ,</i>	· · ·	, ,	, ,	, ,	Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
Manufacturing				Vapor	Inhalation	ONU	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.					
Manufacturing		Import	Repackaging of	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,1,2- trichloroethane.					
				Vapor	Inhalation	Worker	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.					
	Import		import containers	Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.					
				Vapor	Inhalation	ONU	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.					

Life Cycle Stage	Category	Subcategory	Release / Exposure Scenario	Exposure Pathway	Exposure Route	Receptor / Population	Plans to Evaluate?	Rationale
		Intermediate in plastic manufacturing		Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,1,2- trichloroethane.
	Processing as a	Intermediate in all other	Manufacture of plastics,	Vapor	Inhalation	Worker	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
	reactant or intermediate	chemical product and preparation manufacturing	petrochemicals, and other chemical products	Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
Processing	manufact	Petrochemical manufacturing		Vapor	Inhalation	ONU	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Trocessing		cling Recycling	Recycling of 1,1,2- trichloroethane or products containing 1,1,2- trichloroethane	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,1,2- trichloroethane.
				Vapor	Inhalation	Worker	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
	Recycling			Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Distribution in commerce	Distribution in commerce	Distribution in commerce	Distribution of bulk shipments of 1,1,2- trichloroethane and formulated products	Liquid Contact, Vapor	Dermal, Inhalation	Worker, ONU	Yes	EPA plans to analyze activities resulting in exposures associated with distribution in commerce ( <i>e.g.</i> , loading, unloading) throughout the various lifecycle stages and conditions of use ( <i>e.g.</i> , manufacturing, processing, industrial use, commercial use, disposal) rather than as a single distribution scenario.

Life Cycle Stage	Category	Subcategory	Release / Exposure Scenario	Exposure Pathway	Exposure Route	Receptor / Population	Plans to Evaluate?	Rationale										
				Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,1,2- trichloroethane.										
				Vapor	Inhalation	Worker	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.										
	Adhesives and		Use of adhesives	Mist	Inhalation/ Dermal	Worker	Yes	Method of application for this type of adhesive is not known. If the adhesive is spray applied, there is potential for mist generation and associated exposure.										
Industrial Use	sealants	Adhesives and sealants	Adhesives and sealants in aerosp industry			in aerospace industry								Mist	Inhalation	ONU	Yes	Method of application for this type of adhesive is not known. If the adhesive is spray applied, there is potential for mist generation and associated exposure.
				Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.										
				Vapor	Inhalation	ONU	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.										
		Impurity in adhesives Two-component glues and adhesives	Application of glues and adhesives	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,1,2- trichloroethane. The chemical may be present in adhesives at up to 5% concentration. In some cases, it is present as only an impurity.										
Commercial Use	Adhesives and sealants			Vapor	Inhalation	Worker/ONU	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.										
				Mist	Inhalation, Dermal	Worker, ONU	No	Known adhesive products are supplied in tubes and similar packaging. Because the product will not be spray-applied, mist generation is expected to be negligible.										

Life Cycle Stage	Category	Subcategory	Release / Exposure Scenario	Exposure Pathway	Exposure Route	Receptor / Population	Plans to Evaluate?	Rationale
				Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,1,2-trichloroethane.
	Other use	Laboratory chemical	Laboratory chemical use (Use Report	Vapor	Inhalation	Worker	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
			indicates this as industrial use)	Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
		Disposal	Worker handling of wastes	Liquid Contact	Dermal	Worker	Yes	Workers are expected to routinely handle liquids containing 1,1,2-trichloroethane.
				Vapor	Inhalation	Worker	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.
Disposal	Disposal			Liquid Contact	Dermal	ONU	No	Dermal exposure by ONU is not expected for this condition of use as they are not expected to directly handle the chemical.
				Vapor	Inhalation	ONU	Yes	Due to high volatility (23 mmHg at room temperature), EPA plans to evaluate inhalation exposure to vapor.

## Appendix G SUPPORTING INFORMATION - CONCEPTUAL MODEL FOR CONSUMER ACTIVITIES AND USES

Life Cycle Stage	Category	Subcategory	Release from source	Exposure Pathway	Route	Receptor	Proposed for Further Analysis	Rationale
Consumer Use;	Adhesives and	Adhesives and	Application of glues and adhesives	Liquid	Dermal	Consumers and Bystanders	Yes	Consumers can potentially handle liquids containing 1,1,2-trichloroethane. The chemical may be present in adhesives at up to 5% concentration. In some cases, it is present only as an impurity.
Consumer Reuse and Recycling	Sealants	Sealants		•	Vapor	Inhalation	Consumers and Bystanders	Yes
				Mist	Inhalation/ Dermal	Consumers and Bystanders	No	Known adhesive products are supplied in tubes and similar packaging. Because the product will not be spray-applied, mist generation is expected to be negligible.

## Table\_Apx G-1. Consumer Exposure Conceptual Model Supporting Table

## Appendix H SUPPORTING INFORMATION – CONCEPTUAL MODEL FOR ENVIRONMENTAL RELEASES AND WASTES

## Table\_Apx H-1. General Population and Environmental Exposure Conceptual Model Supporting Table

Life Cycle Stage	Category	Release	Exposure Pathway / Media	Exposure Routes	Receptor / Population	Plans to Evaluate	Rationale
			Near facility ambient air concentrations	Inhalation	General Population	No	1,1,2-Trichloroethane is a HAP.
	Emissions to Air	Emissions to Air	Indirect deposition to	Oral Dermal	General Population	No	Stationary source releases of 1,1,2- trichloroethane to ambient air are under
			nearby bodies of water and soil catchments	TBD	Aquatic and Terrestrial Receptors	No	the jurisdiction of the CAA
			Direct release into surface water and	TBD	Aquatic Receptors	Yes	EPA has developed Ambient Water
All			indirect partitioning to sediment	Oral Dermal	General Population	No	Quality Criteria for protection of human health for 1,1,2-trichloroethane.
	Wastewater or Liquid	treatment or POTW	treatment and wastewater Via Surface or	Oral Dermal and Inhalation ( <i>e.g.</i> , showering)	General Population	No	The drinking water exposure pathway for 1,1,2-trichloroethane is currently addressed in the SDWA regulatory analytical process for public water systems
	-			Oral ( <i>e.g.</i> , ingestion of soil) Inhalation	General Population	No	Unlikely to be a route to general population. CWA (AWQC) and SDWA (NPDWR) cover ambient and drinking water.
			and/or surface water	TBD	Aquatic Receptors	Yes	
		Underground injection	Migration to groundwater, potential	Oral Dermal Inhalation	General Population	No	1,1,2-trichloroethane is released to Class I Underground Injection Wells which are covered by SDWA and RCRA.

Life Cycle Stage	Category	Release	Exposure Pathway / Media	Exposure Routes	Receptor / Population	Plans to Evaluate	Rationale
			surface/drinking water	TBD	Aquatic and Terrestrial Species		
	Solid and	Municipal landfill	Leachate to soil, ground	Oral Dermal	General Population	No	1,1,2-trichloroethane is included on the list of hazardous wastes pursuant to RCRA 3001 (40 CFR §§ 261.33).
Disposal	Liquid Wastes	Municipal landfill and other land disposal	water and/or mitigation to surface water	TBD	Aquatic and Terrestrial Receptors		