

2. Trends in Greenhouse Gas Emissions


2.1 Trends in U.S. Greenhouse Gas Emissions and Sinks

In 2019, total gross U.S. greenhouse gas emissions were 6,577.2 million metric tons carbon dioxide equivalent (MMT CO₂ Eq).¹ Total U.S. emissions have increased by 2.0 percent from 1990 to 2019, down from a high of 15.7 percent above 1990 levels in 2007. Emissions decreased from 2018 to 2019 by 1.7 percent (116.0 MMT CO₂ Eq.). Net emissions (i.e., including sinks) were 5,788.3 MMT CO₂ Eq. Overall, net emissions decreased 1.8 percent from 2018 to 2019 and decreased 12.9 percent from 2005 levels, as shown in Table 2-1. The decline reflects many long-term trends, including population, economic growth, energy market trends, technological changes including energy efficiency, and carbon intensity of energy fuel choices. Between 2018 and 2019, the decrease in total greenhouse gas emissions was driven largely by a decrease in CO₂ emissions from fossil fuel combustion. The decrease in CO₂ emissions from fossil fuel combustion was a result of a 1.3 percent decrease in total energy use and reflects a continued shift from coal to less carbon intensive natural gas and renewables.


Since 1990, U.S. emissions have increased at an average annual rate of 0.1 percent. Figure 2-1 through Figure 2-3 illustrate the overall trend in total U.S. emissions by gas, annual changes, and relative changes since 1990.

¹ The gross emissions total presented in this report for the United States excludes emissions and sinks from removals from Land Use, Land-Use Change, and Forestry (LULUCF). The net emissions total presented in this report for the United States includes emissions and sinks from removals from LULUCF.

1 **Figure 2-1: U.S. Greenhouse Gas Emissions by Gas**




2
3 **Figure 2-2: Annual Percent Change in Gross U.S. Greenhouse Gas Emissions Relative to the**
4 **Previous Year**



5

1 **Figure 2-3: Change in Annual Gross U.S. Greenhouse Gas Emissions Relative to 1990**
 2 **(1990=0)**



3

4 Overall, from 1990 to 2019, total emissions of CO₂ increased by 157.9 MMT CO₂ Eq. (3.1 percent), while total
 5 emissions of methane (CH₄) decreased by 119.9 MMT CO₂ Eq. (15.4 percent), and total emissions of nitrous oxide
 6 (N₂O) remained constant despite fluctuations throughout the time series. During the same period, aggregate
 7 weighted emissions of hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), sulfur hexafluoride (SF₆), and nitrogen
 8 trifluoride (NF₃) rose by 85.2 MMT CO₂ Eq. (85.5 percent). Despite being emitted in smaller quantities relative to
 9 the other principal greenhouse gases, emissions of HFCs, PFCs, SF₆, and NF₃ are significant because many of them
 10 have extremely high global warming potentials (GWPs), and, in the cases of PFCs, SF₆, and NF₃, long atmospheric
 11 lifetimes. Conversely, U.S. greenhouse gas emissions were partly offset by carbon (C) sequestration in managed
 12 forests, trees in urban areas, agricultural soils, landfilled yard trimmings, and coastal wetlands. These were
 13 estimated to offset 12.4 percent (812.4 MMT CO₂ Eq.) of total emissions in 2019.

14 Table 2-1 summarizes emissions and sinks from all U.S. anthropogenic sources in weighted units of MMT CO₂ Eq.,
 15 while unweighted gas emissions and sinks in kilotons (kt) are provided in Table 2-2.

16 **Table 2-1: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (MMT CO₂ Eq.)**

Gas/Source	1990	2005	2015	2016	2017	2018	2019
CO₂	5,116.6	6,141.0	5,384.2	5,262.6	5,219.4	5,397.3	5,274.4
Fossil Fuel Combustion	4,731.5	5,753.8	5,013.8	4,922.0	4,870.4	5,023.1	4,888.5
<i>Transportation</i>	1,469.1	1,868.8	1,750.5	1,797.3	1,812.6	1,821.2	1,843.2
<i>Electric Power</i>	1,820.0	2,400.1	1,900.6	1,808.9	1,732.0	1,752.9	1,606.0
<i>Industrial</i>	853.8	844.0	790.3	787.0	794.1	838.0	837.6
<i>Residential</i>	338.6	359.1	318.1	292.3	294.6	339.5	338.8
<i>Commercial</i>	228.3	226.0	224.9	210.4	212.3	246.9	238.3
<i>U.S. Territories</i>	21.7	56.0	29.4	26.2	24.7	24.6	24.6
Non-Energy Use of Fuels	112.8	130.6	105.3	93.6	99.7	110.9	110.1
Petroleum Systems	9.7	12.1	32.4	21.9	24.8	36.6	43.3
Iron and Steel Production & Metallurgical Coke Production	104.7	70.1	47.9	43.6	40.6	42.6	41.3
Cement Production	33.5	46.2	39.9	39.4	40.3	39.0	40.9
Natural Gas Systems	32.0	25.2	29.2	30.1	31.2	33.9	36.9
Petrochemical Production	21.6	27.4	28.1	28.3	28.9	29.3	30.8
Incineration of Waste	11.2	17.4	21.5	21.8	21.3	20.9	20.2
Lime Production	11.7	14.6	13.3	12.5	12.9	13.1	13.0
Ammonia Production	13.0	9.2	10.6	10.2	11.1	12.2	12.3
Other Process Uses of Carbonates	6.3	7.6	12.2	11.0	9.9	7.5	7.5

Urea Consumption for Non-Agricultural Purposes	3.8	3.7	4.6	5.1	5.0	6.1	6.6
Urea Fertilization	2.4	3.5	4.7	4.9	5.1	5.2	5.3
Carbon Dioxide Consumption	1.5	1.4	4.9	4.6	4.6	4.1	4.9
Liming	4.7	4.3	3.7	3.1	3.1	2.2	2.4
Aluminum Production	6.8	4.1	2.8	1.3	1.2	1.5	1.9
Soda Ash Production	1.4	1.7	1.7	1.7	1.8	1.7	1.8
Ferroalloy Production	2.2	1.4	2.0	1.8	2.0	2.1	1.6
Titanium Dioxide Production	1.2	1.8	1.6	1.7	1.7	1.5	1.5
Glass Production	1.5	1.9	1.3	1.2	1.3	1.3	1.3
Zinc Production	0.6	1.0	0.9	0.9	1.0	1.0	1.0
Phosphoric Acid Production	1.5	1.3	1.0	1.0	1.0	0.9	0.9
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.2
Abandoned Oil and Gas Wells	+	+	+	+	+	+	+
Magnesium Production and Processing	+	+	+	+	+	+	+
<i>Wood Biomass, Ethanol, and Biodiesel Consumption^a</i>	219.4	230.7	317.7	316.6	312.3	319.6	316.2
<i>International Bunker Fuels^b</i>	103.5	113.2	110.8	116.5	120.0	122.0	116.1
CH₄^c	780.0	688.1	652.1	643.5	649.2	655.9	660.1
Enteric Fermentation	164.7	169.3	166.9	172.2	175.8	178.0	178.6
Natural Gas Systems	187.7	163.7	148.6	146.7	147.6	150.8	156.4
Landfills	179.6	131.4	111.4	108.0	109.4	112.1	114.5
Manure Management	37.1	51.6	57.9	59.6	59.9	61.7	62.4
Coal Mining	96.5	65.7	61.8	54.3	55.4	53.4	47.4
Petroleum Systems	48.5	40.6	43.0	40.8	40.9	38.5	40.2
Wastewater Treatment	20.2	20.1	18.8	18.7	18.5	18.4	18.4
Rice Cultivation	16.0	18.0	16.2	15.8	14.9	15.6	15.1
Stationary Combustion	8.6	7.8	8.4	7.8	7.6	8.6	8.6
Abandoned Oil and Gas Wells	6.6	7.0	7.1	7.2	7.1	7.0	7.0
Abandoned Underground Coal Mines	7.2	6.6	6.4	6.7	6.4	6.2	5.9
Mobile Combustion	6.4	3.9	2.6	2.5	2.5	2.4	2.4
Composting	0.4	1.9	2.1	2.3	2.4	2.3	2.3
Field Burning of Agricultural Residues	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Petrochemical Production	0.2	0.1	0.2	0.2	0.3	0.3	0.3
Anaerobic Digestion at Biogas Facilities	+	0.1	0.2	0.2	0.2	0.2	0.2
Ferroalloy Production	+	+	+	+	+	+	+
Carbide Production and Consumption	+	+	+	+	+	+	+
Iron and Steel Production & Metallurgical Coke Production	+	+	+	+	+	+	+
Incineration of Waste	+	+	+	+	+	+	+
<i>International Bunker Fuels^b</i>	0.2	0.1	0.1	0.1	0.1	0.1	0.1
N₂O^c	453.1	456.2	468.4	451.1	446.7	459.7	457.8
Agricultural Soil Management	315.9	313.4	348.5	330.1	327.6	338.2	344.6
Wastewater Treatment	18.7	23.0	25.4	25.9	26.4	26.1	26.4
Stationary Combustion	25.1	34.3	30.4	29.9	28.4	28.2	24.9
Manure Management	14.0	16.4	17.5	18.1	18.7	19.4	19.6
Mobile Combustion	45.2	42.0	21.9	21.1	20.2	19.1	18.6
Nitric Acid Production	12.1	11.3	11.6	10.1	9.3	9.6	10.0
Adipic Acid Production	15.2	7.1	4.3	7.0	7.4	10.3	5.3

N ₂ O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Composting	0.3	1.7	1.9	2.0	2.2	2.0	2.0
Caprolactam, Glyoxal, and Glyoxylic Acid Production	1.7	2.1	1.9	1.7	1.5	1.4	1.4
Incineration of Waste	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Electronics Industry	+	0.1	0.2	0.2	0.3	0.3	0.2
Field Burning of Agricultural Residues	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Petroleum Systems	+	+	+	+	+	+	+
Natural Gas Systems	+	+	+	+	+	+	+
<i>International Bunker Fuels^b</i>	0.9	1.0	1.0	1.0	1.1	1.1	1.0
HFCs	46.5	127.5	168.3	168.1	170.1	169.3	173.8
Substitution of Ozone Depleting Substances ^d	0.2	107.3	163.6	164.9	164.5	165.5	169.7
HCFC-22 Production	46.1	20.0	4.3	2.8	5.2	3.3	3.7
Electronics Industry	0.2	0.2	0.3	0.3	0.4	0.4	0.3
Magnesium Production and Processing	+	+	0.1	0.1	0.1	0.1	0.1
PFCs	24.3	6.7	5.2	4.4	4.1	4.7	4.5
Electronics Industry	2.8	3.3	3.1	2.9	2.9	3.0	2.7
Aluminum Production	21.5	3.4	2.1	1.4	1.1	1.6	1.8
Substitution of Ozone Depleting Substances ^d	0.0	+	+	+	+	0.1	0.1
SF₆	28.8	11.8	5.5	6.0	5.9	5.8	6.0
Electrical Transmission and Distribution	23.2	8.4	3.8	4.1	4.2	3.9	4.2
Magnesium Production and Processing	5.2	2.7	1.0	1.1	1.1	1.1	1.0
Electronics Industry	0.5	0.7	0.7	0.8	0.7	0.8	0.8
NF₃	+	0.5	0.6	0.6	0.6	0.6	0.6
Electronics Industry	+	0.5	0.6	0.6	0.6	0.6	0.6
Unspecified Mix of HFCs, PFCs, SF₆, and NF₃	+						
Electronics Industry	+	+	+	+	+	+	+
Total Emissions (Sources)	6,449.4	7,431.9	6,684.3	6,536.3	6,496.0	6,693.2	6,577.2
LULUCF Emissions (Sources)^c	7.9	16.8	27.9	13.2	26.0	23.4	23.4
LULUCF CH ₄ Emissions	4.9	9.3	16.6	7.7	15.3	13.8	13.8
LULUCF N ₂ O Emissions	3.0	7.5	11.3	5.5	10.6	9.7	9.7
LULUCF Carbon Stock Change^e	(908.7)	(804.8)	(791.5)	(855.8)	(791.8)	(824.6)	(812.4)
LULUCF Sector Net Total^f	(900.8)	(788.0)	(763.6)	(842.6)	(765.8)	(801.2)	(788.9)
Net Emissions (Sources and Sinks)	5,548.6	6,643.8	5,920.7	5,693.8	5,730.2	5,892.0	5,788.3

Notes: Total emissions presented without LULUCF. Net emissions presented with LULUCF. Totals may not sum due to independent rounding. Parentheses indicate negative values or sequestration.

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Emissions from Wood Biomass, Ethanol, and Biodiesel Consumption are not included specifically in summing Energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for LULUCF.

^b Emissions from International Bunker Fuels are not included in totals.

^c LULUCF emissions of CH₄ and N₂O are reported separately from gross emissions totals. LULUCF emissions include the CH₄ and N₂O emissions reported for *Peatlands Remaining Peatlands*, *Forest Fires*, *Drained Organic Soils*, *Grassland Fires*, and *Coastal Wetlands Remaining Coastal Wetlands*; CH₄ emissions from *Land Converted to Coastal Wetlands*; and N₂O emissions from *Forest Soils* and *Settlement Soils*. Refer to Table 2-8 for a breakout of emissions and removals for LULUCF by gas and source category.

^d Small amounts of PFC emissions also result from this source.

^e LULUCF Carbon Stock Change is the net C stock change from the following categories: *Forest Land Remaining Forest Land*, *Land Converted to Forest Land*, *Cropland Remaining Cropland*, *Land Converted to Cropland*, *Grassland Remaining Grassland*, *Land Converted to Grassland*, *Wetlands Remaining Wetlands*, *Land Converted to Wetlands*, *Settlements*

Remaining Settlements, and Land Converted to Settlements. Refer to Table 2-8 for a breakout of emissions and removals for LULUCF by gas and source category.

^f The LULUCF Sector Net Total is the net sum of all LULUCF CH₄ and N₂O emissions to the atmosphere plus net carbon stock changes.

1 **Table 2-2: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (kt)**

Gas/Source	1990	2005	2015	2016	2017	2018	2019
CO₂	5,116,575	6,141,018	5,384,235	5,262,571	5,219,364	5,397,305	5,274,426
Fossil Fuel Combustion	4,731,460	5,753,835	5,013,834	4,922,044	4,870,374	5,023,102	4,888,463
<i>Transportation</i>	<i>1,469,108</i>	<i>1,868,757</i>	<i>1,750,510</i>	<i>1,797,265</i>	<i>1,812,632</i>	<i>1,821,191</i>	<i>1,843,175</i>
<i>Electric Power</i>	<i>1,819,951</i>	<i>2,400,057</i>	<i>1,900,637</i>	<i>1,808,871</i>	<i>1,732,031</i>	<i>1,752,936</i>	<i>1,606,024</i>
<i>Industrial</i>	<i>853,807</i>	<i>843,952</i>	<i>790,319</i>	<i>787,026</i>	<i>794,051</i>	<i>838,004</i>	<i>837,591</i>
<i>Residential</i>	<i>338,578</i>	<i>359,086</i>	<i>318,085</i>	<i>292,338</i>	<i>294,649</i>	<i>339,538</i>	<i>338,826</i>
<i>Commercial</i>	<i>228,298</i>	<i>226,019</i>	<i>224,866</i>	<i>210,394</i>	<i>212,288</i>	<i>246,870</i>	<i>238,280</i>
<i>U.S. Territories</i>	<i>21,718</i>	<i>55,963</i>	<i>29,417</i>	<i>26,150</i>	<i>24,723</i>	<i>24,563</i>	<i>24,567</i>
Non-Energy Use of Fuels	112,766	130,567	105,340	93,590	99,686	110,880	110,108
Petroleum Systems	9,732	12,071	32,434	21,855	24,813	36,588	43,308
Iron and Steel Production & Metallurgical Coke Production	104,730	70,076	47,941	43,621	40,566	42,627	41,290
Cement Production	33,484	46,194	39,907	39,439	40,324	38,971	40,896
Natural Gas Systems	32,026	25,182	29,159	30,067	31,207	33,864	36,854
Petrochemical Production	21,611	27,383	28,062	28,310	28,910	29,314	30,792
Incineration of Waste	11,183	17,434	21,478	21,830	21,281	20,886	20,199
Lime Production	11,700	14,552	13,333	12,545	12,875	13,112	12,963
Ammonia Production	13,047	9,177	10,616	10,245	11,112	12,163	12,272
Other Process Uses of Carbonates	6,297	7,644	12,182	10,972	9,933	7,469	7,457
Urea Consumption for Non-Agricultural Purposes	3,784	3,653	4,578	5,132	4,972	6,056	6,569
Urea Fertilization	2,417	3,504	4,728	4,877	5,051	5,192	5,341
Carbon Dioxide Consumption	1,472	1,375	4,940	4,640	4,580	4,130	4,870
Liming	4,667	4,349	3,737	3,081	3,080	2,248	2,442
Aluminum Production	6,831	4,142	2,767	1,334	1,205	1,451	1,880
Soda Ash Production	1,431	1,655	1,714	1,723	1,753	1,714	1,792
Ferroalloy Production	2,152	1,392	1,960	1,796	1,975	2,063	1,598
Titanium Dioxide Production	1,195	1,755	1,635	1,662	1,688	1,541	1,474
Glass Production	1,535	1,928	1,299	1,249	1,296	1,305	1,280
Zinc Production	632	1,030	933	882	951	982	964
Phosphoric Acid Production	1,529	1,342	999	998	1,028	940	891
Lead Production	516	553	473	500	513	513	540
Carbide Production and Consumption	370	213	176	170	181	184	175
Abandoned Oil and Gas Wells	6	7	7	7	7	7	7
Magnesium Production and Processing	1	3	3	3	3	1	1
<i>Wood Biomass, Ethanol, and Biodiesel Consumption^a</i>	<i>219,413</i>	<i>230,700</i>	<i>317,742</i>	<i>316,610</i>	<i>312,304</i>	<i>319,647</i>	<i>316,191</i>
<i>International Bunker Fuels^b</i>	<i>103,471</i>	<i>113,188</i>	<i>110,803</i>	<i>116,501</i>	<i>120,009</i>	<i>122,005</i>	<i>116,056</i>
CH₄^c	31,201	27,523	26,084	25,741	25,968	26,235	26,404
Enteric Fermentation	6,588	6,772	6,675	6,890	7,032	7,119	7,142
Natural Gas Systems	7,508	6,547	5,944	5,868	5,905	6,034	6,258
Landfills	7,182	5,255	4,456	4,321	4,375	4,482	4,580

Manure Management	1,485	2,062	2,316	2,385	2,395	2,467	2,495
Coal Mining	3,860	2,629	2,471	2,172	2,216	2,136	1,895
Petroleum Systems	1,941	1,624	1,720	1,631	1,635	1,540	1,607
Wastewater Treatment	806	803	753	747	739	737	736
Rice Cultivation	640	720	648	631	596	623	602
Stationary Combustion	344	313	336	311	303	343	345
Abandoned Oil and Gas Wells	263	278	286	289	282	281	281
Abandoned Underground							
Coal Mines	288	264	256	268	257	247	237
Mobile Combustion	254	157	105	101	99	97	95
Composting	15	75	85	91	98	90	91
Field Burning of Agricultural							
Residues	15	17	18	17	17	17	17
Petrochemical Production	9	3	7	10	10	12	13
Anaerobic Digestion at Biogas							
Facilities	1	3	8	8	8	8	8
Ferroalloy Production	1	+	1	1	1	1	+
Carbide Production and							
Consumption	1	+	+	+	+	+	+
Iron and Steel Production &							
Metallurgical Coke							
Production	1	1	+	+	+	+	+
Incineration of Waste	+	+	+	+	+	+	+
<i>International Bunker Fuels^b</i>	7	5	4	4	4	4	4
N₂O^c	1,521	1,531	1,572	1,514	1,499	1,542	1,536
Agricultural Soil Management	1,060	1,052	1,169	1,108	1,099	1,135	1,156
Wastewater Treatment	63	77	85	87	89	88	88
Stationary Combustion	84	115	102	100	95	95	83
Manure Management	47	55	59	61	63	65	66
Mobile Combustion	152	141	73	71	68	64	62
Nitric Acid Production	41	38	39	34	31	32	34
Adipic Acid Production	51	24	14	23	25	35	18
N ₂ O from Product Uses	14	14	14	14	14	14	14
Composting	1	6	6	7	7	7	7
Caprolactam, Glyoxal, and							
Glyoxylic Acid Production	6	7	6	6	5	5	5
Incineration of Waste	2	1	1	1	1	1	1
Electronics Industry	+	+	1	1	1	1	1
Field Burning of Agricultural							
Residues	1	1	1	1	1	1	1
Petroleum Systems	+	+	+	+	+	+	+
Natural Gas Systems	+	+	+	+	+	+	+
<i>International Bunker Fuels^b</i>	3	3	3	3	4	4	3
HFCs	M						
Substitution of Ozone							
Depleting Substances ^d	M	M	M	M	M	M	M
HCFC-22 Production	3	1	+	+	+	+	+
Electronics Industry	M	M	M	M	M	M	M
Magnesium Production and							
Processing	0	0	+	+	+	+	+
PFCs	M						
Electronics Industry	M	M	M	M	M	M	M
Aluminum Production	M	M	M	M	M	M	M
Substitution of Ozone							
Depleting Substances ^d	0	+	+	+	+	+	+

SF₆	1	1	+	+	+	+	+
Electrical Transmission and Distribution	1	+	+	+	+	+	+
Magnesium Production and Processing	+	+	+	+	+	+	+
Electronics Industry	+	+	+	+	+	+	+
NF₃	+	+	+	+	+	+	+
Electronics Industry	+	+	+	+	+	+	+
Unspecified Mix of HFCs, PFCs, SF₆, and NF₃	M						
Electronics Industry	M	M	M	M	M	M	M

Notes: Totals may not sum due to independent rounding. Parentheses indicate negative values or sequestration.

+ Does not exceed 0.5 kt.

M - Mixture of multiple gases

^a Emissions from Wood Biomass, Ethanol, and Biodiesel Consumption are not included specifically in summing Energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for LULUCF.


^b Emissions from International Bunker Fuels are not included in totals.

^c LULUCF emissions of LULUCF CH₄ and N₂O are reported separately from gross emissions totals. Refer to Table 2-8 for a breakout of emissions and removals for LULUCF by gas and source category.

^d Small amounts of PFC emissions also result from this source.

- 1 Emissions of all gases can be summed from each source category into a set of five sectors defined by the
- 2 Intergovernmental Panel on Climate Change (IPCC). Figure 2-4 and Table 2-3 illustrate that over the thirty-year
- 3 period of 1990 to 2019, total emissions from the Energy, Industrial Processes and Product Use, and Agriculture
- 4 sectors grew by 81.4 MMT CO₂ Eq. (1.5 percent), 28.5 MMT CO₂ Eq. (8.2 percent), and 73.3 MMT CO₂ Eq. (13.2
- 5 percent), respectively. Emissions from the Waste sector decreased by 55.4 MMT CO₂ Eq. (25.3 percent). Over the
- 6 same period, total C sequestration in the Land Use, Land-Use Change, and Forestry (LULUCF) sector decreased by
- 7 96.3 MMT CO₂ (10.6 percent decrease in total C sequestration), and emissions from the LULUCF sector increased
- 8 by 15.5 MMT CO₂ Eq. (196.1 percent).

1 **Figure 2-4: U.S. Greenhouse Gas Emissions and Sinks by Chapter/IPCC Sector**



2
3 **Table 2-3: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks by Chapter/IPCC**
4 **Sector (MMT CO₂ Eq.)**

Chapter/IPCC Sector	1990	2005	2015	2016	2017	2018	2019
Energy	5,329.4	6,311.1	5,532.9	5,406.9	5,363.8	5,540.1	5,410.8
Fossil Fuel Combustion	4,731.5	5,753.8	5,013.8	4,922.0	4,870.4	5,023.1	4,888.5
Natural Gas Systems	219.7	188.9	177.8	176.8	178.8	184.7	193.3
Non-Energy Use of Fuels	112.8	130.6	105.3	93.6	99.7	110.9	110.1
Petroleum Systems	58.3	52.7	75.5	62.7	65.7	75.1	83.5
Coal Mining	96.5	65.7	61.8	54.3	55.4	53.4	47.4
Stationary Combustion	33.7	42.2	38.8	37.7	35.9	36.8	33.5
Mobile Combustion	51.6	45.9	24.5	23.6	22.7	21.6	20.9
Incineration of Waste	11.6	17.8	21.9	22.3	21.7	21.3	20.6
Abandoned Oil and Gas Wells	6.6	7.0	7.2	7.2	7.1	7.0	7.0
Abandoned Underground Coal Mines	7.2	6.6	6.4	6.7	6.4	6.2	5.9
Industrial Processes and Product Use	345.5	365.7	375.4	367.9	367.6	371.0	374.0
Substitution of Ozone Depleting Substances	0.2	107.3	163.6	164.9	164.6	165.6	169.7
Iron and Steel Production & Metallurgical Coke Production	104.8	70.1	47.9	43.6	40.6	42.6	41.3
Cement Production	33.5	46.2	39.9	39.4	40.3	39.0	40.9
Petrochemical Production	21.8	27.5	28.2	28.6	29.2	29.6	31.1
Lime Production	11.7	14.6	13.3	12.5	12.9	13.1	13.0
Ammonia Production	13.0	9.2	10.6	10.2	11.1	12.2	12.3
Nitric Acid Production	12.1	11.3	11.6	10.1	9.3	9.6	10.0
Other Process Uses of Carbonates	6.3	7.6	12.2	11.0	9.9	7.5	7.5
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	4.6	5.1	5.0	6.1	6.6
Adipic Acid Production	15.2	7.1	4.3	7.0	7.4	10.3	5.3
Carbon Dioxide Consumption	1.5	1.4	4.9	4.6	4.6	4.1	4.9
Electronics Industry	3.6	4.8	5.0	5.0	4.9	5.1	4.6

Electrical Transmission and Distribution	23.2	8.4	3.8	4.1	4.2	3.9	4.2
N ₂ O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
HCFC-22 Production	46.1	20.0	4.3	2.8	5.2	3.3	3.7
Aluminum Production	28.3	7.6	4.9	2.7	2.3	3.1	3.6
Soda Ash Production	1.4	1.7	1.7	1.7	1.8	1.7	1.8
Ferroalloy Production	2.2	1.4	2.0	1.8	2.0	2.1	1.6
Titanium Dioxide Production	1.2	1.8	1.6	1.7	1.7	1.5	1.5
Caprolactam, Glyoxal, and Glyoxylic Acid Production	1.7	2.1	1.9	1.7	1.5	1.4	1.4
Glass Production	1.5	1.9	1.3	1.2	1.3	1.3	1.3
Magnesium Production and Processing	5.2	2.7	1.1	1.2	1.2	1.2	1.0
Zinc Production	0.6	1.0	0.9	0.9	1.0	1.0	1.0
Phosphoric Acid Production	1.5	1.3	1.0	1.0	1.0	0.9	0.9
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.2
Agriculture	555.3	577.1	616.1	604.4	605.5	621.0	628.6
Agricultural Soil Management	315.9	313.4	348.5	330.1	327.6	338.2	344.6
Enteric Fermentation	164.7	169.3	166.9	172.2	175.8	178.0	178.6
Manure Management	51.1	67.9	75.4	77.7	78.5	81.1	82.0
Rice Cultivation	16.0	18.0	16.2	15.8	14.9	15.6	15.1
Urea Fertilization	2.4	3.5	4.7	4.9	5.1	5.2	5.3
Liming	4.7	4.3	3.7	3.1	3.1	2.2	2.4
Field Burning of Agricultural Residues	0.5	0.6	0.6	0.6	0.6	0.6	0.6
Waste	219.2	178.0	159.8	157.1	159.1	161.1	163.8
Landfills	179.6	131.4	111.4	108.0	109.4	112.1	114.5
Wastewater Treatment	38.9	43.0	44.2	44.6	44.9	44.6	44.8
Composting	0.7	3.5	4.0	4.3	4.6	4.3	4.3
Anaerobic Digestion at Biogas Facilities	+	0.1	0.2	0.2	0.2	0.2	0.2
Total Emissions (Sources)^a	6,449.4	7,431.9	6,684.3	6,536.3	6,496.0	6,693.2	6,577.2
LULUCF Sector Net Total^b	(900.8)	(788.0)	(763.6)	(842.6)	(765.8)	(801.2)	(788.9)
Forest land	(884.1)	(751.4)	(749.5)	(814.7)	(740.0)	(781.4)	(774.6)
Cropland	28.6	23.2	43.2	31.7	32.3	37.7	39.7
Grassland	2.2	(29.4)	(10.1)	(13.7)	(12.5)	(11.9)	(8.0)
Wetlands	(2.8)	(1.9)	(3.7)	(3.7)	(3.6)	(3.6)	(3.6)
Settlements	(44.7)	(28.5)	(43.5)	(42.2)	(42.1)	(42.0)	(42.4)
Net Emission (Sources and Sinks)^c	5,548.6	6,643.8	5,920.7	5,693.8	5,730.2	5,892.0	5,788.3

Notes: Total emissions presented without LULUCF. Net emissions presented with LULUCF. Totals may not sum due to independent rounding. Parentheses indicate negative values or sequestration.

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Total emissions without LULUCF.

^b LULUCF emissions of CH₄ and N₂O are reported separately from gross emissions totals. LULUCF emissions include the CH₄ and N₂O emissions reported for *Peatlands Remaining Peatlands*, *Forest Fires*, *Drained Organic Soils*, *Grassland Fires*, and *Coastal Wetlands Remaining Coastal Wetlands*; CH₄ emissions from *Land Converted to Coastal Wetlands*; and N₂O emissions from *Forest Soils* and *Settlement Soils*. Refer to Table 2-8 for a breakout of emissions and removals for LULUCF by gas and source category.

^c Net emissions with LULUCF.

1 Energy


- 2 Energy-related activities, primarily fossil fuel combustion, accounted for the vast majority of U.S. CO₂ emissions for
3 the period of 1990 through 2019. Fossil fuel combustion is the largest source of energy-related emissions, with CO₂

1 being the primary gas emitted (see Figure 2-5). Due to their relative importance, fossil fuel combustion-related CO₂
2 emissions are considered in detail in the Energy chapter (see Energy chapter).

3 In 2019, approximately 80 percent of the energy used in the United States (on a Btu basis) was produced through
4 the combustion of fossil fuels. The remaining 20 percent came from other energy sources such as hydropower,
5 biomass, nuclear, wind, and solar energy. A discussion of specific trends related to CO₂ as well as other greenhouse
6 gas emissions from energy use is presented here with more detail in the Energy chapter. Energy-related activities
7 are also responsible for CH₄ and N₂O emissions (41 percent and 10 percent of total U.S. emissions of each gas,
8 respectively). Table 2-4 presents greenhouse gas emissions from the Energy chapter, by source and gas.

9


10 **Figure 2-5: 2019 Energy Chapter Greenhouse Gas Sources**



11

12

1 **Figure 2-6: Trends in Energy Chapter Greenhouse Gas Sources**



2

3 **Table 2-4: Emissions from Energy (MMT CO₂ Eq.)²**

Gas/Source	1990	2005	2015	2016	2017	2018	2019
CO₂	4,897.2	5,939.1	5,202.3	5,089.4	5,047.4	5,225.3	5,098.9
Fossil Fuel Combustion	4,731.5	5,753.8	5,013.8	4,922.0	4,870.4	5,023.1	4,888.5
<i>Transportation</i>	1,469.1	1,868.8	1,750.5	1,797.3	1,812.6	1,821.2	1,843.2
<i>Electric Power</i>	1,820.0	2,400.1	1,900.6	1,808.9	1,732.0	1,752.9	1,606.0
<i>Industrial</i>	853.8	844.0	790.3	787.0	794.1	838.0	837.6
<i>Residential</i>	338.6	359.1	318.1	292.3	294.6	339.5	338.8
<i>Commercial</i>	228.3	226.0	224.9	210.4	212.3	246.9	238.3
<i>U.S. Territories</i>	21.7	56.0	29.4	26.2	24.7	24.6	24.6
Non-Energy Use of Fuels	112.8	130.6	105.3	93.6	99.7	110.9	110.1
Petroleum Systems	9.7	12.1	32.4	21.9	24.8	36.6	43.3
Natural Gas Systems	32.0	25.2	29.2	30.1	31.2	33.9	36.9
Incineration of Waste	11.2	17.4	21.5	21.8	21.3	20.9	20.2
Abandoned Oil and Gas Wells	+	+	+	+	+	+	+
<i>Biomass-Wood^a</i>	215.2	206.9	224.7	215.7	211.5	219.8	216.5
<i>International Bunker Fuels^b</i>	103.5	113.2	110.8	116.5	120.0	122.0	116.1
<i>Biofuels-Ethanol^a</i>	4.2	22.9	78.9	81.2	82.1	81.9	82.6
<i>Biofuels-Biodiesel^a</i>	+	0.9	14.1	19.6	18.7	17.9	17.1
CH₄	361.4	295.3	277.9	266.0	267.4	266.9	268.0
Natural Gas Systems	187.7	163.7	148.6	146.7	147.6	150.8	156.4
Coal Mining	96.5	65.7	61.8	54.3	55.4	53.4	47.4

² The full time-series data is available in Common Reporting Format (CRF) Tables included in the U.S. UNFCCC submission and in CSV format available at <<https://www.epa.gov/ghgemissions/inventory-us-greenhouse-gas-emissions-and-sinks>>.

Petroleum Systems	48.5	40.6	43.0	40.8	40.9	38.5	40.2
Stationary Combustion	8.6	7.8	8.4	7.8	7.6	8.6	8.6
Abandoned Oil and Gas Wells	6.6	7.0	7.1	7.2	7.1	7.0	7.0
Abandoned Underground Coal Mines	7.2	6.6	6.4	6.7	6.4	6.2	5.9
Mobile Combustion	6.4	3.9	2.6	2.5	2.5	2.4	2.4
Incineration of Waste	+	+	+	+	+	+	+
<i>International Bunker Fuels^b</i>	0.2	0.1	0.1	0.1	0.1	0.1	0.1
N₂O	70.8	76.7	52.8	51.5	49.0	47.9	43.9
Stationary Combustion	25.1	34.3	30.4	29.9	28.4	28.2	24.9
Mobile Combustion	45.2	42.0	21.9	21.1	20.2	19.1	18.6
Incineration of Waste	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Petroleum Systems	+	+	+	+	+	+	+
Natural Gas Systems	+	+	+	+	+	+	+
<i>International Bunker Fuels^b</i>	0.9	1.0	1.0	1.0	1.1	1.1	1.0
Total	5,329.4	6,311.1	5,532.9	5,406.9	5,363.8	5,540.1	5,410.8

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Emissions from Wood Biomass and Biofuel Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for LULUCF.

^b Emissions from International Bunker Fuels are not included in totals.

1 CO₂ Emissions from Fossil Fuel Combustion

2 As the largest contributor to U.S. greenhouse gas emissions, CO₂ from fossil fuel combustion has accounted for
3 approximately 76 percent of gross emissions across the time series. Within the United States, fossil fuel
4 combustion accounted for 92.7 percent of CO₂ emissions in 2019. Emissions from this source category grew by 3.3
5 percent (157.0 MMT CO₂ Eq.) from 1990 to 2019 and were responsible for most of the increase in national
6 emissions during this period. Conversely, CO₂ emissions from fossil fuel combustion decreased by 865.4 MMT CO₂
7 Eq. from 2005 and by 458.2 MMT CO₂ Eq. from 2010, representing decreases of approximately 15.0 percent
8 between 2005 and 2019 and 8.6 percent between 2010 and 2019. From 2018 to 2019, these emissions decreased
9 by 2.7 percent (134.6 MMT CO₂ Eq.). Historically, changes in emissions from fossil fuel combustion have been the
10 main factor influencing U.S. emission trends.

11 Changes in CO₂ emissions from fossil fuel combustion since 1990 are affected by many long-term and short-term
12 factors, including population and economic growth, energy price fluctuations and market trends, technological
13 changes, carbon intensity of energy fuel choices, and seasonal temperatures. CO₂ emissions from coal combustion
14 gradually increased between 1990 and 2007, then began to decrease at a faster rate from 2008 to 2019. CO₂
15 emissions from natural gas combustion remained relatively constant, with a slight increase between 1990 and
16 2009, then began to consistently increase between 2010 and 2019. The replacement of coal combustion with
17 natural gas combustion was largely driven by new discoveries of natural gas fields and advancements in drilling
18 technologies, which led to lower natural gas prices. On an annual basis, the overall consumption and mix of fossil
19 fuels in the United States fluctuates primarily in response to changes in general economic conditions, overall
20 energy prices, the relative price of different fuels, weather, and the availability of non-fossil alternatives. For
21 example, coal consumption for electric power is influenced by a number of factors, including the relative price of
22 coal and alternative sources, the ability to switch fuels, and longer-term trends in coal markets. Likewise, warmer
23 winters lead to a decrease in heating degree days and result in a decreased demand for heating fuel and electricity
24 for heat in the residential and commercial sectors, which leads to a decrease in emissions from reduced fuel
25 consumption.

26 Fossil fuel combustion CO₂ emissions also depend on the type of fuel consumed or energy used and its carbon
27 intensity. Producing a unit of heat or electricity using natural gas instead of coal, for example, reduces CO₂
28 emissions because of the lower C content of natural gas (see Table A-29 in Annex 2.1 for more detail on the C
29 Content Coefficient of different fossil fuels).

1 Recent trends in CO₂ emissions from fossil fuel combustion have been strongly influenced by trends in the electric
2 power sector, which historically has accounted for the largest share of emissions from this source (see Figure 2-7).
3 Electric power sector emissions are driven by the total amount of electricity produced and the carbon intensity of
4 the fuels used to produce the electricity. The types of fuel consumed to produce electricity have shifted over time,
5 impacting emission trends. Electricity generation from lower carbon intensity renewable energy sources increased
6 by 115 percent from 2005 to 2019 and natural gas generation increased by 116 percent while coal generation
7 decreased by 52 percent over the same time period (see Table 3-12 for more detail on electricity generation by
8 source). The decrease in coal-powered electricity generation and increase in natural gas and renewable energy
9 electricity generation have contributed to a 33 percent decrease in overall CO₂ emissions from electric power
10 generation from 2005 to 2019 (see Figure 2-9). Between 2018 and 2019, emissions from the electric power sector
11 decreased 8.4 percent due to a decrease in electric power generation of 1.4 percent and the continued shift in the
12 share of electric power generation from coal to natural gas and renewable energy.

13 Petroleum use is another major driver of CO₂ emissions from fossil fuel combustion, particularly in the
14 transportation sector, which represents the largest source of CO₂ emissions from fossil fuel combustion in 2019.
15 Emissions from petroleum consumption for transportation (including bunker fuels) have increased by 4.5 percent
16 since 2015; this trend can be primarily attributed to a 5.6 percent increase in vehicle miles traveled (VMT) over the
17 same time period. Fuel economy of light-duty vehicles is another important factor. The decline in new light-duty
18 vehicle fuel economy between 1990 and 2004 reflected the increasing market share of light-duty trucks, which
19 grew from about 30 percent of new vehicle sales in 1990 to 48 percent in 2004. Starting in 2005, average new
20 vehicle fuel economy began to increase while light-duty VMT grew only modestly for much of the period and has
21 slowed the rate of increase of CO₂ emissions.

22 Overall, across all sectors, there was a 2.7 percent decrease in total CO₂ emissions from fossil fuel combustion from
23 2018 to 2019 and a 2.5 percent reduction since 2015. Carbon dioxide emissions from fossil fuel combustion,
24 separated by end-use sector, are presented in Table 2-5 and Figure 2-7 based on the underlying U.S. energy
25 consumer data collected by the U.S. Energy Information Administration (EIA). Figure 2-8 further describes direct
26 and indirect CO₂ emissions from fossil fuel combustion, separated by end-use sector. Estimates of CO₂ emissions
27 from fossil fuel combustion are calculated from these EIA “end-use sectors” based on total fuel consumption and
28 appropriate fuel properties described below. (Any additional analysis and refinement of the EIA data is further
29 explained in the Energy chapter of this report.)

- 30 • *Transportation.* EIA’s fuel consumption data for the transportation sector consists of all vehicles whose
31 primary purpose is transporting people and/or goods from one physical location to another.
- 32 • *Industry.* EIA statistics for the industrial sector include fossil fuel consumption that occurs in the fields of
33 manufacturing, agriculture, mining, and construction. EIA’s fuel consumption data for the industrial sector
34 consist of all facilities and equipment used for producing, processing, or assembling goods. (EIA includes
35 generators that produce electricity and/or useful thermal output primarily to support on-site industrial
36 activities in this sector.)
- 37 • *Electric Power.* EIA’s fuel consumption data for the electric power sector are comprised of electricity-only
38 and combined-heat-and-power (CHP) plants within the North American Industry Classification System
39 (NAICS) 22 category whose primary business is to sell electricity, or electricity and heat, to the public.
40 (Non-utility power producers are included in this sector as long as they meet the electric power sector
41 definition.)
- 42 • *Residential.* EIA’s fuel consumption data for the residential sector consist of living quarters for private
43 households.
- 44 • *Commercial.* EIA’s fuel consumption data for the commercial sector consist of service-providing facilities
45 and equipment from private and public organizations and businesses. (EIA includes generators that
46 produce electricity and/or useful thermal output primarily to support the activities at commercial
47 establishments in this sector.)


48 **Table 2-5: CO₂ Emissions from Fossil Fuel Combustion by End-Use Sector (MMT CO₂ Eq.)**

End-Use Sector	1990	2005	2015	2016	2017	2018	2019
Transportation	1,472.2	1,873.5	1,754.8	1,801.5	1,816.9	1,825.9	1,847.9
Combustion	1,469.1	1,868.8	1,750.5	1,797.3	1,812.6	1,821.2	1,843.2
Electricity	3.0	4.7	4.3	4.2	4.3	4.7	4.7
Industrial	1,540.2	1,580.3	1,339.9	1,304.6	1,298.5	1,339.3	1,302.9
Combustion	853.8	844.0	790.3	787.0	794.1	838.0	837.6
Electricity	686.4	736.3	549.5	517.6	504.4	501.3	465.3
Residential	931.3	1,215.1	1,001.9	945.8	911.7	981.7	922.3
Combustion	338.6	359.1	318.1	292.3	294.6	339.5	338.8
Electricity	592.7	856.0	683.8	653.5	617.1	642.1	583.5
Commercial	766.0	1,029.0	887.9	844.0	818.5	851.7	790.7
Combustion	228.3	226.0	224.9	210.4	212.3	246.9	238.3
Electricity	537.7	803.0	663.0	633.6	606.2	604.8	552.4
U.S. Territories^a	21.7	56.0	29.4	26.2	24.7	24.6	24.6
Total	4,731.5	5,753.8	5,013.8	4,922.0	4,870.4	5,023.1	4,888.5
Electric Power	1,820.0	2,400.1	1,900.6	1,808.9	1,732.0	1,752.9	1,606.0

Notes: Combustion-related emissions from electric power are allocated based on aggregate national electricity use by each end-use sector. Totals may not sum due to independent rounding.

^a Fuel consumption by U.S. Territories (i.e., American Samoa, Guam, Puerto Rico, U.S. Virgin Islands, Wake Island, and other U.S. Pacific Islands) is included in this report.


1 **Figure 2-7: 2019 CO₂ Emissions from Fossil Fuel Combustion by Sector and Fuel Type**



2


3 Note on Figure 2-7: Fossil Fuel Combustion for electric power also includes emissions of less than 0.5 MMT CO₂ Eq. from
 4 geothermal-based generation.

1 **Figure 2-8: 2019 End-Use Sector Emissions of CO₂ from Fossil Fuel Combustion**



2
 3 Electric power was the second largest emitter of CO₂ in 2019 (surpassed by transportation); electric power
 4 generators used 30 percent of U.S. energy from fossil fuels and emitted 33 percent of the CO₂ from fossil fuel
 5 combustion in 2019. Changes in electricity demand and the carbon intensity of fuels used for electric power
 6 generation have a significant impact on CO₂ emissions. Carbon dioxide emissions from the electric power sector
 7 have decreased by approximately 11.8 percent since 1990, and the carbon intensity of the electric power sector, in
 8 terms of CO₂ Eq. per QBtu input, has significantly decreased by 16 percent during that same timeframe. This
 9 decoupling of electric power generation and the resulting CO₂ emissions is shown below in Figure 2-9.

10 **Figure 2-9: Electric Power Generation (Billion kWh) and Emissions**



11
 12 Electric power CO₂ emissions can also be allocated to the end-use sectors that use electricity, as presented in Table
 13 2-5. With electricity CO₂ emissions allocated to end-use sectors, the transportation end-use sector represents the
 14 largest source of fossil fuel combustion emissions accounting for 1,847.9 MMT CO₂ Eq. in 2019 or approximately 38
 15 percent of total CO₂ emissions from fossil fuel combustion. The industrial end-use sector accounted for 27 percent
 16 of CO₂ emissions from fossil fuel combustion when including allocated electricity emissions. The residential and
 17 commercial end-use sectors accounted for 19 and 16 percent, respectively, of CO₂ emissions from fossil fuel

1 combustion when including allocated electricity emissions. Both of these end-use sectors were heavily reliant on
2 electricity for meeting energy needs, with electricity use for lighting, heating, air conditioning, and operating
3 appliances contributing 63 and 70 percent of emissions from the residential and commercial end-use sectors,
4 respectively.

5 **Other Significant Trends in Energy**

6 Other significant trends in emissions from energy source categories (Figure 2-6) over the thirty-year period from
7 1990 through 2019 included the following:

- 8 • Methane emissions from natural gas systems and petroleum systems (combined here) decreased 39.6
9 MMT CO₂ Eq. (16.8 percent decrease from 1990 to 2019) or from 236.2 MMT CO₂ Eq. in 1990 to 196.6
10 MMT CO₂ Eq. in 2019. Natural gas systems CH₄ emissions decreased by 31.3 MMT CO₂ Eq. (16.7 percent)
11 since 1990, largely due to a decrease in emissions from distribution, transmission and storage, processing,
12 and exploration. The decrease in distribution is largely due to decreased emissions from pipelines and
13 distribution station leaks, and the decrease in transmission and storage emissions is largely due to
14 reduced compressor station emissions (including emissions from compressors and leaks). Petroleum
15 systems CH₄ emissions decreased by 8.3 MMT CO₂ Eq. (or 17.2 percent) since 1990. This decrease is due
16 primarily to decreases in emissions from offshore platforms, tanks, and pneumatic controllers. Carbon
17 dioxide emissions from natural gas and petroleum systems increased by 38.4 MMT CO₂ Eq. (92.0 percent)
18 from 1990 to 2019. This increase is due primarily to increases in the production segment, where flaring
19 emissions from associated gas flaring, tanks, and miscellaneous production flaring have increased over
20 time.
- 21 • Methane emissions from coal mining decreased by 49.1 MMT CO₂ Eq. (50.9 percent) from 1990 through
22 2019, primarily due to a decrease in the number of active mines and annual coal production over the time
23 period.
- 24 • Nitrous oxide emissions from mobile combustion decreased by 26.6 MMT CO₂ Eq. (58.9 percent) from
25 1990 through 2019, primarily as a result of national vehicle emissions standards and emission control
26 technologies for on-road vehicles.
- 27 • Carbon dioxide emissions from non-energy uses of fossil fuels decreased by 2.7 MMT CO₂ Eq. (2.4
28 percent) from 1990 through 2019. Emissions from non-energy uses of fossil fuels were 110.1 MMT CO₂
29 Eq. in 2019, which constituted 2.1 percent of total national CO₂ emissions, approximately the same
30 proportion as in 1990.
- 31 • Nitrous oxide emissions from stationary combustion decreased by 0.2 MMT CO₂ Eq. (0.8 percent) from
32 1990 through 2019. Nitrous oxide emissions from this source decreased primarily as a result of a decrease
33 in overall coal combustion in the electric power sector, particularly from tangentially-fired coal and dry
34 bottom wall-fired boilers.
- 35 • Carbon dioxide emissions from incineration of waste (20.2 MMT CO₂ Eq. in 2019) increased by 9.0 MMT
36 CO₂ Eq. (80.6 percent) from 1990 through 2019, as the volume of scrap tires and other fossil C-containing
37 materials in waste increased.


38 **Industrial Processes and Product Use**

39 In many cases, greenhouse gas emissions are generated and emitted in two different ways. First, they are
40 generated and emitted as the byproducts of many non-energy-related industrial activities. For example, industrial
41 processes can chemically or physically transform raw materials, which often release waste gases such as CO₂, CH₄,
42 N₂O, and fluorinated gases (e.g., HFC-23). In the case of byproduct emissions, the emissions are generated by an
43 industrial process itself, and are not directly a result of energy consumed during the process.

44 Second, industrial manufacturing processes and use by end-consumers also release HFCs, PFCs, SF₆, and NF₃ and
45 other fluorinated compounds. In addition to the use of HFCs and some PFCs as substitutes for ozone depleting


1 substances (ODS), fluorinated compounds such as HFCs, PFCs, SF₆, NF₃, and others are also emitted through use by
 2 a number of other industrial sources in the United States. These industries include the electronics industry, electric
 3 power transmission and distribution, and magnesium metal production and processing. In addition, N₂O is used in
 4 and emitted by the electronics industry and anesthetic and aerosol applications. Figure 2-10 and Table 2-6
 5 presents greenhouse gas emissions from industrial processes and product use by source category. Overall,
 6 emission sources in the Industrial Processes and Product Use (IPPU) chapter account for 5.7 percent of U.S.
 7 greenhouse gas emissions in 2019.

8 **Figure 2-10: 2019 Industrial Processes and Product Use Chapter Greenhouse Gas Source**



9

1 **Figure 2-11: Trends in Industrial Processes and Product Use Chapter Greenhouse Gas**
 2 **Sources**



3
 4 **Table 2-6: Emissions from Industrial Processes and Product Use (MMT CO₂ Eq.)**

Gas/Source	1990	2005	2015	2016	2017	2018	2019
CO₂	212.3	194.1	173.5	165.2	163.9	164.5	167.7
Iron and Steel Production & Metallurgical Coke Production	104.7	70.1	47.9	43.6	40.6	42.6	41.3
<i>Iron and Steel Production</i>	99.1	66.2	43.5	41.0	38.6	41.3	39.9
<i>Metallurgical Coke Production</i>	5.6	3.9	4.4	2.6	2.0	1.3	1.4
Cement Production	33.5	46.2	39.9	39.4	40.3	39.0	40.9
Petrochemical Production	21.6	27.4	28.1	28.3	28.9	29.3	30.8
Lime Production	11.7	14.6	13.3	12.5	12.9	13.1	13.0
Ammonia Production	13.0	9.2	10.6	10.2	11.1	12.2	12.3
Other Process Uses of Carbonates	6.3	7.6	12.2	11.0	9.9	7.5	7.5
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	4.6	5.1	5.0	6.1	6.6
Carbon Dioxide Consumption	1.5	1.4	4.9	4.6	4.6	4.1	4.9
Ferroalloy Production	2.2	1.4	2.0	1.8	2.0	2.1	1.6
Soda Ash Production	1.4	1.7	1.7	1.7	1.8	1.7	1.8
Titanium Dioxide Production	1.2	1.8	1.6	1.7	1.7	1.5	1.5
Aluminum Production	6.8	4.1	2.8	1.3	1.2	1.5	1.9
Glass Production	1.5	1.9	1.3	1.2	1.3	1.3	1.3
Zinc Production	0.6	1.0	0.9	0.9	1.0	1.0	1.0
Phosphoric Acid Production	1.5	1.3	1.0	1.0	1.0	0.9	0.9
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.2

Magnesium Production and Processing	+	+	+	+	+	+	+
CH₄	0.3	0.1	0.2	0.3	0.3	0.3	0.4
Petrochemical Production	0.2	0.1	0.2	0.2	0.3	0.3	0.3
Ferroalloy Production	+	+	+	+	+	+	+
Carbide Production and Consumption	+	+	+	+	+	+	+
Iron and Steel Production & Metallurgical Coke Production	+	+	+	+	+	+	+
N₂O	33.3	24.9	22.2	23.3	22.7	25.8	21.1
Adipic Acid Production	15.2	7.1	4.3	7.0	7.4	10.3	5.3
Nitric Acid Production	12.1	11.3	11.6	10.1	9.3	9.6	10.0
N ₂ O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Caprolactam, Glyoxal, and Glyoxylic Acid Production	1.7	2.1	1.9	1.7	1.5	1.4	1.4
Electronics Industry	+	0.1	0.2	0.2	0.3	0.3	0.2
HFCs	46.5	127.5	168.3	168.1	170.1	169.3	173.8
Substitution of Ozone Depleting Substances ^a	0.2	107.3	163.6	164.9	164.5	165.5	169.7
HCFC-22 Production	46.1	20.0	4.3	2.8	5.2	3.3	3.7
Electronics Industry	0.2	0.2	0.3	0.3	0.4	0.4	0.3
Magnesium Production and Processing	+	+	0.1	0.1	0.1	0.1	0.1
PFCs	24.3	6.7	5.2	4.4	4.1	4.7	4.5
Electronics Industry	2.8	3.3	3.1	2.9	2.9	3.0	2.7
Aluminum Production	21.5	3.4	2.1	1.4	1.1	1.6	1.8
Substitution of Ozone Depleting Substances	0.0	+	+	+	+	0.1	0.1
SF₆	28.8	11.8	5.5	6.0	5.9	5.8	6.0
Electrical Transmission and Distribution	23.2	8.4	3.8	4.1	4.2	3.9	4.2
Magnesium Production and Processing	5.2	2.7	1.0	1.1	1.1	1.1	1.0
Electronics Industry	0.5	0.7	0.7	0.8	0.7	0.8	0.8
NF₃	+	0.5	0.6	0.6	0.6	0.6	0.6
Electronics Industry	+	0.5	0.6	0.6	0.6	0.6	0.6
Unspecified Mix of HFCs, NF₃, PFCs and SF₆	+						
Electronics Industry	+	+	+	+	+	+	+
Total	345.5	365.7	375.4	367.9	367.6	371.0	374.0

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Small amounts of PFC emissions also result from this source.

- 1 Overall, emissions from the IPPU sector increased by 8.2 percent from 1990 to 2019. Significant trends in emissions
- 2 from IPPU source categories (Figure 2-11) over the thirty-year period from 1990 through 2019 included the
- 3 following:
- 4
 - HFC and PFC emissions resulting from the substitution of ODS (e.g., chlorofluorocarbons [CFCs]) have
 - 5 been increasing from small amounts in 1990 to 169.7 MMT CO₂ Eq. in 2019 and accounted for 45.4
 - 6 percent of total IPPU emissions.
 - 7
 - Combined CO₂ and CH₄ emissions from iron and steel production and metallurgical coke production
 - 8 decreased by 3.1 percent to 41.3 MMT CO₂ Eq. from 2018 to 2019, and have declined overall by 63.5
 - 9 MMT CO₂ Eq. (60.6 percent) from 1990 through 2019, due to restructuring of the industry. The trend in
 - 10 the United States has been a shift towards fewer integrated steel mills and more EAFs. EAFs use scrap
 - 11 steel as their main input and generally have less on-site emissions.
 - 12
 - Carbon dioxide emissions from petrochemicals increased by 42.5 percent between 1990 and 2019 from
 - 13 21.6 MMT CO₂ Eq. to 30.8 MMT CO₂ Eq. Emissions are driven primarily from ethylene production and the
 - 14 increase in emissions from 1990 to 2019 is largely driven by an almost doubling of production of ethylene
 - 15 over that time period.


- 1 • Carbon dioxide emission from lime production (13 MMT CO₂ Eq. in 2019) increased 10.8 percent
2 respectively between 1990 and 2019. Emissions have increased as the environmental uses for lime have
3 increased, the most significant use being flue gas treatment at utility power plants.
- 4 • Carbon dioxide emissions from ammonia production (12.3 MMT CO₂ Eq. in 2019) decreased by 5.9
5 percent (0.8 MMT CO₂ Eq.) since 1990. Ammonia production relies on natural gas as both a feedstock and
6 a fuel, and as such, market fluctuations and volatility in natural gas prices affect the production of
7 ammonia from year to year. Emissions from ammonia production have increased steadily since 2016, due
8 to the addition of new ammonia production facilities and new production units at existing facilities.
9 Agricultural demands continue to drive demand for nitrogen fertilizers and the need for new ammonia
10 production capacity.
- 11 • Carbon dioxide emissions from cement production increased by 22.1 percent (7.4 MMT CO₂ Eq.) from
12 1990 through 2019. They rose from 1990 through 2006 and then fell until 2009 due to a decrease in
13 demand for construction materials during the economic recession. Since 2010, CO₂ emissions from
14 cement production have risen 30.0 percent (9.4 MMT CO₂ Eq.).
- 15 • PFC emissions from aluminum production decreased by 91.8 percent (19.7 MMT CO₂ Eq.) from 1990 to
16 2019, due to both industry emission reduction efforts and lower domestic aluminum production.

17 Agriculture

18 Agricultural activities contribute directly to emissions of greenhouse gases through a variety of processes,
19 including the following source categories: enteric fermentation in domestic livestock, livestock manure
20 management, rice cultivation, agricultural soil management, liming, urea fertilization, and field burning of
21 agricultural residues. Methane, N₂O, and CO₂ were the primary greenhouse gases emitted by agricultural activities.


22 In 2019, agricultural activities were responsible for emissions of 628.6 MMT CO₂ Eq., or 9.6 percent of total U.S.
23 greenhouse gas emissions. Methane emissions from enteric fermentation and manure management represented
24 approximately 27.1 percent and 9.4 percent of total CH₄ emissions from anthropogenic activities, respectively, in
25 2019. Agricultural soil management activities, such as application of synthetic and organic fertilizers, deposition of
26 livestock manure, and growing N-fixing plants, were the largest contributors to U.S. N₂O emissions in 2019,
27 accounting for 75.3 percent. Carbon dioxide emissions from the application of crushed limestone and dolomite
28 (i.e., soil liming) and urea fertilization represented 0.1 percent of total CO₂ emissions from anthropogenic
29 activities. Figure 2-12 and Table 2-7 illustrate agricultural greenhouse gas emissions by source.

30 **Figure 2-12: 2019 Agriculture Chapter Greenhouse Gas Sources**



31

1 **Figure 2-13: Trends in Agriculture Chapter Greenhouse Gas Sources**



2
3 **Table 2-7: Emissions from Agriculture (MMT CO₂ Eq.)**

Gas/Source	1990	2005	2015	2016	2017	2018	2019
CO₂	7.1	7.9	8.5	8.0	8.1	7.4	7.8
Urea Fertilization	2.4	3.5	4.7	4.9	5.1	5.2	5.3
Liming	4.7	4.3	3.7	3.1	3.1	2.2	2.4
CH₄	218.2	239.3	241.4	248.1	251.0	255.7	256.4
Enteric Fermentation	164.7	169.3	166.9	172.2	175.8	178.0	178.6
Manure Management	37.1	51.6	57.9	59.6	59.9	61.7	62.4
Rice Cultivation	16.0	18.0	16.2	15.8	14.9	15.6	15.1
Field Burning of Agricultural Residues	0.4	0.4	0.4	0.4	0.4	0.4	0.4
N₂O	330.1	329.9	366.2	348.4	346.4	357.9	364.4
Agricultural Soil Management	315.9	313.4	348.5	330.1	327.6	338.2	344.6
Manure Management	14.0	16.4	17.5	18.1	18.7	19.4	19.6
Field Burning of Agricultural Residues	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Total	555.3	577.1	616.1	604.4	605.5	621.0	628.6

Note: Totals may not sum due to independent rounding.

4 Some significant trends in U.S. emissions from Agriculture source categories (Figure 2-13) over the thirty-year
5 period from 1990 through 2019 included the following:

- 6 • Agricultural soils are the largest anthropogenic source of N₂O emissions in the United States, accounting
7 for approximately 75.3 percent of N₂O emissions in 2019 and 5.2 percent of total emissions in the United
8 States in 2019. Estimated emissions from this source in 2019 were 344.6 MMT CO₂ Eq. Annual N₂O
9 emissions from agricultural soils fluctuated between 1990 and 2019, although overall emissions were 28.7

1 MMT CO₂ Eq. or 9.1 percent higher in 2019 than in 1990. Year-to-year fluctuations are largely a reflection
2 of annual variation in weather patterns, synthetic fertilizer use, and crop production.

- 3 • Enteric fermentation is the largest anthropogenic source of CH₄ emissions in the United States. In 2019,
4 enteric fermentation CH₄ emissions were 27.1 percent of total CH₄ emissions (178.6 MMT CO₂ Eq.), which
5 represents an increase of 13.9 MMT CO₂ Eq. (8.4 percent) since 1990. This increase in emissions from
6 1990 to 2019 in enteric fermentation generally follows the increasing trends in cattle populations. For
7 example, from 1990 to 1995, emissions increased and then generally decreased from 1996 to 2004,
8 mainly due to fluctuations in beef cattle populations and increased digestibility of feed for feedlot cattle.
9 Emissions increased from 2005 to 2007, as both dairy and beef populations increased. Research indicates
10 that the feed digestibility of dairy cow diets decreased during this period. Emissions decreased again from
11 2008 to 2014 as beef cattle populations again decreased. Emissions increased from 2014 to 2019,
12 consistent with an increase in beef cattle population over those same years.
- 13 • Overall, emissions from manure management increased 60.3 percent between 1990 and 2019. This
14 encompassed an increase of 67.9 percent for CH₄, from 37.1 MMT CO₂ Eq. in 1990 to 62.4 MMT CO₂ Eq. in
15 2019; and an increase of 40.2 percent for N₂O, from 14.0 MMT CO₂ Eq. in 1990 to 19.6 MMT CO₂ Eq. in
16 2019. The majority of the increase observed in CH₄ resulted from swine and dairy cattle manure, where
17 emissions increased 48.6 and 117.3 percent, respectively, from 1990 to 2019. From 2018 to 2019, there
18 was a 1.1 percent increase in total CH₄ emissions from manure management, mainly due to minor shifts in
19 the animal populations and the resultant effects on manure management system allocations.
- 20 • Liming and urea fertilization are the only sources of CO₂ emissions reported in the Agriculture sector.
21 Estimated emissions from these sources were 2.4 and 5.3 MMT CO₂ Eq., respectively. Liming emissions
22 increased by 8.6 percent relative to 2018 and decreased 2.3 MMT CO₂ Eq. or 52 percent relative to 1990,
23 while urea fertilization emissions increased by 1.9 percent relative to 2018 and 2.9 MMT CO₂ Eq. or 121.0
24 percent relative to 1990.

25 Land Use, Land-Use Change, and Forestry

26 When humans alter the terrestrial biosphere through land use, changes in land use, and land management
27 practices, they also influence the carbon (C) stock fluxes on these lands and cause emissions of CH₄ and N₂O.
28 Overall, managed land is a net sink for CO₂ (C sequestration) in the United States. The primary driver of fluxes on
29 managed lands is from management of forest lands, but also includes trees in settlements (i.e., urban areas),
30 afforestation, conversion of forest lands to settlements and croplands the management of croplands and
31 grasslands, and the landfilling of yard trimmings and food scraps. The main drivers for net forest sequestration
32 include net forest growth, increasing forest area, and a net accumulation of C stocks in harvested wood pools. The
33 net sequestration in *Settlements Remaining Settlements*, is driven primarily by C stock gains in urban forests (i.e.,
34 Settlement Trees) through net tree growth and increased urban area, as well as long-term accumulation of C in
35 landfills from additions of yard trimmings and food scraps.

36 The LULUCF sector in 2019 resulted in a net increase in C stocks (i.e., net CO₂ removals) of 812.4 MMT CO₂ Eq.
37 (Table 2-8).³ This represents an offset of approximately 12.3 percent of total (i.e., gross) greenhouse gas emissions
38 in 2019. Emissions of CH₄ and N₂O from LULUCF activities in 2019 were 23.4 MMT CO₂ Eq. and represent 0.4
39 percent of total greenhouse gas emissions.⁴ Between 1990 and 2019, total C sequestration in the LULUCF sector

³ LULUCF Carbon Stock Change is the net C stock change from the following categories: *Forest Land Remaining Forest Land*, *Land Converted to Forest Land*, *Cropland Remaining Cropland*, *Land Converted to Cropland*, *Grassland Remaining Grassland*, *Land Converted to Grassland*, *Wetlands Remaining Wetlands*, *Land Converted to Wetlands*, *Settlements Remaining Settlements*, and *Land Converted to Settlements*.

⁴ LULUCF emissions include the CH₄ and N₂O emissions reported for *Peatlands Remaining Peatlands*, *Forest Fires*, *Drained Organic Soils*, *Grassland Fires*, and *Coastal Wetlands Remaining Coastal Wetlands*; CH₄ emissions from *Land Converted to Coastal Wetlands*; and N₂O emissions from *Forest Soils* and *Settlement Soils*.


1 decreased by 10.6 percent, primarily due to a decrease in the rate of net C accumulation in forests and *Cropland*
 2 *Remaining Cropland*, as well as an increase in CO₂ emissions from *Land Converted to Settlements*.

3 Forest fires were the largest source of CH₄ emissions from LULUCF in 2019, totaling 9.5 MMT CO₂ Eq. (379 kt of
 4 CH₄). *Coastal Wetlands Remaining Coastal Wetlands* resulted in CH₄ emissions of 3.8 MMT CO₂ Eq. (153 kt of CH₄).
 5 Grassland fires resulted in CH₄ emissions of 0.3 MMT CO₂ Eq. (12 kt of CH₄). *Land Converted to Wetlands, Drained*
 6 *Organic Soils, and Peatlands Remaining Peatlands* resulted in CH₄ emissions of less than 0.05 MMT CO₂ Eq. each.


7 Forest fires were also the largest source of N₂O emissions from LULUCF in 2019, totaling 6.2 MMT CO₂ Eq. (21 kt of
 8 N₂O). Nitrous oxide emissions from fertilizer application to settlement soils in 2019 totaled to 2.4 MMT CO₂ Eq. (8
 9 kt of N₂O). Additionally, the application of synthetic fertilizers to forest soils in 2019 resulted in N₂O emissions of
 10 0.5 MMT CO₂ Eq. (2 kt of N₂O). Grassland fires resulted in N₂O emissions of 0.3 MMT CO₂ Eq. (1 kt of N₂O). *Coastal*
 11 *Wetlands Remaining Coastal Wetlands* and *Drained Organic Soils* resulted in N₂O emissions of 0.1 MMT CO₂ Eq.
 12 each (0.5 kt of N₂O). *Peatlands Remaining Peatlands* resulted in N₂O emissions of less than 0.05 MMT CO₂ Eq.

13 Carbon dioxide removals from C stock changes are presented (green) in Figure 2-14 and Table 2-8 along with CH₄
 14 and N₂O emissions (purple) for LULUCF source categories.

15 **Figure 2-14: 2019 LULUCF Chapter Greenhouse Gas Sources and Sinks**



1 **Figure 2-15: Trends in Emissions and Removals (Net CO₂ Flux) from Land Use, Land-Use**
 2 **Change, and Forestry^a**



3
 4 ^a In Figure 2-15, the values above stacked bars represent only non-CO₂ LULUCF emission. LULUCF emissions include the CH₄ and
 5 N₂O emissions reported for Peatlands Remaining Peatlands, Forest Fires, Drained Organic Soils, Grassland Fires, and Coastal
 6 Wetlands Remaining Coastal Wetlands; CH₄ emissions from Land Converted to Coastal Wetlands; and N₂O emissions from
 7 Forest Soils and Settlement Soils

8 **Table 2-8: U.S. Greenhouse Gas Emissions and Removals (Net Flux) from Land Use, Land-**
 9 **Use Change, and Forestry (MMT CO₂ Eq.)**

Land-Use Category	1990	2005	2015	2016	2017	2018	2019
Forest Land Remaining Forest Land	(785.9)	(652.8)	(650.6)	(715.7)	(640.9)	(682.4)	(675.5)
Changes in Forest Carbon Stocks ^a	(787.6)	(661.5)	(671.4)	(721.9)	(659.7)	(698.6)	(691.8)
Non-CO ₂ Emissions from Forest Fires ^b	1.5	8.2	20.3	5.6	18.3	15.7	15.7
N ₂ O Emissions from Forest Soils ^c	0.1	0.5	0.5	0.5	0.5	0.5	0.5
Non-CO ₂ Emissions from Drained Organic Soils ^d	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Land Converted to Forest Land	(98.2)	(98.7)	(98.9)	(99.0)	(99.1)	(99.1)	(99.1)
Changes in Forest Carbon Stocks ^e	(98.2)	(98.7)	(98.9)	(99.0)	(99.1)	(99.1)	(99.1)
Cropland Remaining Cropland	(23.2)	(29.0)	(12.8)	(22.7)	(22.3)	(16.6)	(14.5)
Changes in Mineral and Organic Soil Carbon Stocks	(23.2)	(29.0)	(12.8)	(22.7)	(22.3)	(16.6)	(14.5)
Land Converted to Cropland	51.8	52.2	56.1	54.4	54.6	54.3	54.2
Changes in all Ecosystem Carbon Stocks ^f	51.8	52.2	56.1	54.2	54.6	54.3	54.2

Grassland Remaining Grassland	8.5	10.7	13.8	10.4	11.9	12.3	15.1
Changes in Mineral and Organic Soil Carbon Stocks	8.3	10.0	13.1	9.8	11.3	11.7	14.5
Non-CO ₂ Emissions from Grassland Fires ^g	0.2	0.7	0.7	0.6	0.6	0.6	0.6
Land Converted to Grassland	(6.2)	(40.1)	(23.9)	(24.0)	(24.4)	(24.1)	(23.2)
Changes in all Ecosystem Carbon Stocks ^f	(6.2)	(40.1)	(23.9)	(24.0)	(24.4)	(24.1)	(23.2)
Wetlands Remaining Wetlands	(3.5)	(2.6)	(3.8)	(3.8)	(3.7)	(3.7)	(3.7)
Changes in Organic Soil Carbon Stocks in Peatlands	1.1	1.1	0.8	0.7	0.8	0.8	0.8
Changes in Aboveground and Soil Carbon Stocks in Coastal Wetlands	(8.5)	(7.7)	(8.5)	(8.5)	(8.5)	(8.5)	(8.5)
CH ₄ Emissions from Coastal Wetlands Remaining Coastal Wetlands	3.7	3.8	3.8	3.8	3.8	3.8	3.8
N ₂ O Emissions from Coastal Wetlands Remaining Coastal Wetlands	0.1	0.2	0.1	0.1	0.1	0.1	0.1
Non-CO ₂ Emissions from Peatlands Remaining Peatlands	+	+	+	+	+	+	+
Land Converted to Wetlands	0.7	0.7	0.2	0.2	0.2	0.2	0.2
Changes in Aboveground and Soil Carbon Stocks	0.4	0.4	(0.1)	(+)	(+)	(+)	(+)
CH ₄ Emissions from Land Converted to Coastal Wetlands	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Settlements Remaining Settlements	(107.6)	(113.5)	(123.7)	(121.5)	(121.4)	(121.2)	(121.7)
Changes in Organic Soil Carbon Stocks	11.3	12.2	15.7	16.0	16.0	15.9	15.9
Changes in Settlement Tree Carbon Stocks	(96.4)	(117.4)	(130.4)	(129.8)	(129.8)	(129.8)	(129.8)
Changes in Yard Trimming and Food Scrap Carbon Stocks in Landfills	(24.5)	(11.4)	(11.1)	(10.0)	(9.8)	(9.8)	(10.2)
N ₂ O Emissions from Settlement Soils ^h	2.0	3.1	2.2	2.2	2.3	2.4	2.4
Land Converted to Settlements	62.9	85.0	80.1	79.4	79.3	79.3	79.2
Changes in all Ecosystem Carbon Stocks ^f	62.9	85.0	80.1	79.4	79.3	79.3	79.2
LULUCF Emissionsⁱ	7.9	16.8	27.9	13.2	26.0	23.4	23.4
LULUCF Carbon Stock Change^j	(908.7)	(804.8)	(791.5)	(855.8)	(791.8)	(824.6)	(812.4)
LULUCF Sector Net Total^k	(900.8)	(788.0)	(763.6)	(842.6)	(765.8)	(801.2)	(788.9)

Notes: Totals may not sum due to independent rounding. Parentheses indicate net sequestration.

+ Absolute value does not exceed 0.05 MMT CO₂ Eq.

^a Includes the net changes to carbon stocks stored in all forest ecosystem pools and harvested wood products.

^b Estimates include emissions from fires on both *Forest Land Remaining Forest Land* and *Land Converted to Forest Land*.

^c Estimates include emissions from N fertilizer additions on both *Forest Land Remaining Forest Land* and *Land Converted to Forest Land*.

^d Estimates include emissions from drained organic soils on both *Forest Land Remaining Forest Land* and *Land Converted to Forest Land*.

^e Includes the net changes to carbon stocks stored in all forest ecosystem pools.

^f Includes changes in mineral and organic soil carbon stocks for all land use conversions to cropland, grassland, and settlements, respectively. Also includes aboveground/belowground biomass, dead wood, and litter carbon stock changes for conversion of forest land to cropland, grassland, and settlements, respectively.

^g Estimates include emissions from fires on both *Grassland Remaining Grassland* and *Land Converted to Grassland*.

^h Estimates include emissions from N fertilizer additions on both *Settlements Remaining Settlements* and *Land Converted to Settlements* because it is not possible to separate the activity data at this time.

ⁱ LULUCF emissions include the CH₄ and N₂O emissions reported for *Peatlands Remaining Peatlands*, *Forest Fires*, *Drained Organic Soils*, *Grassland Fires*, and *Coastal Wetlands Remaining Coastal Wetlands*; CH₄ emissions from *Land Converted to Coastal Wetlands*; and N₂O emissions from *Forest Soils* and *Settlement Soils*.

^j LULUCF Carbon Stock Change includes any C stock gains and losses from all land use and land use conversion categories.

^k The LULUCF Sector Net Total is the net sum of all LULUCF CH₄ and N₂O emissions to the atmosphere plus net carbon stock changes in units of MMT CO₂ Eq.


1 Other significant trends from 1990 to 2019 in emissions from LULUCF categories (Figure 2-15) over the thirty-year
2 period from 1990 through 2019 included the following:

- 3 • Annual carbon (C) sequestration by forest land (i.e., annual C stock accumulation in the five ecosystem C
4 pools and harvested wood products for *Forest Land Remaining Forest Land* and *Land Converted to Forest
5 Land*) has decreased by approximately 10.7 percent since 1990. This is primarily due to decreased C stock
6 gains in *Land Converted to Forest Land* and the harvested wood products pools within *Forest Land
7 Remaining Forest Land*.
- 8 • Annual C sequestration from *Settlements Remaining Settlements* (which includes organic soils, settlement
9 trees, and landfilled yard trimmings and food scraps) has increased by 13.2 percent over the period from
10 1990 to 2019. This is primarily due to an increase in urbanized land area in the United States with trees
11 growing on it.
- 12 • Annual emissions from *Land Converted to Settlements* increased by approximately 26.0 percent from
13 1990 to 2019 due primarily to C stock losses from *Forest Land Converted to Settlements* and mineral soils
14 C stocks from *Grassland Converted to Settlements*.

15 Waste

16 Waste management and treatment activities are sources of greenhouse gas emissions (see Figure 2-16). In 2019,
17 landfills were the third-largest source of U.S. anthropogenic CH₄ emissions, generating 114.5 MMT CO₂ Eq. and
18 accounting for 17.3 percent of total U.S. CH₄ emissions.⁵ Additionally, wastewater treatment generates emissions
19 of 44.8 MMT CO₂ Eq. and accounts for 27.3 percent of waste emissions, 2.8 percent of U.S. CH₄ emissions, and 5.8
20 percent of U.S. N₂O emissions. Emissions of CH₄ and N₂O from composting are also accounted for in this chapter,
21 generating emissions of 2.3 MMT CO₂ Eq. and 2.0 MMT CO₂ Eq., respectively. Overall, emission sources accounted
22 for in the Waste chapter generated 163.8 MMT CO₂ Eq., or 2.5 percent of total U.S. greenhouse gas emissions in
23 2019. A summary of greenhouse gas emissions from the Waste chapter is presented in Table 2-9.


24 **Figure 2-16: 2019 Waste Sector Greenhouse Gas Sources**



25

⁵ Landfills also store carbon, due to incomplete degradation of organic materials such as wood products and yard trimmings, as described in the Land Use, Land-Use Change, and Forestry chapter.

1 **Figure 2-17: Trends in Waste Chapter Greenhouse Gas Sources**



2

3 **Table 2-9: Emissions from Waste (MMT CO₂ Eq.)**

Gas/Source	1990	2005	2015	2016	2017	2018	2019
CH₄	200.1	153.4	132.6	129.2	130.5	132.9	135.4
Landfills	179.6	131.4	111.4	108.0	109.4	112.1	114.5
Wastewater Treatment	20.2	20.1	18.8	18.7	18.5	18.4	18.4
Composting	0.4	1.9	2.1	2.3	2.4	2.3	2.3
Anaerobic Digester at Biogas Facilities	+	0.1	0.2	0.2	0.2	0.2	0.2
N₂O	19.0	24.6	27.3	27.9	28.6	28.2	28.4
Wastewater Treatment	18.7	23.0	25.4	25.9	26.4	26.1	26.4
Composting	0.3	1.7	1.9	2.0	2.2	2.0	2.0
Total	219.2	178.0	159.8	157.1	159.1	161.1	163.8

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO₂ Eq.

4 Some significant trends in U.S. emissions from waste source categories (Figure 2-17) over the thirty-year period
 5 from 1990 through 2019 included the following:

- 6 • From 1990 to 2019, net CH₄ emissions from landfills decreased by 65.1 MMT CO₂ Eq. (36.2 percent), with
 7 small increases occurring in interim years. This downward trend in emissions coincided with increased
 8 landfill gas collection and control systems, and a reduction of decomposable materials (i.e., paper and
 9 paperboard, food scraps, and yard trimmings) discarded in municipal solid waste (MSW) landfills over the
 10 time series.
- 11 • From 1990 to 2019, CH₄ and N₂O emissions from wastewater treatment decreased by 1.8 MMT CO₂ Eq.
 12 (8.7 percent) and increased by 7.7 MMT CO₂ Eq. (41.0 percent), respectively. Methane emissions from
 13 domestic wastewater treatment have decreased since 1999 due to decreasing percentages of wastewater

1 being treated in anaerobic systems, including reduced use of on-site septic systems and central anaerobic
2 treatment systems. Nitrous oxide emissions from wastewater treatment processes gradually increased
3 across the time series as a result of increasing U.S. population and protein consumption.

- 4 • Combined CH₄ and N₂O emissions from composting have generally increased approximately 3.6 MMT CO₂
5 Eq. since 1990, from 0.7 MMT CO₂ Eq. to 4.3 MMT CO₂ Eq. in 2019, which represents more than a five-fold
6 increase over the time series. The growth in composting since the 1990s is attributable to primarily four
7 factors: (1) the enactment of legislation by state and local governments that discouraged the disposal of
8 yard trimmings and food waste in landfills; (2) yard trimming collection and yard trimming drop off sites
9 provided by local solid waste management districts; (3) an increased awareness of the environmental
10 benefits of composting; and (4) loans or grant programs to establish or expand composting infrastructure.


11 2.2 Emissions by Economic Sector

12 Throughout this report, emission estimates are grouped into five sectors (i.e., chapters) defined by the IPCC and
13 detailed above: Energy, IPPU, Agriculture, LULUCF, and Waste. It is also useful to characterize emissions according
14 to commonly used economic sector categories: residential, commercial, industry, transportation, electric power,
15 and agriculture. Emissions from U.S. Territories are reported as their own end-use sector due to a lack of specific
16 consumption data for the individual end-use sectors within U.S. Territories. See Box 2-1 for more information on
17 how economic sectors are defined. For more information on trends in the Land Use, Land Use Change, and
18 Forestry sector, see Section 2.1.

19 Using this categorization, transportation activities, in aggregate, accounted for the largest portion (28.9 percent) of
20 total U.S. greenhouse gas emissions in 2019. Emissions from electric power accounted for the second largest
21 portion (25.2 percent), while emissions from industry accounted for the third largest portion (22.8 percent) of total
22 U.S. greenhouse gas emissions in 2019. Emissions from industry have in general declined over the past decade due
23 to a number of factors, including structural changes in the U.S. economy (i.e., shifts from a manufacturing-based to
24 a service-based economy), fuel switching, and efficiency improvements.

25 The remaining 23.1 percent of U.S. greenhouse gas emissions were contributed by, in order of magnitude, the
26 agriculture, commercial, and residential sectors, plus emissions from U.S. Territories. Activities related to
27 agriculture accounted for roughly 10.2 percent of emissions; unlike other economic sectors, agricultural sector
28 emissions were dominated by N₂O emissions from agricultural soil management and CH₄ emissions from enteric
29 fermentation, rather than CO₂ from fossil fuel combustion. An increasing amount of carbon is stored in agricultural
30 soils each year, but this CO₂ sequestration is assigned to the LULUCF sector rather than the agriculture economic
31 sector. The commercial and residential sectors accounted for roughly 6.7 percent and 5.8 percent of greenhouse
32 gas emissions, respectively, and U.S. Territories accounted for 0.4 percent of emissions; emissions from these
33 sectors primarily consisted of CO₂ emissions from fossil fuel combustion. Carbon dioxide was also emitted and
34 sequestered (in the form of C) by a variety of activities related to forest management practices, tree planting in
35 urban areas, the management of agricultural soils, landfilling of yard trimmings, and changes in C stocks in coastal
36 wetlands. Table 2-10 presents a detailed breakdown of emissions from each of these economic sectors by source
37 category, as they are defined in this report. Figure 2-18 shows the trend in emissions by sector from 1990 to 2019.

1 **Figure 2-18: U.S. Greenhouse Gas Emissions Allocated to Economic Sectors**



2
 3 Note: Emissions and removals from Land Use, Land Use Change, and Forestry are excluded from figure above. Excludes U.S.
 4 Territories.

5
 6 **Table 2-10: U.S. Greenhouse Gas Emissions Allocated to Economic Sectors (MMT CO₂ Eq. and**
 7 **Percent of Total in 2019)**

Sector/Source	1990	2005	2015	2016	2017	2018	2019	Percent ^a
Transportation	1,527.1	1,986.1	1,825.4	1,867.6	1,877.9	1,883.1	1,902.3	28.9%
CO ₂ from Fossil Fuel Combustion	1,469.1	1,868.8	1,750.5	1,797.3	1,812.6	1,821.2	1,843.2	28.0%
Substitution of Ozone Depleting Substances	+	69.3	46.3	43.3	40.1	38.5	36.7	0.6%
Mobile Combustion	46.1	37.9	17.6	16.7	15.5	14.2	13.5	0.2%
Non-Energy Use of Fuels	11.8	10.2	11.0	10.4	9.6	9.2	8.9	0.1%
Electric Power Industry	1,878.8	2,461.0	1,960.1	1,868.0	1,788.7	1,807.5	1,656.9	25.2%
CO ₂ from Fossil Fuel Combustion	1,820.0	2,400.1	1,900.6	1,808.9	1,732.0	1,752.9	1,606.0	24.4%
Stationary Combustion	20.9	30.9	27.7	27.4	25.9	25.6	22.3	0.3%
Incineration of Waste	11.6	17.8	21.9	22.3	21.7	21.3	20.6	0.3%
Electrical Transmission and Distribution	23.2	8.4	3.8	4.1	4.2	3.9	4.2	0.1%
Other Process Uses of Carbonates	3.1	3.8	6.1	5.5	5.0	3.7	3.7	0.1%
Industry	1,640.9	1,513.3	1,432.5	1,392.0	1,414.6	1,488.9	1,498.9	22.8%
CO ₂ from Fossil Fuel Combustion	810.4	793.1	749.5	747.2	754.5	798.5	798.3	12.1%
Natural Gas Systems	219.7	188.9	177.8	176.8	178.8	184.7	193.3	2.9%
Non-Energy Use of Fuels	97.6	112.8	93.7	82.5	89.4	101.0	100.5	1.5%
Petroleum Systems	58.3	52.7	75.5	62.7	65.7	75.1	83.5	1.3%
Coal Mining	96.5	65.7	61.8	54.3	55.4	53.4	47.4	0.7%
Iron and Steel Production	104.8	70.1	47.9	43.6	40.6	42.6	41.3	0.6%
Cement Production	33.5	46.2	39.9	39.4	40.3	39.0	40.9	0.6%
Substitution of Ozone Depleting Substances	+	8.7	27.6	29.7	31.6	33.1	34.3	0.5%
Petrochemical Production	21.8	27.5	28.2	28.6	29.2	29.6	31.1	0.5%
Lime Production	11.7	14.6	13.3	12.5	12.9	13.1	13.0	0.2%
Ammonia Production	13.0	9.2	10.6	10.2	11.1	12.2	12.3	0.2%
Nitric Acid Production	12.1	11.3	11.6	10.1	9.3	9.6	10.0	0.2%
Abandoned Oil and Gas Wells	6.6	7.0	7.2	7.2	7.1	7.0	7.0	0.1%

Wastewater Treatment (Industrial)	5.8	6.4	6.4	6.6	6.7	6.8	6.9	0.1%
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	4.6	5.1	5.0	6.1	6.6	0.1%
Mobile Combustion	4.0	6.2	5.6	5.7	6.0	6.1	6.3	0.1%
Abandoned Underground Coal Mines	7.2	6.6	6.4	6.7	6.4	6.2	5.9	0.1%
Adipic Acid Production	15.2	7.1	4.3	7.0	7.4	10.3	5.3	0.1%
Carbon Dioxide Consumption	1.5	1.4	4.9	4.6	4.6	4.1	4.9	0.1%
Electronics Industry	3.6	4.8	5.0	5.0	4.9	5.1	4.6	0.1%
N ₂ O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2	0.1%
Stationary Combustion	4.9	4.6	4.2	4.1	4.0	4.1	4.0	0.1%
Other Process Uses of Carbonates	3.1	3.8	6.1	5.5	5.0	3.7	3.7	0.1%
HCFC-22 Production	46.1	20.0	4.3	2.8	5.2	3.3	3.7	0.1%
Aluminum Production	28.3	7.6	4.9	2.7	2.3	3.1	3.6	0.1%
Soda Ash Production	1.4	1.7	1.7	1.7	1.8	1.7	1.8	+
Ferroalloy Production	2.2	1.4	2.0	1.8	2.0	2.1	1.6	+
Titanium Dioxide Production	1.2	1.8	1.6	1.7	1.7	1.5	1.5	+
Caprolactam, Glyoxal, and Glyoxylic Acid Production	1.7	2.1	1.9	1.7	1.5	1.4	1.4	+
Glass Production	1.5	1.9	1.3	1.2	1.3	1.3	1.3	+
Magnesium Production and Processing	5.2	2.7	1.1	1.2	1.2	1.2	1.0	+
Zinc Production	0.6	1.0	0.9	0.9	1.0	1.0	1.0	+
Phosphoric Acid Production	1.5	1.3	1.0	1.0	1.0	0.9	0.9	+
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5	+
Landfills (Industrial)	10.9	14.4	15.0	15.0	15.0	15.0	15.1	0.2%
Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.2	+
Agriculture	600.2	629.7	658.2	645.5	646.3	661.6	669.1	10.2%
N ₂ O from Agricultural Soil Management	315.9	313.4	348.5	330.1	327.6	338.2	344.6	5.2%
Enteric Fermentation	164.7	169.3	166.9	172.2	175.8	178.0	178.6	2.7%
Manure Management	51.1	67.9	75.4	77.7	78.5	81.1	82.0	1.2%
CO ₂ from Fossil Fuel Combustion	43.4	50.9	40.8	39.9	39.5	39.5	39.3	0.6%
Rice Cultivation	16.0	18.0	16.2	15.8	14.9	15.6	15.1	0.2%
Urea Fertilization	2.4	3.5	4.7	4.9	5.1	5.2	5.3	0.1%
Liming	4.7	4.3	3.7	3.1	3.1	2.2	2.4	+
Mobile Combustion	1.5	1.8	1.2	1.2	1.2	1.2	1.2	+
Field Burning of Agricultural Residues	0.5	0.6	0.6	0.6	0.6	0.6	0.6	+
Stationary Combustion	+	+	+	+	+	+	+	+
Commercial	432.2	406.8	425.6	408.9	412.1	448.5	443.9	6.7%
CO ₂ from Fossil Fuel Combustion	228.3	226.0	224.9	210.4	212.3	246.9	238.3	3.6%
Landfills (Municipal)	168.7	117.0	96.4	93.1	94.4	97.0	99.4	1.5%
Substitution of Ozone Depleting Substances	+	22.1	60.8	61.5	61.0	60.8	62.3	0.9%
Wastewater Treatment (Domestic)	33.0	36.6	37.8	38.0	38.2	37.8	37.9	0.6%
Composting	0.7	3.5	4.0	4.3	4.6	4.3	4.3	0.1%
Stationary Combustion	1.5	1.4	1.4	1.4	1.4	1.6	1.5	+
Anaerobic Digestion at Biogas Facilities	+	0.1	0.2	0.2	0.2	0.2	0.2	+
Residential	345.1	371.2	352.3	327.4	331.0	378.2	380.7	5.8%
CO ₂ from Fossil Fuel Combustion	338.6	359.1	318.1	292.3	294.6	339.5	338.8	5.2%

Substitution of Ozone Depleting Substances	0.2	7.2	28.9	30.4	31.8	33.2	36.3	0.6%
Stationary Combustion	6.3	4.9	5.3	4.7	4.5	5.4	5.5	0.1%
U.S. Territories	25.2	63.8	30.2	26.9	25.5	25.4	25.4	0.4%
CO ₂ from Fossil Fuel Combustion	21.7	56.0	29.4	26.2	24.7	24.6	24.6	0.4%
Non-Energy Use of Fuels	3.4	7.6	0.7	0.7	0.7	0.7	0.7	+
Stationary Combustion	0.1	0.2	0.1	0.1	0.1	0.1	0.1	+
Total Emissions (Sources)	6,449.4	7,431.9	6,684.3	6,536.3	6,496.0	6,693.2	6,577.2	100.0%
LULUCF Sector Net Total^b	(900.8)	(788.0)	(763.6)	(842.6)	(765.8)	(801.2)	(788.9)	-12.0%
Net Emissions (Sources and Sinks)	5,548.6	6,643.8	5,920.7	5,693.8	5,730.2	5,892.0	5,788.3	88.0%

Notes: Total emissions presented without LULUCF. Total net emissions presented with LULUCF. Totals may not sum due to independent rounding. Parentheses indicate negative values or sequestration.

+ Does not exceed 0.05 MMT CO₂ Eq. or 0.05 percent.

^a Percent of total (gross) emissions excluding emissions from LULUCF for 2019.

^b The LULUCF Sector Net Total is the net sum of all LULUCF CH₄ and N₂O emissions to the atmosphere plus net carbon stock changes.

1

Box 2-1: Methodology for Aggregating Emissions by Economic Sector

In presenting the Economic Sectors in the annual *Inventory of U.S. Greenhouse Gas Emissions and Sinks*, the Inventory expands upon the standard IPCC sectors common for UNFCCC reporting. Discussing greenhouse gas emissions relevant to U.S.-specific economic sectors improves communication of the report's findings.

The *Electric Power* economic sector includes CO₂, CH₄ and N₂O emissions from the combustion of fossil fuels that are included in the EIA electric power sector. Carbon dioxide, CH₄, and N₂O emissions from waste incineration are included in the Electric Power economic sector, as the majority of municipal solid waste is combusted in plants that produce electricity. The Electric Power economic sector also includes SF₆ from Electrical Transmission and Distribution, and a portion of CO₂ from Other Process Uses of Carbonates (from pollution control equipment installed in electric power plants).

The *Transportation* economic sector includes CO₂ emissions from the combustion of fossil fuels that are included in the EIA transportation fuel-consuming sector. (Additional analyses and refinement of the EIA data are further explained in the Energy chapter of this report.) Emissions of CH₄ and N₂O from mobile combustion are also apportioned to the Transportation economic sector based on the EIA transportation fuel-consuming sector. Substitution of Ozone Depleting Substances emissions are apportioned to the Transportation economic sector based on emissions from refrigerated transport and motor vehicle air-conditioning systems. Finally, CO₂ emissions from Non-Energy Uses of Fossil Fuels identified as lubricants for transportation vehicles are included in the Transportation economic sector.

The *Industry* economic sector includes CO₂ emissions from the combustion of fossil fuels that are included in the EIA industrial fuel-consuming sector, minus the agricultural use of fuel explained below. The CH₄ and N₂O emissions from stationary and mobile combustion are also apportioned to the Industry economic sector based on the EIA industrial fuel-consuming sector, minus emissions apportioned to the Agriculture economic sector. Substitution of Ozone Depleting Substances emissions are apportioned based on their specific end-uses within the source category, with most emissions falling within the Industry economic sector. Finally, CH₄ emissions from industrial landfills and CH₄ and N₂O from industrial wastewater treatment are included in the Industry economic sector.

Additionally, all process-related emissions from sources with methods considered within the IPCC IPPU sector are apportioned to the Industry economic sector. This includes the process-related emissions (i.e., emissions from the actual process to make the material, not from fuels to power the plant) from activities such as Cement Production, Iron and Steel Production and Metallurgical Coke Production, and Ammonia Production. Additionally, fugitive emissions from energy production sources, such as Natural Gas Systems, Coal Mining, and Petroleum Systems are included in the Industry economic sector. A portion of CO₂ from Other Process Uses of Carbonates (from pollution control equipment installed in large industrial facilities) is also included in the Industry economic sector. Finally, all remaining CO₂ emissions from Non-Energy Uses of Fossil Fuels are assumed

to be industrial in nature (besides the lubricants for transportation vehicles specified above) and are attributed to the Industry economic sector.

The *Agriculture* economic sector includes CO₂ emissions from the combustion of fossil fuels that are based on supplementary sources of agriculture fuel use data, because EIA does not include an agriculture fuel-consuming sector. Agriculture equipment is included in the EIA industrial fuel-consuming sector. Agriculture fuel use estimates are obtained from U.S. Department of Agriculture survey data, in combination with separate EIA fuel sales reports (USDA 2019; EIA 2020a). These supplementary data are subtracted from the industrial fuel use reported by EIA to obtain agriculture fuel use. CO₂ emissions from fossil fuel combustion, and CH₄ and N₂O emissions from stationary and mobile combustion, are then apportioned to the Agriculture economic sector based on agricultural fuel use.

The other IPCC Agriculture emission source categories apportioned to the Agriculture economic sector include N₂O emissions from Agricultural Soils, CH₄ from Enteric Fermentation, CH₄ and N₂O from Manure Management, CH₄ from Rice Cultivation, CO₂ emissions from Liming and Urea Application, and CH₄ and N₂O from Field Burning of Agricultural Residues.

The *Residential* economic sector includes CO₂ emissions from the combustion of fossil fuels that are included in the EIA residential fuel-consuming sector. Stationary combustion emissions of CH₄ and N₂O are also based on the EIA residential fuel-consuming sector. Substitution of Ozone Depleting Substances are apportioned to the Residential economic sector based on emissions from residential air-conditioning systems. Nitrous oxide emissions from the application of fertilizers to developed land (termed “settlements” by the IPCC) are also included in the Residential economic sector.

The *Commercial* economic sector includes CO₂ emissions from the combustion of fossil fuels that are included in the EIA commercial fuel-consuming sector. Emissions of CH₄ and N₂O from Mobile Combustion are also apportioned to the Commercial economic sector based on the EIA commercial fuel-consuming sector. Substitution of Ozone Depleting Substances emissions are apportioned to the Commercial economic sector based on emissions from commercial refrigeration/air-conditioning systems. Public works sources, including direct CH₄ from municipal landfills, CH₄ from anaerobic digestion at biogas facilities, CH₄ and N₂O from domestic wastewater treatment, and composting, are also included in the Commercial economic sector.

1

2 Emissions with Electricity Distributed to Economic Sectors

3 It is also useful to view greenhouse gas emissions from economic sectors with emissions related to electric power
4 distributed into end-use categories (i.e., emissions from electric power are allocated to the economic sectors in
5 which the electricity is used).

6 The generation, transmission, and distribution of electricity accounted for 25.2 percent of total U.S. greenhouse
7 gas emissions in 2019. Electric power-related emissions decreased by 11.8 percent since 1990 and by 8.3 percent
8 from 2018 to 2019, primarily due to a significantly colder winter and a hotter summer in 2019 compared to 2018,
9 which increased the amount of energy required for heating and cooling. Between 2018 to 2019, the consumption
10 of natural gas for electric power generation increased by 6.7, while the consumption of coal and petroleum
11 decreased by 15.5 and 27.6 percent, respectively, reflecting a continued shift from coal to natural gas for electricity
12 generation.

13 From 2018 to 2019, electricity sales to the residential and commercial end-use sectors decreased by 2.0 percent
14 and 1.5 percent, respectively. Electricity sales to the industrial sector increased by approximately 0.2 percent.
15 Overall, from 2018 to 2019, the amount of electricity retail sales (in kWh) decreased by 1.2 percent. Table 2-11
16 provides a detailed summary of emissions from electric power-related activities.

17 **Table 2-11: Electric Power-Related Greenhouse Gas Emissions (MMT CO₂ Eq.)**

Gas/Fuel Type or Source	1990	2005	2015	2016	2017	2018	2019
-------------------------	------	------	------	------	------	------	------

CO₂	1,834.3	2,421.3	1,928.2	1,836.2	1,758.3	1,777.6	1,630.0
Fossil Fuel Combustion	1,820.0	2,400.1	1,900.6	1,808.9	1,732.0	1,752.9	1,606.0
<i>Coal</i>	<i>1,546.5</i>	<i>1,982.8</i>	<i>1,351.4</i>	<i>1,242.0</i>	<i>1,207.1</i>	<i>1,152.9</i>	<i>973.5</i>
<i>Natural Gas</i>	<i>175.4</i>	<i>318.9</i>	<i>525.2</i>	<i>545.0</i>	<i>505.6</i>	<i>577.4</i>	<i>616.0</i>
<i>Petroleum</i>	<i>97.5</i>	<i>98.0</i>	<i>23.7</i>	<i>21.5</i>	<i>18.9</i>	<i>22.2</i>	<i>16.2</i>
<i>Geothermal</i>	<i>0.5</i>	<i>0.5</i>	<i>0.4</i>	<i>0.4</i>	<i>0.4</i>	<i>0.4</i>	<i>0.4</i>
Incineration of Waste	11.2	17.4	21.5	21.8	21.3	20.9	20.2
Other Process Uses of							
Carbonates	3.1	3.8	6.1	5.5	5.0	3.7	3.7
CH₄	0.4	0.9	1.2	1.2	1.1	1.2	1.3
Stationary Sources ^a	0.4	0.9	1.2	1.2	1.1	1.2	1.3
Incineration of Waste	+	+	+	+	+	+	+
N₂O	21.0	30.4	26.9	26.6	25.2	24.8	21.5
Stationary Sources ^a	20.5	30.1	26.5	26.2	24.8	24.4	21.1
Incineration of Waste	0.5	0.4	0.4	0.4	0.4	0.4	0.4
SF₆	23.2	8.4	3.8	4.1	4.2	3.9	4.2
Electrical Transmission and							
Distribution	23.2	8.4	3.8	4.1	4.2	3.9	4.2
Total	1,878.8	2,461.0	1,960.1	1,868.0	1,788.7	1,807.5	1,656.9

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO₂ Eq.


^a Includes only stationary combustion emissions related to the generation of electricity.

1 To distribute electricity emissions among economic end-use sectors, emissions from the source categories
2 assigned to the electric power sector were allocated to the residential, commercial, industry, transportation, and
3 agriculture economic sectors according to each economic sector's share of retail sales of electricity (EIA 2020b;
4 Duffield 2006). These source categories include CO₂ from Fossil Fuel Combustion, CH₄ and N₂O from Stationary
5 Combustion, Incineration of Waste, Other Process Uses of Carbonates, and SF₆ from Electrical Transmission and
6 Distribution Systems. Note that only 50 percent of the Other Process Uses of Carbonates emissions were
7 associated with electric power and distributed as described; the remainder of Other Process Uses of Carbonates
8 emissions were attributed to the industrial processes economic end-use sector.⁶

9 When emissions from electricity use are distributed among these economic end-use sectors, industrial activities
10 account for the largest share of total U.S. greenhouse gas emissions (29.6 percent), followed closely by emissions
11 from transportation (29.0 percent). Emissions from the commercial and residential sectors also increase
12 substantially when emissions from electricity are included (15.4 and 14.9 percent, respectively). In all economic
13 end-use sectors except agriculture, CO₂ accounts for more than 78.8 percent of greenhouse gas emissions,
14 primarily from the combustion of fossil fuels. Table 2-12 presents a detailed breakdown of emissions from each of
15 these economic sectors, with emissions from electric power distributed to them. Figure 2-14 shows the trend in
16 these emissions by sector from 1990 to 2019.

⁶ Emissions were not distributed to U.S. Territories, since the electric power sector only includes emissions related to the generation of electricity in the 50 states and the District of Columbia.

1 **Figure 2-19: U.S. Greenhouse Gas Emissions with Electricity-Related Emissions Distributed**
 2 **to Economic Sectors**



3
 4 Note: Emissions and removals from Land Use, Land Use Change, and Forestry are excluded from figure above. Excludes U.S.
 5 Territories.

6 **Table 2-12: U.S. Greenhouse Gas Emissions by Economic Sector and Gas with Electricity-**
 7 **Related Emissions Distributed (MMT CO₂ Eq.) and Percent of Total in 2019**

Sector/Gas	1990	2005	2015	2016	2017	2018	2019	Percent ^a
Industry	2,314.4	2,229.9	1,957.8	1,887.1	1,896.6	1,966.4	1,943.6	29.6%
Direct Emissions	1,640.9	1,513.3	1,432.5	1,392.0	1,414.6	1,488.9	1,498.9	22.8%
CO ₂	1,158.9	1,133.4	1,072.2	1,041.3	1,058.9	1,130.7	1,142.9	17.4%
CH ₄	365.4	306.4	290.3	279.3	281.1	279.9	281.1	4.3%
N ₂ O	40.3	33.8	30.2	31.4	30.9	34.3	29.6	0.5%
HFCs, PFCs, SF ₆ , and NF ₃	76.3	39.6	39.8	39.9	43.6	44.0	45.2	0.7%
Electricity-Related	673.5	716.6	525.3	495.2	482.0	477.5	444.7	6.8%
CO ₂	657.5	705.1	516.7	486.7	473.8	469.6	437.5	6.7%
CH ₄	0.2	0.3	0.3	0.3	0.3	0.3	0.3	+
N ₂ O	7.5	8.9	7.2	7.1	6.8	6.6	5.8	0.1%
SF ₆	8.3	2.4	1.0	1.1	1.1	1.0	1.1	+
Transportation	1,530.2	1,990.9	1,829.8	1,872.0	1,882.3	1,887.9	1,907.2	29.0%
Direct Emissions	1,527.1	1,986.1	1,825.4	1,867.6	1,877.9	1,883.1	1,902.3	28.9%
CO ₂	1,480.9	1,878.9	1,761.5	1,807.6	1,822.2	1,830.4	1,852.1	28.2%
CH ₄	5.7	2.9	1.8	1.6	1.6	1.5	1.4	+
N ₂ O	40.4	35.0	15.9	15.0	14.0	12.7	12.1	0.2%
HFCs ^b	+	69.3	46.3	43.3	40.1	38.5	36.7	0.6%
Electricity-Related	3.1	4.8	4.4	4.3	4.5	4.8	4.9	0.1%
CO ₂	3.1	4.8	4.3	4.3	4.4	4.7	4.8	0.1%
CH ₄	+	+	+	+	+	+	+	+
N ₂ O	+	0.1	0.1	0.1	0.1	0.1	0.1	+
SF ₆	+	+	+	+	+	+	+	+
Commercial	987.3	1,230.2	1,109.4	1,063.2	1,038.1	1,072.2	1,013.9	15.4%
Direct Emissions	432.2	406.8	425.6	408.9	412.1	448.5	443.9	6.7%
CO ₂	228.3	226.0	224.9	210.4	212.3	246.9	238.3	3.6%
CH ₄	184.9	134.2	112.9	109.3	110.5	112.9	115.2	1.8%
N ₂ O	19.0	24.5	27.0	27.7	28.3	27.9	28.2	0.4%
HFCs	+	22.1	60.8	61.5	61.0	60.8	62.3	0.9%

Electricity-Related	555.1	823.4	683.8	654.4	626.0	623.7	569.9	8.7%
CO ₂	542.0	810.1	672.7	643.2	615.4	613.3	560.7	8.5%
CH ₄	0.1	0.3	0.4	0.4	0.4	0.4	0.4	+
N ₂ O	6.2	10.2	9.4	9.3	8.8	8.6	7.4	0.1%
SF ₆	6.8	2.8	1.3	1.4	1.5	1.3	1.5	+
Residential	957.0	1,248.9	1,057.5	1,002.3	968.3	1,040.3	982.7	14.9%
Direct Emissions	345.1	371.2	352.3	327.4	331.0	378.2	380.7	5.8%
CO ₂	338.6	359.1	318.1	292.3	294.6	339.5	338.8	5.2%
CH ₄	5.2	4.1	4.5	3.9	3.8	4.5	4.6	0.1%
N ₂ O	1.0	0.9	0.9	0.8	0.8	0.9	0.9	+
HFCs	0.2	7.2	28.9	30.4	31.8	33.2	36.3	0.6%
Electricity-Related	611.9	877.7	705.2	674.8	637.3	662.1	602.0	9.2%
CO ₂	597.4	863.6	693.7	663.3	626.5	651.1	592.2	9.0%
CH ₄	0.1	0.3	0.4	0.4	0.4	0.4	0.5	+
N ₂ O	6.8	10.9	9.7	9.6	9.0	9.1	7.8	0.1%
SF ₆	7.5	3.0	1.4	1.5	1.5	1.4	1.5	+
Agriculture	635.3	668.1	699.6	684.8	685.2	701.0	704.5	10.7%
Direct Emissions	600.2	629.7	658.2	645.5	646.3	661.6	669.1	10.2%
CO ₂	50.5	58.7	49.3	47.8	47.6	46.9	47.1	0.7%
CH ₄	218.3	239.5	241.5	248.2	251.1	255.8	256.5	3.9%
N ₂ O	331.4	331.5	367.3	349.5	347.5	358.9	365.5	5.6%
Electricity-Related	35.1	38.4	41.5	39.3	39.0	39.4	35.4	0.5%
CO ₂	34.3	37.8	40.8	38.7	38.3	38.7	34.8	0.5%
CH ₄	+	+	+	+	+	+	+	+
N ₂ O	0.4	0.5	0.6	0.6	0.5	0.5	0.5	+
SF ₆	0.4	0.1	0.1	0.1	0.1	0.1	0.1	+
U.S. Territories	25.2	63.8	30.2	26.9	25.5	25.4	25.4	0.4%
Total Emissions (Sources)	6,449.4	7,431.9	6,684.3	6,536.3	6,496.0	6,693.2	6,577.2	100.0%
LULUCF Sector Net Total^c	(900.8)	(788.0)	(763.6)	(842.6)	(765.8)	(801.2)	(788.9)	(+)%
Net Emissions (Sources and Sinks)	5,548.6	6,643.8	5,920.7	5,693.8	5,730.2	5,892.0	5,788.3	88.0%

Notes: Total emissions presented without LULUCF. Net emissions presented with LULUCF. Emissions from electric power are allocated based on aggregate electricity use in each end-use sector. Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO₂ Eq. or 0.05 percent.

^a Percent of total (gross) emissions excluding emissions from LULUCF for year 2019.

^b Includes primarily HFC-134a.

^c The LULUCF Sector Net Total is the net sum of all LULUCF CH₄ and N₂O emissions to the atmosphere plus net carbon stock changes.

1 Industry

2 The industry end-use sector includes CO₂ emissions from fossil fuel combustion from all manufacturing facilities, in
3 aggregate, and with the distribution of electricity-related emissions, accounts for 29.6 percent of U.S. greenhouse
4 gas emissions in 2019. This end-use sector also includes emissions that are produced as a byproduct of the non-
5 energy-related industrial process activities. The variety of activities producing these non-energy-related emissions
6 includes CH₄ emissions from petroleum and natural gas systems, fugitive CH₄ emissions from coal mining,
7 byproduct CO₂ emissions from cement manufacture, and HFC, PFC, SF₆, and NF₃ byproduct emissions from the
8 electronics industry, to name a few.

9 Since 1990, industrial sector emissions have declined by 16.0 percent. The decline has occurred both in direct
10 emissions and indirect emissions associated with electricity use. Structural changes within the U.S. economy that
11 led to shifts in industrial output away from energy-intensive manufacturing products to less energy-intensive
12 products (e.g., from steel to computer equipment) have had a significant effect on industrial emissions.

1 Transportation

2 When electricity-related emissions are distributed to economic end-use sectors, transportation activities
3 accounted for 29.0 percent of U.S. greenhouse gas emissions in 2019. The largest sources of transportation
4 greenhouse gas emissions in 2019 were passenger cars (41.0 percent); freight trucks (23.2 percent); light-duty
5 trucks, which include sport utility vehicles, pickup trucks, and minivans (17.3 percent); commercial aircraft (6.9
6 percent); pipelines (2.8 percent); other aircraft (2.6 percent); ships and boats (2.2 percent); and rail (2.1 percent).
7 These figures include direct CO₂, CH₄, and N₂O emissions from fossil fuel combustion used in transportation,
8 indirect emissions from electricity use and emissions from non-energy use (i.e., lubricants) used in transportation,
9 as well as HFC emissions from mobile air conditioners and refrigerated transport allocated to these vehicle types.

10 In terms of the overall trend, from 1990 to 2019, total transportation emissions increased due, in large part, to
11 increased demand for travel. The number of VMT by light-duty motor vehicles (passenger cars and light-duty
12 trucks) increased 47.4 percent from 1990 to 2019, as a result of a confluence of factors including population
13 growth, economic growth, urban sprawl, and periods of low fuel prices.

14 The decline in new light-duty vehicle fuel economy between 1990 and 2004 reflected the increasing market share
15 of light-duty trucks, which grew from about 30 percent of new vehicle sales in 1990 to 48 percent in 2004. Starting
16 in 2005, average new vehicle fuel economy began to increase while light-duty VMT grew only modestly for much
17 of the period. Light-duty VMT grew by less than one percent or declined each year between 2005 and 2013,⁷ then
18 grew at a faster rate until 2016 (2.6 percent from 2014 to 2015, and 2.5 percent from 2015 to 2016). Since 2016,
19 the rate of light-duty VMT growth has slowed to less than one percent each year. Average new vehicle fuel
20 economy has increased almost every year since 2005, while light-duty truck market share decreased to about 33
21 percent in 2009 and has since varied from year to year between 36 and 56 percent. Light-duty truck market share
22 was about 56 percent of new vehicles in model year 2019 (EPA 2020a).

23 Table 2-13 provides a detailed summary of greenhouse gas emissions from transportation-related activities with
24 electricity-related emissions included in the totals. Historically, the majority of electricity use in the transportation
25 sector was for rail transport. However, more recently there has been increased electricity use in on-road electric
26 and plug-in hybrid vehicles. For a more detailed breakout of emissions by fuel type by vehicle see Table A106 in
27 Annex 3.


28 Almost all of the energy used for transportation was supplied by petroleum-based products, with more than half
29 being related to gasoline consumption in automobiles and other highway vehicles. Other fuel uses, especially
30 diesel fuel for freight trucks and jet fuel for aircraft, accounted for the remainder. The primary driver of
31 transportation-related emissions was CO₂ from fossil fuel combustion, which increased by 26 percent from 1990 to
32 2019.⁸ This rise in CO₂ emissions, combined with an increase in HFCs from close to zero emissions in 1990 to 36.7
33 MMT CO₂ Eq. in 2019, led to an increase in overall greenhouse gas emissions from transportation activities of 25
34 percent.⁹

⁷ VMT estimates are based on data from FHWA Highway Statistics Table VM-1 (FHWA 1996 through 2018). In 2007 and 2008 light-duty VMT decreased 3.0 percent and 2.3 percent, respectively. Note that the decline in light-duty VMT from 2006 to 2007 is due at least in part to a change in FHWA's methods for estimating VMT. In 2011, FHWA changed its methods for estimating VMT by vehicle class, which led to a shift in VMT and emissions among on-road vehicle classes in the 2007 to 2018 time period. In absence of these method changes, light-duty VMT growth between 2006 and 2007 would likely have been higher.

⁸ See previous footnote.

⁹ See previous footnote.

1 **Figure 2-20: Trends in Transportation-Related Greenhouse Gas Emissions¹⁰**



2

3 **Table 2-13: Transportation-Related Greenhouse Gas Emissions (MMT CO₂ Eq.)**

Gas/Vehicle	1990	2005	2015	2016	2017	2018	2019
Passenger Cars	639.6	697.6	775.8	787.6	784.9	777.5	782.1
CO ₂	612.2	647.3	752.4	766.8	766.9	761.5	768.0
CH ₄	3.2	1.3	0.6	0.6	0.5	0.5	0.4
N ₂ O	24.1	17.3	8.1	7.1	6.1	5.1	4.4
HFCs	+	31.7	14.7	13.2	11.4	10.4	9.3
Light-Duty Trucks	326.7	542.0	329.8	339.7	333.7	328.3	329.5
CO ₂	312.2	494.2	303.3	315.6	311.9	308.0	310.6
CH ₄	1.7	0.8	0.3	0.2	0.2	0.2	0.2
N ₂ O	12.8	13.6	3.2	2.9	2.4	2.0	1.8
HFCs	+	33.3	23.0	21.1	19.2	18.1	16.9
Medium- and Heavy-Duty Trucks	230.3	404.0	412.9	420.8	428.2	437.5	443.1
CO ₂	229.3	399.3	404.9	412.6	419.6	428.6	433.7
CH ₄	0.3	0.1	0.1	0.1	0.1	0.1	0.1
N ₂ O	0.7	1.2	2.4	2.6	2.8	2.9	3.2
HFCs	+	3.4	5.5	5.5	5.7	5.9	6.1
Buses	8.5	12.3	19.5	19.2	20.4	21.9	22.2

¹⁰ In 2011 FHWA changed its methods for estimating VMT and related data. These methodological changes included how vehicles are classified, moving from a system based on body-type to one that is based on wheelbase. These changes were first incorporated for the 1990 through 2008 Inventory and apply to the 2007 to 2019 time period. This resulted in large changes in VMT data by vehicle class, leading to a shift in emissions among on-road vehicle classes. This change in vehicle classification has moved some smaller trucks and sport utility vehicles from the light truck category to the passenger vehicle category in this Inventory.

CO ₂	8.4	11.8	18.8	18.5	19.7	21.1	21.4
CH ₄	+	0.2	0.2	0.2	0.2	0.2	0.2
N ₂ O	+	+	0.1	0.1	0.1	0.1	0.1
HFCs	+	0.3	0.4	0.4	0.4	0.4	0.4
Motorcycles	1.7	1.6	3.8	4.0	3.9	3.9	3.9
CO ₂	1.7	1.6	3.7	3.9	3.8	3.8	3.8
CH ₄	+	+	+	+	+	+	+
N ₂ O	+	+	+	+	+	+	+
Commercial Aircraft^a	110.9	134.0	120.1	121.5	129.2	130.8	130.8
CO ₂	109.9	132.7	119.0	120.4	128.0	129.6	129.6
CH ₄	+	+	+	+	+	+	+
N ₂ O	1.0	1.2	1.1	1.1	1.2	1.2	1.2
Other Aircraft^b	78.3	59.7	40.4	47.5	45.6	44.7	50.2
CO ₂	77.5	59.1	40.0	47.0	45.2	44.3	49.7
CH ₄	0.1	0.1	+	+	+	+	+
N ₂ O	0.7	0.5	0.4	0.4	0.4	0.4	0.5
Ships and Boats^c	47.4	45.9	34.1	41.4	44.4	41.4	42.4
CO ₂	46.3	44.4	30.8	37.6	40.2	36.9	37.6
CH ₄	0.3	0.4	0.4	0.4	0.4	0.4	0.4
N ₂ O	0.8	0.7	0.3	0.5	0.6	0.5	0.6
HFCs	+	0.5	2.6	2.9	3.3	3.6	3.9
Rail	39.0	51.4	43.9	40.5	41.1	42.8	40.4
CO ₂	38.5	50.7	43.3	39.9	40.5	42.2	39.7
CH ₄	0.1	0.1	0.1	0.1	0.1	0.1	0.1
N ₂ O	0.3	0.4	0.4	0.3	0.4	0.4	0.3
HFCs	+	0.1	0.1	0.1	0.1	0.1	0.1
Other Emissions from Electric Power ^d	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Pipelines^e	36.0	32.4	38.5	39.2	41.3	49.9	53.7
CO ₂	36.0	32.4	38.5	39.2	41.3	49.9	53.7
Lubricants	11.8	10.2	11.0	10.4	9.6	9.2	8.9
CO ₂	11.8	10.2	11.0	10.4	9.6	9.2	8.9
Total Transportation	1,530.2	1,990.9	1,829.8	1,872.0	1,882.3	1,887.9	1,907.2
<i>International Bunker Fuels^f</i>	<i>54.8</i>	<i>44.7</i>	<i>31.6</i>	<i>35.0</i>	<i>34.6</i>	<i>32.5</i>	<i>26.4</i>
<i>Ethanol CO₂^g</i>	<i>4.1</i>	<i>21.8</i>	<i>76.6</i>	<i>79.4</i>	<i>80.2</i>	<i>78.6</i>	<i>80.2</i>
<i>Biodiesel CO₂^g</i>	<i>+</i>	<i>0.9</i>	<i>14.1</i>	<i>19.6</i>	<i>18.7</i>	<i>17.9</i>	<i>17.1</i>

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Consists of emissions from jet fuel consumed by domestic operations of commercial aircraft (no bunkers).

^b Consists of emissions from jet fuel and aviation gasoline consumption by general aviation and military aircraft.

^c Fluctuations in emission estimates are associated with fluctuations in reported fuel consumption and may reflect issues with data sources.

^d Other emissions from electric power are a result of waste incineration (as the majority of municipal solid waste is combusted in “trash-to-steam” electric power plants), electrical transmission and distribution, and a portion of Other Process Uses of Carbonates (from pollution control equipment installed in electric power plants).

^e CO₂ estimates reflect natural gas used to power pipelines, but not electricity. While the operation of pipelines produces CH₄ and N₂O, these emissions are not directly attributed to pipelines in the Inventory.

^f Emissions from International Bunker Fuels include emissions from both civilian and military activities; these emissions are not included in the transportation totals.

^g Ethanol and biodiesel CO₂ estimates are presented for informational purposes only. See Section 3.11 and the estimates in Land Use, Land-Use Change, and Forestry (see Chapter 6), in line with IPCC methodological guidance and UNFCCC reporting obligations, for more information on ethanol and biodiesel.

Notes: Passenger cars and light-duty trucks include vehicles typically used for personal travel and less than 8,500 lbs; medium- and heavy-duty trucks include vehicles larger than 8,500 lbs. HFC emissions primarily reflect HFC-134a. Totals may not sum due to independent rounding.

1 Commercial

2 The commercial end-use sector, with electricity-related emissions distributed, accounts for 15.4 percent of U.S.
3 greenhouse gas emissions in 2019 and is heavily reliant on electricity for meeting energy needs, with electricity use
4 for lighting, heating, air conditioning, and operating appliances. The remaining emissions were largely due to the
5 direct consumption of natural gas and petroleum products, primarily for heating and cooking needs. Energy-
6 related emissions from the commercial sector have generally been increasing since 1990, and annual variations are
7 often correlated with short-term fluctuations in energy use caused by weather conditions, rather than prevailing
8 economic conditions. Decreases in energy-related emissions in the commercial sector in recent years can be
9 largely attributed to an overall reduction in energy use driven by a reduction in heating degree days and increases
10 in energy efficiency.

11 Landfills and wastewater treatment are included in the commercial sector, with landfill emissions decreasing since
12 1990 and wastewater treatment emissions decreasing slightly.

13 Residential

14 The residential end-use sector, with electricity-related emissions distributed, accounts for 14.9 percent of U.S.
15 greenhouse gas emissions in 2019 and similarly, is heavily reliant on electricity for meeting energy needs, with
16 electricity use for lighting, heating, air conditioning, and operating appliances. The remaining emissions were
17 largely due to the direct consumption of natural gas and petroleum products, primarily for heating and cooking
18 needs. Emissions from the residential sector have generally been increasing since 1990, and annual variations are
19 often correlated with short-term fluctuations in energy use caused by weather conditions, rather than prevailing
20 economic conditions. In the long term, the residential sector is also affected by population growth, migration
21 trends toward warmer areas, and changes in housing and building attributes (e.g., larger sizes and improved
22 insulation). A shift toward energy-efficient products and more stringent energy efficiency standards for household
23 equipment has also contributed to recent trends in energy demand in households (EIA 2018).

24 Agriculture

25 The agriculture end-use sector accounts for 10.7 percent of U.S. greenhouse gas emissions in 2019 when
26 electricity-related emissions are distributed, and includes a variety of processes, including enteric fermentation in
27 domestic livestock, livestock manure management, and agricultural soil management. In 2019, agricultural soil
28 management was the largest source of N₂O emissions, and enteric fermentation was the largest source of CH₄
29 emissions in the United States. This sector also includes small amounts of CO₂ emissions from fossil fuel
30 combustion by motorized farm equipment such as tractors.

31 **Box 2-2: Trends in Various U.S. Greenhouse Gas Emissions-Related Data**

Total greenhouse gas emissions can be compared to other economic and social indices to highlight changes over time. These comparisons include: (1) emissions per unit of aggregate energy use, because energy-related activities are the largest sources of emissions; (2) emissions per unit of fossil fuel consumption, because almost all energy-related emissions involve the combustion of fossil fuels; (3) emissions per unit of total gross domestic product as a measure of national economic activity; and (4) emissions per capita.

Table 2-14 provides data on various statistics related to U.S. greenhouse gas emissions normalized to 1990 as a baseline year. These values represent the relative change in each statistic since 1990. Greenhouse gas emissions in the United States have grown at an average annual rate of 0.1 percent since 1990, although changes from year to year have been significantly larger. This growth rate is slightly slower than that for total energy use, overall gross domestic product (GDP) and national population (see Table 2-14 and Figure 2-21). The direction of these trends started to change after 2005, when greenhouse gas emissions, total energy use and associated fossil fuel consumption began to peak. Greenhouse gas emissions in the United States have decreased at an average annual rate of 0.8 percent since 2005. Fossil fuel consumption has also decreased at a slower rate than

emissions since 2005, while total energy use, GDP, and national population continued to increase.

Table 2-14: Recent Trends in Various U.S. Data (Index 1990 = 100)

Variable	1990	2005	2014	2015	2016	2017	2018	Avg. Annual Change Since 1990 ^a	Avg. Annual Change Since 2005 ^a
Greenhouse Gas Emissions ^b	100	115	104	101	101	104	102	0.1%	-0.8%
Energy Use ^c	100	119	116	116	116	120	119	0.6%	0.1%
GDP ^d	100	159	186	189	194	200	204	2.5%	1.8%
Population ^e	100	118	128	129	130	131	132	1.0%	0.8%

^a Average annual growth rate.


^b GWP-weighted values.

^c Energy-content-weighted values (EIA 2020b).

^d GDP in chained 2009 dollars (BEA 2020).

^e U.S. Census Bureau (2020).

Figure 2-21: U.S. Greenhouse Gas Emissions Per Capita and Per Dollar of Gross Domestic Product



Source: BEA (2020), U.S. Census Bureau (2020), and emission estimates in this report.

1

2

3

2.3 Precursor Greenhouse Gas Emissions (CO, NO_x, NMVOCs, and SO₂)

4

The reporting requirements of the UNFCCC¹¹ request that information be provided on precursor greenhouse gases, which include carbon monoxide (CO), nitrogen oxides (NO_x), non-CH₄ volatile organic compounds

5

¹¹ See <<http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf>>.

1 (NMVOCs), and sulfur dioxide (SO₂). These gases are not direct greenhouse gases, but indirectly affect terrestrial
 2 radiation absorption by influencing the formation and destruction of tropospheric and stratospheric ozone, or, in
 3 the case of SO₂, by affecting the absorptive characteristics of the atmosphere. Additionally, some of these gases
 4 may react with other chemical compounds in the atmosphere to form compounds that are greenhouse gases.
 5 Carbon monoxide is produced when carbon-containing fuels are combusted incompletely. Nitrogen oxides (i.e., NO
 6 and NO₂) are created by lightning, fires, fossil fuel combustion, and in the stratosphere from N₂O. Non-methane
 7 volatile organic compounds—which include hundreds of organic compounds that participate in atmospheric
 8 chemical reactions (i.e., propane, butane, xylene, toluene, ethane, and many others)—are emitted primarily from
 9 transportation, industrial processes, and non-industrial consumption of organic solvents. In the United States, SO₂
 10 is primarily emitted from coal combustion for electric power generation and the metals industry. Sulfur-containing
 11 compounds emitted into the atmosphere tend to exert a negative radiative forcing (i.e., cooling) and therefore are
 12 discussed separately.

13 One important indirect climate change effect of NMVOCs and NO_x is their role as precursors for tropospheric
 14 ozone formation. They can also alter the atmospheric lifetimes of other greenhouse gases. Another example of
 15 indirect greenhouse gas formation into greenhouse gases is the interaction of CO with the hydroxyl radical—the
 16 major atmospheric sink for CH₄ emissions—to form CO₂. Therefore, increased atmospheric concentrations of CO
 17 limit the number of hydroxyl molecules (OH) available to destroy CH₄.

18 Since 1970, the United States has published estimates of emissions of CO, NO_x, NMVOCs, and SO₂ (EPA 2020b),¹²
 19 which are regulated under the Clean Air Act. Table 2-15 shows that fuel combustion accounts for the majority of
 20 emissions of these indirect greenhouse gases. Industrial processes—such as the manufacture of chemical and
 21 allied products, metals processing, and industrial uses of solvents—are also significant sources of CO, NO_x, and
 22 NMVOCs.

23

24 **Table 2-15: Emissions of NO_x, CO, NMVOCs, and SO₂ (kt)**

Gas/Activity	1990	2005	2015	2016	2017	2018	2019
NO_x	21,739	17,339	10,187	8,792	8,642	8,145	7,754
Mobile Fossil Fuel Combustion	10,862	10,295	5,634	4,739	4,563	4,123	3,862
Stationary Fossil Fuel Combustion	10,023	5,858	3,084	2,856	2,728	2,711	2,581
Oil and Gas Activities	139	321	622	594	565	565	565
Industrial Processes and Product Use	592	572	408	402	397	397	397
Forest Fires	22	126	312	87	281	242	242
Waste Combustion	82	128	88	80	71	71	71
Grassland Fires	5	21	21	19	21	20	20
Agricultural Burning	13	15	14	14	14	14	14
Waste	+	2	2	1	1	1	1
CO	130,969	71,781	51,525	39,287	45,314	42,355	41,524
Mobile Fossil Fuel Combustion	119,360	58,615	32,635	28,789	28,124	26,590	25,749
Forest Fires	800	4,511	11,136	3,080	10,036	8,626	8,626
Stationary Fossil Fuel Combustion	5,000	4,648	3,688	3,690	3,692	3,692	3,692
Waste Combustion	978	1,403	1,576	1,375	1,175	1,175	1,175
Industrial Processes and Product Use	4,129	1,557	1,163	1,075	1,006	1,006	1,006
Oil and Gas Activities	302	318	622	607	592	592	592
Grassland Fires	84	358	356	324	345	331	341
Agricultural Burning	315	363	342	340	339	338	337
Waste	1	7	7	6	5	5	5
NMVOCs	20,930	13,154	10,596	9,774	9,444	9,228	9,123
Industrial Processes and Product Use	7,638	5,849	3,796	3,776	3,767	3,767	3,767
Mobile Fossil Fuel Combustion	10,932	5,724	3,458	2,873	2,758	2,543	2,437

¹² NO_x and CO emission estimates from Field Burning of Agricultural Residues were estimated separately, and therefore not taken from EPA (2019b).

Oil and Gas Activities	554	510	2,656	2,459	2,262	2,262	2,262
Stationary Fossil Fuel Combustion	912	716	493	489	496	496	496
Waste Combustion	222	241	132	121	109	109	109
Waste	673	114	63	57	52	52	52
Agricultural Burning	NA	NA	NA	NA	NA	NA	NA
SO₂	20,935	13,196	3,578	2,906	2,313	2,233	1,966
Stationary Fossil Fuel Combustion	18,407	11,541	2,901	2,269	1,638	1,569	1,304
Industrial Processes and Product Use	1,307	831	482	466	509	509	509
Oil and Gas Activities	390	180	92	89	86	86	86
Mobile Fossil Fuel Combustion	793	619	78	57	58	47	45
Waste Combustion	38	25	26	24	22	22	22
Waste	+	1	1	1	1	1	1
Agricultural Burning	NA	NA	NA	NA	NA	NA	NA

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.5 kt.

NA (Not Available)

Source: (EPA 2020b) except for estimates from Forest Fires, Grassland Fires, and Field Burning of Agricultural Residues.

1

Box 2-3: Sources and Effects of Sulfur Dioxide

Sulfur dioxide (SO₂) emitted into the atmosphere through natural and anthropogenic processes affects the earth's radiative budget through its photochemical transformation into sulfate aerosols that can:

- (1) scatter radiation from the sun back to space, thereby reducing the radiation reaching the earth's surface;
- (2) affect cloud formation; and
- (3) affect atmospheric chemical composition (e.g., by providing surfaces for heterogeneous chemical reactions).

The indirect effect of sulfur-derived aerosols on radiative forcing can be considered in two parts. The first indirect effect is the aerosols' tendency to decrease water droplet size and increase water droplet concentration in the atmosphere. The second indirect effect is the tendency of the reduction in cloud droplet size to affect precipitation by increasing cloud lifetime and thickness. Although still highly uncertain, the radiative forcing estimates from both the first and the second indirect effect are believed to be negative, as is the combined radiative forcing of the two (IPCC 2013).

Sulfur dioxide is also a major contributor to the formation of regional haze, which can cause significant increases in acute and chronic respiratory diseases. Once SO₂ is emitted, it is chemically transformed in the atmosphere and returns to the earth as the primary source of acid rain. Because of these harmful effects, the United States has regulated SO₂ emissions in the Clean Air Act.

Electric power is the largest anthropogenic source of SO₂ emissions in the United States, accounting for 46.9 percent in 2019. Coal combustion contributes nearly all of those emissions (approximately 92 percent). Sulfur dioxide emissions have decreased in recent years, primarily as a result of electric power generators switching from high-sulfur to low-sulfur coal and installing flue gas desulfurization equipment.

2