

Methods for Childhood Cancer

Indicator H4 Cancer incidence and mortality for children ages 0 to 19 years, 1992-2018.

See below for Indicator H5 Cancer incidence for children ages 0 to 19 years, by type, 1993-2018.

Summary

Since 1973, the Surveillance, Epidemiology, and End Results (SEER) Program of the National Cancer Institute (NCI) has been collecting and publishing cancer incidence data from population-based cancer registries that currently cover a total of 18 geographical areas in the United States and one quarter of the population. Since the coverage area has expanded over time, the trend analyses for indicator H4 were based on the SEER 13 registries that cover 13.4% of the U.S. population based on 2010 Census data. Data include the type of cancer, age at diagnosis, year of diagnosis, sex, race, and ethnicity. For indicator H4, cancer incidence rates for malignant cancers in children ages 0 to 19 years were calculated using the SEER*Stat software provided by the NCI. SEER*Stat calculates incidence rates for each age group by dividing the number of new cancer cases in a given year by the total population for that age group in the cancer registry geographical area in that year. SEER*Stat then calculates age-adjusted incidence rates as a weighted average of the rates for each five-year age group, where the weights are the proportions of persons in each age group in the selected standard population; for these analyses the incidence rates were age-adjusted to the 2000 U.S. Standard Population.

National cancer mortality rates were also obtained from SEER, which uses data from the National Vital Statistics System, administered by the National Center for Health Statistics. Data include the type of cancer, age at death, year of death, sex, race, and ethnicity. For indicator H4, cancer mortality rates for malignant cancers in children ages 0 to 19 years were also calculated using SEER*Stat. The mortality rate is calculated by dividing the number of deaths attributed to a malignant cancer in a given year by the total population for that age group in the cancer registry geographical area in that year. The mortality rates were age-adjusted to the 2000 U.S. Standard Population.

Supplementary Tables: Tables H4a, H4b, and H4c provide the age-adjusted cancer incidence and mortality rates for children 0 to 19 years, stratified by race/ethnicity, sex, and age for 2016–2018.

Health: Childhood Cancer

Data Summary

Indicator H4. Cancer incidence and mortality for children ages 0 to 19 years.

Data	Cancer incidence of malignant cancers in SEER 13 registries for children ages 0 to 19 years. U.S. mortality from malignant cancers in children ages 0 to 19 years.						
Years	1992	1993	1994	1995	1996	1997	1998
Cancer incidence	1,654	1,723	1,654	1,684	1,775	1,714	1,833
Cancer mortality	2,417	2,419	2,354	2,275	2,224	2,261	2,176
Years	1999	2000	2001	2002	2003	2004	2005
Cancer incidence	1,779	1,838	1,912	1,982	1,807	1,934	2,029
Cancer mortality	2,243	2,271	2,226	2,271	2,233	2,223	2,183
Years	2006	2007	2008	2009	2010	2011	2012
Cancer incidence	1,837	2,007	2,036	2,103	2,116	2,093	2,061
Cancer mortality	2,035	2,068	2,039	1,964	1,925	1,922	1,995
Years	2013	2014	2015	2016	2017	2018	
Cancer incidence	2,043	2,159	2,229	2,143	2,054	2,047	
Cancer mortality	1,914	1,837	1,855	1,920	1,797	1,841	

Calculation of Indicator

All the calculations for this indicator were carried out using SEER*Stat software, version 8.3.9.2. In this section we detail the menu options required for carrying out these analyses.

Note that any User-Defined variable previously created using the File and Dictionary menus in the same or any earlier session will not need to be recreated.

Incidence

Click the Rate Session button “Σ”.

Data tab. Select “Incidence – SEER Research Data, 13 Registries, Nov 2020 Sub (1992–2018).” For “Age Variable” select “Age recode with <1 year olds.”

Statistic tab. Check “Rates (Age – Adjusted)” and “Show Standard Errors and Confidence Intervals.” For “Standard Population” select “2000 US Std Population (19 age groups – Census P25-1130).” For “Age Variable” select “Age recode with <1 year olds.”

Selection tab. Check “Malignant Behavior.” For “Age at Diagnosis” click the Edit button and then select the age group 0–19 using “Age recode with <1 year olds” = 00 years, 01-04 years, 05-09 years, 10-14 years, and 15-19 years. This restricts the data to malignant cancers in children ages 0 to 19 years.

Table tab. From the “Available Variables” panel, expand “Race, Sex, Year Dx,” and select “Year of diagnosis.” Click the Row button. This tabulates the rates by calendar year.

Output tab. For “Display Rates as Cases Per” select “1,000,000.” For “Number of Decimal Places for” select “0.001.”

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the data to a text file.

Mortality

Click the Rate Session button “Σ”.

Data tab. Select “Mortality - All COD, Aggregated With State, Total U.S. (1969–2018) <Katrina/Rita Population Adjustment>.” For “Age Variable” select “Age recode with <1 year olds.”

Statistic tab. Check “Rates (Age – Adjusted)” and “Show Standard Errors and Confidence Intervals.” For “Standard Population” select “2000 US Std Population (19 age groups – Census P25-1130).” For “Age Variable” select “Age recode with <1 year olds.”

Health: Childhood Cancer

Selection tab. For “Age at Death” click the Edit button and then select the age group 0–19 using “Age recode with <1 year olds” = 00 years, 01-04 years, 05-09 years, 10-14 years, and 15-19 years. For “Other (Case Files)” click the Edit button and then follow the menus to select “Site and Morphology,” “Cause of death recode,” and “All Malignant Cancers.” This restricts the data to malignant cancers in children ages 0 to 19 years.

Table tab. From the “Available Variables” panel, expand “Race, Sex, Year Dth, State, Registry” and select “Year of Death.” Click the Row button. This tabulates the rates by calendar year.

Output tab. For “Display Rates as Cases Per” select “1,000,000.” For “Number of Decimal Places for” select “0.001.”

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the data to a text file.

Age adjustment

For indicator H4, all cancer incidence and mortality rates were age-adjusted to the 2000 U.S. Standard Population using 5 age groups for children. The calculations were carried out automatically by the SEER*Stat software. The age-adjusted rates are often preferred when comparing different populations (such as for different calendar years or different race groups) since differences in the age-adjusted cancer rates are mainly attributable to factors other than age. In general, very young children and adolescents will tend to have a higher crude (unadjusted) cancer rate than children ages 5 to 14. The age adjustment replaces the distribution of ages in each given population by the same standard age distribution, in this case the age distribution of the U.S. population in 2000.

The calculation is shown by the following example which gives the age-adjusted cancer incidence rate for 2018 for children ages 0 to 19 years.

Age-adjusted Cancer Incidence for 2018.

<i>Age Group [1]</i>	<i>Cases [2]</i>	<i>SEER 13 Population [3]</i>	<i>Crude Rate (Cases per Million)[4] = [2]/[3] × 1,000,000</i>	<i>U.S. 2000 Standard Population [5]</i>	<i>U.S. 2000 Proportion of Age Group 0-19 [6]</i>	<i>Adjusted Rate (Cases per Million) = [4] × [6]</i>
00 years	126	513,509	245.371	3,794,901	0.048	11.819
01-04 years	468	2,148,693	217.807	15,191,619	0.193	42.000
05-09 years	358	2,718,074	131.711	19,919,840	0.253	33.303

Health: Childhood Cancer

<i>Age Group [1]</i>	<i>Cases [2]</i>	<i>SEER 13 Population [3]</i>	<i>Crude Rate (Cases per Million)[4] = [2]/[3] × 1,000,000</i>	<i>U.S. 2000 Standard Population [5]</i>	<i>U.S. 2000 Proportion of Age Group 0-19 [6]</i>	<i>Adjusted Rate (Cases per Million) = [4] × [6]</i>
10-14 years	446	2,789,246	159.900	20,056,779	0.255	40.708
15-19 years	649	2,775,691	233.816	19,819,518	0.252	58.821
00-19 years	2,047	10,945,213	187.022	78,782,657	1.000	186.651

The first three columns give the number of new malignant cancer cases in 2018 and the total population for each children’s age group within the counties included in the SEER 13 cancer registries. The number of cases divided by the SEER 13 population and multiplied by 1 million gives the crude rate shown in the fourth column. For example, for the age group 00 years, there were 126 cases in a population of 513,509 giving a crude rate of $(126/513509) \times 1000000 = 245.371$ cases per million. The crude rate for all children ages 0 to 19 years is 187.022 (bottom row).

The U.S. 2000 Standard Population for children is shown in the fifth column. This is the age distribution of children for the entire United States in 2000, in five age groups (0, 1–4, 5–9, 10–14, and 15–19). For analyzing children’s populations only five age groups are needed (0, 1–4, 5–9, 10–14, and 15–19). However, for analyzing children and adults the U.S. 2000 Standard Population has 19 age groups, since the U.S. 2000 Standard Population also includes five-year age groups up to 80–84 and the 85+ age group. The standard age distribution for children ages 0 to 19 years is shown in the sixth column. The U.S. 2000 Standard Population in each age group is divided by the total U.S. 2000 Standard Population for children ages 0 to 19 years, i.e., 78,782,657.

The age-adjusted population is a weighted average of the crude rates (column 4) weighted by the U.S. 2000 Standard Population proportions (column 6). Thus, the first five rows of column 7 contain the products of columns 4 and 6, and the age-adjusted rate is given by the total for column 7, shown in the bottom row as 186.651.

For this example, the crude rate for children ages 0 to 19 years, 187.022, is very close to the age-adjusted rate, 186.651, reflecting the fact that in 2018 the age distribution for children 0 to 19 years in the SEER 13 regions closely matches the 2000 U.S. population age distribution. For other years or for different race/ethnicity/sex groups, crude and age-adjusted rates can differ by a much greater percentage.

Relative Standard Error

The uncertainties of the incidence and mortality rates were computed by the SEER*Stat software under the assumption that the counts have Poisson distributions. The relative standard error is the standard error of the rate divided by the estimated rate:

Health: Childhood Cancer

$$\text{Relative Error (\%)} = [\text{Standard Error (Rate)} / \text{Rate}] \times 100\%$$

Rates with a relative error less than 30% were treated as being reliable and were tabulated. For indicator H4, as well as the tables H4a, H4b, and H4c, described below, all the relative standard errors were at most 30% and so all values were tabulated.

Rates by Race, Ethnicity, Sex, and Age

Supplementary Tables: In Tables H4a, H4b, and H4c, cancer incidence and mortality rates are tabulated by race, ethnicity, sex, and age.

The following SEER*Stat data files were used for the analyses in Tables H4a, H4b, and H4c:

- Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Incidence - SEER Research Data, 13 Registries, Nov 2020 Sub (1992-2018) - Linked To County Attributes - Time Dependent (1990-2018) Income/Rurality, 1969-2019 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released April 2021, based on the November 2020 submission. Incidence data from the SEER 13 registries: Connecticut; Hawaii; Iowa; New Mexico; Utah; Atlanta, Georgia; Detroit, Michigan; San Francisco-Oakland, California; Seattle-Puget Sound, Washington; Los Angeles, California; San Jose-Monterey, California; Rural Georgia; and the Alaska Native Tumor Registry. Downloaded November 2021.
- Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Mortality - All COD, Aggregated With State, Total U.S. (1990-2018) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, released May 2020. Underlying mortality data provided by NCHS (www.cdc.gov/nchs). Mortality from All Malignant Cancers was selected. Downloaded November 2021.

The supplementary tables by race, ethnicity, sex, and age for incidence and mortality in 2016-2018 were calculated as follows:

Incidence by Race/Ethnicity and Sex

Click the Rate Session button “Σ”.

Data tab. Select “Incidence – SEER Research Data, 13 Registries, Nov 2020 Sub (1992–2018).” For “Age Variable” select “Age recode with <1 year olds.”

Statistic tab. Check “Rates (Age – Adjusted)” and “Show Standard Errors and Confidence Intervals.” For “Standard Population” select “2000 US Std Population (19 age groups – Census P25-1130).” For “Age Variable” select “Age recode with <1 year olds.”

Health: Childhood Cancer

Selection tab. Check “Malignant Behavior.” For “Age at Diagnosis” click the Edit button and then select the age group 0–19 using “Age recode with <1 year olds” = 00 years, 01-04 years, 05-09 years, 10-14 years, and 15-19 years. For “Race, Sex, Year Dx, Registry, County” click the Edit button and select the years 2016-2018 using “Year of Diagnosis” = 2016, 2017, and 2018. This restricts the data to malignant cancers in children ages 0 to 19 years diagnosed in 2016-2018.

Table tab. From the “Available Variables” panel, expand “Race, Sex, Year Dx” and select “Sex.” Click the Row button. Use the menus “File,” “Dictionary,” and “Merge” to create a User-Defined Merged variable Raceeth from the Race recode (W, B, AI, API) and Origin recode NHIA (Hispanic, Non-Hisp) variables in the database:

- Raceeth = All if Race recode = White, Black, American Indian/Alaskan Native, Asian or Pacific Islander, or Unknown
- Raceeth = WhiteNH if Race recode = White and Origin recode NHIA = Non-Spanish-Hispanic-Latino
- Raceeth = BlackNH if Race recode = Black and Origin recode NHIA = Non-Spanish-Hispanic-Latino
- Raceeth = AIANNH if Race recode = American Indian/Alaskan Native and Origin recode NHIA = Non-Spanish-Hispanic-Latino
- Raceeth = APINH if Race recode = Asian or Pacific Islander and Origin recode NHIA = Non-Spanish-Hispanic-Latino
- Raceeth = Hispanic if Origin recode NHIA = Spanish-Hispanic-Latino

In the Table tab, from the “Available Variables” panel, expand “Merged” and select “Raceeth.” Click the Column button. These options tabulate the rates by sex and race/ethnicity.

Output tab. For “Display Rates as Cases Per” select “1,000,000.” For “Number of Decimal Places for” select “0.001.”

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the race/ethnicity/sex data to a text file.

Incidence by Age

Assuming you are in the same session as above, modify the selections as follows:

Table tab. In the Table tab, from the “Display Variables” panel, select the Row variable “Sex” and click Remove. Also select the Column variable “Raceeth” and click Remove. Use the menus “File” and “Dictionary” to create a User-Defined variable AgesH4 that takes the values 0-4 years, 5-9 years, 10-14 years, 15-19 years, and 0-19 years if the “Age at Diagnosis” is in the selected age group. Age at Diagnosis is defined by the values of “Age recode with <1 year olds.” In the Table tab, from the “Available Variables” panel, expand “User-Defined” and select “AgesH4.” Click the Row button. This tabulates the rates by the selected age groups.

Health: Childhood Cancer

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the data to a text file. Tabulate the rates for selected age groups.

Mortality by Race/Ethnicity and Sex

Click the Rate Session button “Σ”.

Data tab. Select “Mortality - All COD, Aggregated With State, Total U.S. (1990–2018) <Katrina/Rita Population Adjustment>.” For “Age Variable” select “Age recode with <1 year olds.”

Statistic tab. Check “Rates (Age – Adjusted)” and “Show Standard Errors and Confidence Intervals.” For “Standard Population” select “2000 US Std Population (19 age groups – Census P25-1130).” For “Age Variable” select “Age recode with <1 year olds.”

Selection tab. For “Age at Death” click the Edit button and then select the age group 0 to 19 years using “Age recode with <1 year olds” = 00 years, 01-04 years, 05-09 years, 10-14 years, and 15-19 years. For “Race, Sex, Year Dth, State, Registry” click the Edit button and select the years 2016-2018 using “Year of Death” = 2016, 2017 and 2018. For “Other (Case Files)” click the Edit button and then follow the menus to select “Site and Morphology,” “Cause of death recode,” and “All Malignant Cancers.” These options restrict the data to malignant cancers in children ages 0 to 19 years who died in 2016-2018.

Table tab. From the “Available Variables” panel, expand “Race, Sex, Year Dth, State, Registry” and select “Sex.” Click the Row button. Use the menus “File,” “Dictionary,” and “Merge” to create a User-Defined Merged variable Raceeth2 from the Race recode (W, B, AI, API) and Origin recode 1990+ (Hispanic, Non-Hisp variables in the database:

- Raceeth2 = All if Race recode = White, Black, American Indian/Alaskan Native, or Asian or Pacific Islander
- Raceeth2 = WhiteNH if Race recode = White, Origin recode 1990+ (Hispanic, Non-Hisp) = Non-Spanish-Hispanic-Latino or Unknown
- Raceeth2 = BlackNH if Race recode = Black, Origin recode 1990+ (Hispanic, Non-Hisp) = Non-Spanish-Hispanic-Latino or Unknown
- Raceeth2 = AIANNH if Race recode = American Indian/Alaskan Native, Origin recode 1990+ (Hispanic, Non-Hisp) = Non-Spanish-Hispanic-Latino or Unknown
- Raceeth2 = APINH if Race recode = Asian or Pacific Islander, Origin recode 1990+ (Hispanic, Non-Hisp) = Non-Spanish-Hispanic-Latino or Unknown
- Raceeth2 = Hispanic if Origin recode 1990+ (Hispanic, Non-Hisp) = Spanish-Hispanic-Latino

Health: Childhood Cancer

In the Table tab, from the “Available Variables” panel, expand “Merged” and select “Raceeth2.” Click the Column button. These options tabulate the rates by sex and race/ethnicity.

Output tab. For “Display Rates as Cases Per” select “1,000,000.” For “Number of Decimal Places for” select “0.001.”

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the race/ethnicity/sex data to a text file.

Questions and Comments

Questions regarding these methods, and suggestions to improve the description of the methods, are welcome. Please use the “Contact Us” link at the bottom of any page in the America’s Children and the Environment website.

Statistical Comparisons

Statistical analyses of the cancer incidence and mortality rates were used to determine whether the differences between rates for different demographic groups were statistically significant. For these analyses, the incidence and mortality counts and populations for each combination of calendar year, age group, sex, and race/ethnicity were obtained from the SEER database using the same race/ethnicity groups described above (excluding the All category)ⁱ, and the age groups 0-4 years, 5-9 years, 10-14 years, and 15-19 years. Incidence data for malignant cancers for the years 1992-2018 were obtained from the SEER data set “Incidence – SEER Research Data, 13 Registries, Nov 2020 Sub (1992–2018).” U.S. mortality data for 1992-2018 were obtained from the SEER data set “Mortality - All COD, Aggregated With State, Total U.S. (1990–2018) <Katrina/Rita Population Adjustment>.” For the U.S. mortality data, the analysis included data from all U.S. States.ⁱⁱ For the mortality data, SEER*Stat version 8.3.9.2 does not report mortality counts of 9 or fewer deaths. For the statistical analysis, any missing mortality count for a combination of calendar year, age group, sex, and race/ethnicity was replaced by 5.

Using a Poisson regression model adjusted for dispersion, the number of cases for each demographic subgroup was assumed to have a Poisson distribution with a mean equal to the

ⁱ For the incidence data used for these analyses, the race recode variable includes an “Unknown” category. For that category, the SEER database provides case counts, but the reported populations for that category are zero and the corresponding age-adjusted incidence rates could not be calculated. For this reason, the statistical analysis excludes data from persons of unknown race. For the mortality data used for these analyses, the race recode variable includes an “Other unspecified (1978-1991)” category, but those persons are not included in the time period studied.

ⁱⁱ At the url [http://seer.cancer.gov/seerstat/variables/mort/origin_recode_1990+/,](http://seer.cancer.gov/seerstat/variables/mort/origin_recode_1990+/) NCI recommends excluding mortality data from states and years where Hispanic origin was not recorded. This would apply to Louisiana in 1990, New Hampshire in 1990-1992 and Oklahoma in 1990-1996. Applying this rule to individual years would introduce a bias by excluding different states in different years. Thus we decided to not exclude any mortality data for these analyses.

Health: Childhood Cancer

population of that demographic subgroup multiplied by the rate, and a variance equal to the mean multiplied by a dispersion parameter. The logarithm of the rate was assumed to be the sum of explanatory terms for age, sex, and race/ethnicity. Using this model, the difference in the value of a rate between different demographic groups is statistically significant if the difference between the corresponding sums of explanatory terms is statistically significantly different from zero. A p-value at or below 0.05 implies that the difference is statistically significant at the 5% significance level. No adjustment is made for multiple comparisons.

For each type of comparison, we present unadjusted and adjusted analyses. The unadjusted analyses compare a rate between different demographic groups. The adjusted analyses add other demographic explanatory variables to the statistical model and use the statistical model to account for the possible confounding effects of these other demographic variables. For example, the unadjusted race/ethnicity comparisons compare the rates between different race/ethnicity pairs. The adjusted analyses add age and sex terms to the statistical model and compare the rates between different race/ethnicity pairs after accounting for the effects of the other demographic variables. For example, if Hispanic children tend to be younger than White non-Hispanics, and if the rate strongly depends on age only, then the unadjusted differences between these two race/ethnicity groups would be significant but the adjusted difference (taking into account age) would not be significant.

Comparisons of incidence rates between pairs of race/ethnicity groups are shown in Table 1. Comparisons of mortality rates between pairs of race/ethnicity groups are shown in Table 2. In Tables 1 and 2, for the “Unadjusted” comparisons, the only explanatory variables are terms for each race/ethnicity group. For these unadjusted comparisons, the statistical tests compare the rates for each pair of race/ethnicity groups. For the “Adjusted for age, sex” comparisons, the explanatory variables are terms for each race/ethnicity together with terms for each age group and sex. For these adjusted comparisons, the statistical test compares the pair of race/ethnicity groups after accounting for any differences in the age and sex distributions between the race/ethnicity groups.

Additional comparisons are shown in Table 3 for incidence rates and in Table 4 for mortality rates. The Against = “age” unadjusted p-value compares the rates between all the age groups. The adjusted p-value includes adjustment terms for sex and race/ethnicity in the model. The Against = “sex” unadjusted p-value compares the rates between the two sexes. The adjusted p-value includes adjustment terms for age and race/ethnicity in the model. The Against = “year” unadjusted p-value compares the trends in the rates by regressing against the calendar year. The adjusted p-value includes adjustment terms for age, sex, and race/ethnicity in the model. To evaluate whether the difference in rates for Males and Females varies according to the race/ethnicity, p-values for comparing the Male and Female rates by race/ethnicity group are found in the rows where Against = “sex” and Subset = the race/ethnicity group. The adjusted p-value includes an adjustment term for age.

For more details on these statistical analyses, see the memorandum by Cohen (2011).ⁱⁱⁱ

ⁱⁱⁱ Cohen, J. 2011. *Selected statistical methods for testing for trends and comparing years or demographic groups in other ACE health-based indicators*. Memorandum submitted to Dan Axelrad, EPA, 16 June, 2011.

Health: Childhood Cancer

Table 1. Statistical significance tests comparing cancer incidence rates for children ages 0 to 19 years, between pairs of race/ethnicity groups, for 2016-2018.

Variable	First race/ethnicity group	Second race/ethnicity group	P-VALUES	
			Unadjusted	Adjusted for age, sex
Cancer incidence	White non-Hispanic	Black non-Hispanic	0.002	< 0.001
Cancer incidence	White non-Hispanic	API non-Hispanic	0.114	0.001
Cancer incidence	White non-Hispanic	AIAN non-Hispanic	0.561	0.241
Cancer incidence	White non-Hispanic	Hispanic	0.092	0.001
Cancer incidence	Black non-Hispanic	API non-Hispanic	0.227	0.011
Cancer incidence	Black non-Hispanic	AIAN non-Hispanic	0.571	0.219
Cancer incidence	Black non-Hispanic	Hispanic	0.076	< 0.001
Cancer incidence	API non-Hispanic	AIAN non-Hispanic	0.990	0.951
Cancer incidence	API non-Hispanic	Hispanic	0.728	0.380
Cancer incidence	AIAN non-Hispanic	Hispanic	0.902	0.780

Table 2. Statistical significance tests comparing cancer mortality rates for children ages 0 to 19 years, between pairs of race/ethnicity groups, for 2016-2018.

Variable	First race/ethnicity group	Second race/ethnicity group	P-VALUES	
			Unadjusted	Adjusted for age, sex
Cancer mortality	White non-Hispanic	Black non-Hispanic	0.001	< 0.001
Cancer mortality	White non-Hispanic	API non-Hispanic	0.509	< 0.001
Cancer mortality	White non-Hispanic	AIAN non-Hispanic	0.454	< 0.001
Cancer mortality	White non-Hispanic	Hispanic	0.761	0.041
Cancer mortality	Black non-Hispanic	API non-Hispanic	0.016	< 0.001
Cancer mortality	Black non-Hispanic	AIAN non-Hispanic	0.847	0.316
Cancer mortality	Black non-Hispanic	Hispanic	0.002	< 0.001
Cancer mortality	API non-Hispanic	AIAN non-Hispanic	0.326	< 0.001
Cancer mortality	API non-Hispanic	Hispanic	0.637	0.012
Cancer mortality	AIAN non-Hispanic	Hispanic	0.413	< 0.001

Table 3. Other statistical significance tests comparing cancer incidence rates for children ages 0 to 19 years for 2016 to 2018 (trends for 1992-2018).

Variable	From	To	Against	Subset	P-VALUES	
					Unadjusted	Adjusted*
Cancer incidence	2016	2018	age		< 0.001	< 0.001
Cancer incidence	2016	2018	race		0.036	< 0.001
Cancer incidence	2016	2018	sex		0.038	< 0.001
Cancer incidence	2016	2018	sex	White non-Hispanic	0.431	0.003
Cancer incidence	2016	2018	sex	Black non-Hispanic	0.125	0.010
Cancer incidence	2016	2018	sex	API non-Hispanic	0.779	0.550
Cancer incidence	2016	2018	sex	AIAN non-Hispanic	0.556	0.536

Health: Childhood Cancer

Variable	From	To	Against	Subset	P-VALUES	
					Unadjusted	Adjusted*
Cancer incidence	2016	2018	sex	Hispanic	0.163	< 0.001
Cancer incidence	1992	2018	year		< 0.001	< 0.001

*For Against = "age," the p-values are adjusted for sex and race/ethnicity.

For Against = "sex," the p-values are adjusted for age and race/ethnicity.

For Against = "year," the p-values are adjusted for age, sex, and race/ethnicity.

Table 4. Other statistical significance tests comparing cancer mortality rates for children ages 0 to 19 years for 2016 to 2018 (trends for 1992-2018).

Variable	From	To	Against	Subset	P-VALUES	
					Unadjusted	Adjusted*
Cancer mortality	2016	2018	age		< 0.001	< 0.001
Cancer mortality	2016	2018	race		0.004	< 0.001
Cancer mortality	2016	2018	sex		0.042	< 0.001
Cancer mortality	2016	2018	sex	White non-Hispanic	0.325	< 0.001
Cancer mortality	2016	2018	sex	Black non-Hispanic	0.351	< 0.001
Cancer mortality	2016	2018	sex	API non-Hispanic	0.441	< 0.001
Cancer mortality	2016	2018	sex	AIAN non-Hispanic	0.304	< 0.001
Cancer mortality	2016	2018	sex	Hispanic	0.390	< 0.001
Cancer mortality	1992	2018	year		< 0.001	< 0.001

*For Against = "age," the p-values are adjusted for sex and race/ethnicity.

For Against = "sex," the p-values are adjusted for age and race/ethnicity.

For Against = "year," the p-values are adjusted for age, sex, and race/ethnicity.

Data Files

The following files are needed to calculate this indicator. All these files together with the SEER*Stat software used to calculate the incidence and mortality rates are available from the SEER website: <http://seer.cancer.gov/data/access.html>. The data and analyses used for this indicator were obtained using version 8.3.9.2 of SEER*Stat.

- Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Incidence - SEER Research Data, 13 Registries, Nov 2020 Sub (1992-2018) - Linked To County Attributes - Time Dependent (1990-2018) Income/Rurality, 1969-2019 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released April 2021, based on the November 2020 submission. Incidence data from the SEER 13 registries: Connecticut; Hawaii; Iowa; New Mexico; Utah; Atlanta, Georgia; Detroit, Michigan; San Francisco-Oakland, California; Seattle-Puget Sound, Washington; Los Angeles, California; San Jose-Monterey, California; Rural Georgia; and the Alaska Native Tumor Registry. Downloaded November 2021.

Health: Childhood Cancer

- Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Mortality - All COD, Aggregated With State, Total U.S. (1990-2018) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, released May 2020. Underlying mortality data provided by NCHS (www.cdc.gov/nchs). Mortality from All Malignant Cancers was selected. Downloaded November 2021.

The databases incorporated into the SEER*Stat software include census population data used to compute the incidence and mortality rates and to age-adjust the rates to the 2000 U.S. Standard Population. These population data are obtained by NCI from the Census Bureau. Age-adjustment is explained above.

Method for H5

Indicator H5 Cancer incidence for children ages 0 to 19 years, by type, 1993-2018.

See above for H4 Cancer incidence and mortality for children ages 0 to 19 years, 1992-2018.

Summary

Since 1973, the Surveillance, Epidemiology, and End Results (SEER) Program of the National Cancer Institute (NCI) has been collecting and publishing cancer incidence data from population-based cancer registries that currently cover a total of 18 geographical areas in the United States and one quarter of the population. Since the coverage area has expanded over time, the trend analyses for indicator H5 were based on the SEER 13 registries that cover 13.4% of the U.S. population for 1995 and later. Data include the type of cancer (cancer site), age at diagnosis, and year of diagnosis.

For indicator H5, cancer incidence rates for specific malignant cancers in children ages 0 to 19 years for two-year periods from 1993-2018 were calculated using the SEER*Stat software provided by the NCI. SEER*Stat calculates incidence rates for each age group by dividing the number of new cancer cases by the total population for that age group in the cancer registry geographical area. SEER*Stat then calculates age-adjusted incidence rates as a weighted average of the rates for each five-year age group, where the weights are the proportions of persons in each age group in the selected standard population; for these analyses the incidence rates were age-adjusted to the 2000 U.S. Standard Population. Supplementary Table H5a provides the age-adjusted cancer incidence rates for children ages 0 to 19 years, stratified by age group and cancer type, for 2017-2018.

Health: Childhood Cancer

Data Summary

Data used to support Indicators:

H5. Cancer incidence for children ages 0 to 19 years, by type, 1993-2018.

H4. Cancer incidence and mortality for children ages 0 to 19 years, 1992-2018.

Data	Cancer incidence of malignant cancers in SEER 13 registries for children ages 0 to 19 years.						
Years	1993	1994	1995	1996	1997	1998	1999
Cancer incidence	1,723	1,654	1,684	1,775	1,714	1,833	1,779
Years	2000	2001	2002	2003	2004	2005	2006
Cancer incidence	1,838	1,912	1,982	1,807	1,934	2,029	1,837
Years	2007	2008	2009	2010	2011	2012	2013
Cancer incidence	2,007	2,036	2,103	2,116	2,093	2,061	2,043
Years	2014	2015	2016	2017	2018		
Cancer incidence	2,159	2,229	2,143	2,054	2,047		

Calculation of Indicator

All the calculations for this indicator were carried out using SEER*Stat software, version 8.3.9.2. In this section we detail the menu options required for carrying out these analyses.

Note that any User-Defined variable previously created using the File and Dictionary menus in the same or any earlier session will not need to be recreated.

ICCC site recode ICD-O-3/WHO 2008.

Click the Rate Session button “Σ”.

Data tab. Select “Incidence – SEER Research Data, 13 Registries, Nov 2020 Sub (1992–2018).”

Statistic tab. Check “Rates (Age – Adjusted)” and “Show Standard Errors and Confidence Intervals.” For “Standard Population” select “2000 US Std Population (19 age groups – Census P25-1130).” For “Age Variable” select “Age recode with <1 year olds.”

Selection tab. Check “Malignant Behavior.” For “Age at Diagnosis” click the Edit button and then select the age group 0 to 19 years using “Age recode with <1 year olds” = 00 years, 01-04 years, 05-09 years, 10-14 years, and 15-19 years. This restricts the data to malignant cancers in children ages 0 to 19 years.

Table tab. Use the menus “File” and “Dictionary” to create a User-Defined variable YearsH5 that takes the values 1993-1994, 1995-1996, 1997-1998, 1999-2000, 2001-2002, 2003-2004, 2005-2006, 2007-2008, 2009-2010, 2011-2012, 2013-2014, 2015-2016, and 2017-2018 if the “Year of Diagnosis” is in the selected year group. In the Table tab, from the “Available Variables” panel, expand “User-Defined” and select “YearsH5.” Click the Row button. From the “Available Variables” panel, expand “Site and Morphology” and select “ICCC site recode ICD-O-3/WHO 2008.” Click the Column button. This tabulates the rates by the selected calendar year groups and cancer types.

Output tab. For “Display Rates as Cases Per” select “1,000,000.” For “Number of Decimal Places for” select “0.001.”

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the data to a text file. Tabulate the rates for selected cancer types and year groups.

AYA Site Recode/WHO 2008.

The cancer types acute lymphocytic leukemia and Wilms’ tumor are not specifically included in the ICCC site recode ICD-O-3/WHO 2008 categories. Thus the incidence rate for these two

Health: Childhood Cancer

cancers is calculated separately using the “AYA site recode/WHO 2008,” which codes cancer types for adolescents and young adults.

Modify the above selections as follows:

Table tab. In the Table tab, from the “Display Variables” panel, select the Column variable “ICCC site recode ICD-O-3/WHO 2008” and click Remove. From the “Available Variables” panel, expand “Site and Morphology” and select “AYA site recode/WHO 2008.” Click the Column button. This tabulates the rates by the selected calendar year groups for the cancer types coded in the AYA site recode.

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the AYA site recode data to a text file. Tabulate the rates for selected cancer types and year groups.

Age-adjustment

For indicator H5, all cancer incidence rates were age-adjusted to the 2000 U.S. Standard Population using five age groups for children. The calculations were carried out automatically by the SEER*Stat software. The age-adjusted rates are often preferred when comparing different populations (such as for different calendar years or different race groups) since differences in the age-adjusted cancer rates are mainly attributable to factors other than age. In general, a younger population will tend to have a lower crude (unadjusted) cancer rate. The age-adjustment replaces the distribution of ages in each given population by the same standard age distribution, in this case the age distribution of the U.S. population in 2000.

The calculation is shown by the following example which gives the age-adjusted cancer incidence rate for 2017-2018 for acute myeloid leukemia in children ages 0 to 19 years.

Age-adjusted Acute Myeloid Leukemia Incidence for 2017-2018.

<i>Age Group [1]</i>	<i>Cases [2]</i>	<i>SEER 13 Population [3]</i>	<i>Crude Rate (Cases per Million)[4] = [2]/[3]</i>	<i>U.S. 2000 Standard Population [5]</i>	<i>U.S. 2000 Proportion of Age Group 0-19 [6]</i>	<i>Adjusted Rate (Cases per Million) = [4] × [6]</i>
00 years	23	1,044,300	22.024	3,794,901	0.048	1.061
01-04 years	61	4,319,801	14.121	15,191,619	0.193	2.723
05-09 years	29	5,464,387	5.307	19,919,840	0.253	1.342
10-14 years	37	5,573,113	6.639	20,056,779	0.255	1.690

Health: Childhood Cancer

<i>Age Group [1]</i>	<i>Cases [2]</i>	<i>SEER 13 Population [3]</i>	<i>Crude Rate (Cases per Million)[4] = [2]/[3]</i>	<i>U.S. 2000 Standard Population [5]</i>	<i>U.S. 2000 Proportion of Age Group 0-19 [6]</i>	<i>Adjusted Rate (Cases per Million) = [4] × [6]</i>
15-19 years	56	5,562,810	10.067	19,819,518	0.252	2.533
00-19 years	206	21,964,411	9.379	78,782,657	1.000	9.348

The first three columns give the number of new malignant acute myeloid leukemia cases in 2017-2018 and the total population for each children’s age group within the counties included in the SEER 13 cancer registries. The number of cases divided by the SEER 13 population and multiplied by 1 million gives the crude rate shown in the fourth column. For example, for the age group 00 years, there were 23 cases in a population of 1,044,300 giving a crude rate of $(23/1044300) \times 1000000 = 22.024$ cases per million. The crude rate for all children ages 0 to 19 years is 9.379 (bottom row).

The U.S. 2000 Standard Population for children is shown in the fifth column. This is the age distribution for the entire United States in 2000. For analyzing children’s populations only five age groups are needed (0, 1–4, 5–9, 10–14, and 15–19). However, for analyzing children and adults the U.S. 2000 Standard Population has 19 age groups, since the U.S. 2000 Standard Population also includes five-year age groups up to 80–84 and the 85+ age group. The standard age distribution for children 0 to 19 years is shown in the sixth column. The U.S. 2000 Standard Population in each age group is divided by the total U.S. 2000 Standard Population for children ages 0 to 19 years, i.e., 78,782,657.

The age-adjusted population is a weighted average of the crude rates (column 4) weighted by the U.S. 2000 Standard Population proportions (column 6). Thus, the first five rows of column 7 contain the products of columns 4 and 6, and the age-adjusted rate is given by the total for column 7, shown in the bottom row as 9.348.

For this example, the crude rate for ages 0 to 19 years, 9.379, is very close to the age-adjusted rate, 9.348, reflecting the fact that the age distribution for children 0 to 19 years in the SEER 13 regions closely matches the 2000 U.S. population age distribution. For other years or for different cancer types, crude and age-adjusted rates can differ by a much greater percentage.

Relative Standard Error

The uncertainties of the incidence and mortality rates were computed by the SEER*Stat software under the assumption that the counts have Poisson distributions. The relative standard error is the standard error of the rate divided by the estimated rate:

$$\text{Relative Error (\%)} = [\text{Standard Error (Rate)} / \text{Rate}] \times 100\%$$

Health: Childhood Cancer

Rates with a relative error less than 30% were treated as being reliable and were tabulated. For indicator H5, all the relative standard errors in Table H5 were at most 30% and so all values were tabulated. For Table H5a (see below), some relative standard errors were greater than 30% and less than 40% and those rates were tabulated but flagged as to be interpreted with caution. Rates with a relative standard error greater than or equal to 40%, or without an estimated relative standard error, were treated as being unreliable; these values were not tabulated and were flagged as having a large uncertainty.

Rates by Age Group

For Table H5a, cancer incidence for 2017-2018 is tabulated by cancer type and age group.

The Table H5a was calculated as follows using SEER*Stat:

ICCC site recode ICD-O-3/WHO 2008.

Click the Rate Session button “Σ”.

Data tab. Select “Incidence – SEER Research Data, 13 Registries, Nov 2020 Sub (1992–2018).”

Statistic tab. Check “Rates (Age – Adjusted)” and “Show Standard Errors and Confidence Intervals.” For “Standard Population” select “2000 US Std Population (19 age groups – Census P25-1130).” For “Age Variable” select “Age recode with <1 year olds.”

Selection tab. Check “Malignant Behavior.” For “Age at Diagnosis” click the Edit button and then select the age group 0 to 19 years using “Age recode with <1 year olds” = 00 years, 01-04 years, 05-09 years, 10-14 years, and 15-19 years. For “Race, Sex, Year Dx” click the Edit button and select the years 2017-2018 using “Year of Diagnosis” = 2017 and 2018. This restricts the data to malignant cancers in children ages 0 to 19 years diagnosed in 2017-2018.

Table tab. Use the menus “File” and “Dictionary” to create a User-Defined variable AgesH5 that takes the values 0-4 years, 5-9 years, 10-14 years, 15-19 years, and 0-19 years if the “Age at Diagnosis” is in the selected age group. Age at Diagnosis is defined by the values of “Age recode with <1 year olds.” In the Table tab, from the “Available Variables” panel, expand “User-Defined” and select “AgesH5.” Click the Row button. From the “Available Variables” panel, expand “Site and Morphology” and select “ICCC site recode ICD-O-3/WHO 2008.” Click the Column button. This tabulates the rates by the selected age groups and cancer types.

Output tab. For “Display Rates as Cases Per” select “1,000,000.” For “Number of Decimal Places for” select “0.001.”

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the data to a text file. Tabulate the rates for selected cancer types and age groups.

Health: Childhood Cancer

AYA Site Recode/WHO 2008.

The cancer types acute lymphocytic leukemia and Wilms' tumor are not specifically included in the ICCC site recode ICD-O-3/WHO 2008 categories. Thus the incidence rate for these two cancers is calculated separately using the "AYA site recode/WHO 2008," which codes cancer types for adolescents and young adults.

Modify the above selections as follows:

Table tab. In the Table tab, from the "Display Variables" panel, select the Column variable "ICCC site recode ICD-O-3/WHO 2008" and click Remove. From the "Available Variables" panel, expand "Site and Morphology" and select "AYA site recode/WHO 2008." Click the Column button. This tabulates the rates by the selected age groups for the cancer types coded in the AYA site recode/WHO 2008.

Click the Execute button to run the analyses and create the output Rate Matrix.

Use the Matrix and Export menus to output the AYA site recode data to a text file. Tabulate the rates for selected cancer types and age groups.

Questions and Comments

Questions regarding these methods, and suggestions to improve the description of the methods, are welcome. Please use the "Contact Us" link at the bottom of any page in the America's Children and the Environment website.

Statistical Comparisons

Statistical analyses of the cancer incidence rates for each cancer type were used to determine whether the differences between rates for different demographic groups were statistically significant. Each cancer type was analyzed separately. Incidence data for malignant cancers in 1993-2018 were obtained from the SEER data set "SEER Research Data, 13 Registries, Nov 2020 Sub (1992–2018)." For these analyses, the incidence counts and populations for each combination of cancer type, calendar year, age group, sex, and race/ethnicity were obtained from the SEER database using the age groups 0-4 years, 5-9 years, 10-14 years, and 15-19 years and the following race/ethnicity groups defined by the Race recode and Origin recode NHIA variables in the SEER database:

- Raceeth = WhiteNH if Race recode = White and Origin recode NHIA = Non-Spanish-Hispanic-Latino
- Raceeth = BlackNH if Race recode = Black and Origin recode NHIA = Non-Spanish-Hispanic-Latino
- Raceeth = AIANNH if Race recode = American Indian/Alaskan Native and Origin recode NHIA = Non-Spanish-Hispanic-Latino

Health: Childhood Cancer

- Raceeth = APINH if Race recode = Asian or Pacific Islander and Origin recode NHIA = Non-Spanish-Hispanic-Latino
- Raceeth = Hispanic if Origin recode NHIA = Spanish-Hispanic-Latino

Using a Poisson regression model adjusted for dispersion, the number of cases for each demographic subgroup was assumed to have a Poisson distribution with a mean equal to the population of that demographic subgroup multiplied by the rate, and a variance equal to the mean multiplied by a dispersion parameter. The logarithm of the rate was assumed to be the sum of explanatory terms for age, sex, and race/ethnicity^{iv}. Using this model, the difference in the value of a rate between different demographic groups is statistically significant if the difference between the corresponding sums of explanatory terms is statistically significantly different from zero. A p-value at or below 0.05 implies that the difference is statistically significant at the 5% significance level. No adjustment is made for multiple comparisons.

For each type of comparison, we present unadjusted and adjusted analyses. The unadjusted analyses compare a rate between different demographic groups. The adjusted analyses add other demographic explanatory variables to the statistical model and use the statistical model to account for the possible confounding effects of these other demographic variables. For example, the unadjusted age comparisons compare the rates between the different age groups. The adjusted analyses add sex and race/ethnicity terms to the statistical model and compare the rates between different age groups after accounting for the effects of the other demographic variables. For example, if Hispanic children tend to be younger than White non-Hispanics, and if the cancer incidence rate strongly depends on race/ethnicity only, then the unadjusted differences between age groups would be significant because of their different race/ethnicity breakdowns but the adjusted difference (taking into account race/ethnicity) would not be significant.

The statistical comparisons for each cancer type are shown in Table 1. The Against = “age” unadjusted p-value compares the rates between all the age groups for 2017-2018. The adjusted p-value includes adjustment terms for sex and race/ethnicity in the model. The Against = “year” unadjusted p-value compares the trends in the rates for 1993 -2018 by regressing against the calendar year. Instead of using the two-year rates given in the tables and graphs, the statistical trend analysis used the annual rates to improve the precision of the trend estimates and p-values. The adjusted p-value includes adjustment terms for age, sex, and race/ethnicity in the model.

For more details on these statistical analyses, see the memorandum by Cohen (2011).^v

^{iv} For the incidence data used for these analyses, the race recode variable includes an “Unknown” category. For that category, the SEER database provides case counts, but the reported populations for that category are zero and the corresponding age-adjusted incidence rates could not be calculated. For this reason, the statistical analysis excludes data from persons of unknown race.

^v Cohen, J. 2011. *Selected statistical methods for testing for trends and comparing years or demographic groups in other ACE health-based indicators*. Memorandum submitted to Dan Axelrad, EPA, 16 June, 2011.

Health: Childhood Cancer

Table 1. Statistical significance tests comparing cancer incidence rates for children ages 0 to 19 between age groups for 2017 to 2018 and analyzing trends for 1993-2018.

Type of Cancer	From	To	Against	P-VALUES	
				Unadjusted	Adjusted*
Acute lymphoblastic leukemia	2017	2018	age	< 0.001	< 0.001
Acute lymphoblastic leukemia	1993	2018	year	0.569	0.043
Central nervous system tumors	2017	2018	age	0.931	0.930
Central nervous system tumors	1993	2018	year	< 0.001	< 0.001
Soft tissue sarcomas	2017	2018	age	0.208	NA**
Soft tissue sarcomas	1993	2018	year	0.693	0.320
Hodgkin's lymphoma	2017	2018	age	< 0.001	< 0.001
Hodgkin's lymphoma	1993	2018	year	0.288	0.066
Acute myeloid leukemia	2017	2018	age	0.001	< 0.001
Acute myeloid leukemia	1993	2018	year	0.071	0.083
Non-Hodgkin's lymphoma	2017	2018	age	0.132	0.067
Non-Hodgkin's lymphoma	1993	2018	year	0.293	0.021
Neuroblastoma	2017	2018	age	< 0.001	< 0.001
Neuroblastoma	1993	2018	year	0.009	0.011
Thyroid carcinoma	2017	2018	age	NA**	0.007
Thyroid carcinoma	1993	2018	year	0.932	0.704
Osteosarcoma	2017	2018	age	< 0.001	< 0.001
Osteosarcoma	1993	2018	year	0.485	0.619
Wilms' tumor	2017	2018	age	< 0.001	< 0.001
Wilms' tumor	1993	2018	year	0.913	0.349
Other and unspecified carcinomas†	2017	2018	age	1.000	NA**
Other and unspecified carcinomas†	1993	2018	year	< 0.001	NA**
Ewing's sarcoma	2017	2018	age	0.343	0.301
Ewing's sarcoma	1993	2018	year	< 0.001	< 0.001

*For Against = "age," the p-values are adjusted for sex and race/ethnicity.

For Against = "year," the p-values are adjusted for age, sex, and race/ethnicity.

**NA. Statistical model failed to converge.

†"Other and unspecified carcinomas" represents all carcinomas and other malignant epithelial neoplasms other than thyroid carcinoma and malignant melanoma.

Data Files

The following files are needed to calculate this indicator. All these files together with the SEER*Stat software used to calculate the incidence rates are available from the SEER website: <http://seer.cancer.gov/data/access.html>. The data and analyses used for this indicator were obtained using version 8.3.9.2 of SEER*Stat.

- Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Incidence - SEER Research Data, 13 Registries, Nov 2020 Sub (1992-2018) - Linked To County Attributes - Time Dependent (1990-2018) Income/Rurality, 1969-2019 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released April 2021, based on the November 2020 submission. Incidence data from the SEER 13 registries: Connecticut; Hawaii; Iowa; New Mexico;

Health: Childhood Cancer

Utah; Atlanta, Georgia; Detroit, Michigan; San Francisco-Oakland, California; Seattle-Puget Sound, Washington; Los Angeles, California; San Jose-Monterey, California; Rural Georgia; and the Alaska Native Tumor Registry. Downloaded December 2021.

The databases incorporated into the SEER*Stat software include census population data used to compute the incidence rates and to age-adjust the rates to the 2000 U.S. Standard Population. These population data are obtained by NCI from the Census Bureau. Age-adjustment is explained above.