



2020 National Emissions Inventory Technical Support Document: Waste Disposal – Composting

EPA-454/R-23-001gg
March 2023

2020 National Emissions Inventory Technical Support Document: Waste Disposal - Composting

U.S. Environmental Protection Agency
Office of Air Quality Planning and Standards
Air Quality Assessment Division
Research Triangle Park, NC

Contents

| | |
|--|------|
| List of Tables | i |
| 33 Waste Disposal – Composting | 33-1 |
| 33.1 Sector Descriptions and Overview | 33-1 |
| 33.2 EPA-developed emissions..... | 33-1 |
| 33.2.1 Activity Data | 33-2 |
| 33.2.2 Allocation Procedure | 33-4 |
| 33.2.3 Emissions Factors | 33-6 |
| 33.2.4 Controls..... | 33-6 |
| 33.2.5 Emissions | 33-6 |
| 33.2.6 Sample Calculations..... | 33-7 |
| 33.2.7 Improvements/Changes in the 2020 NEI..... | 33-8 |
| 33.2.8 Puerto Rico and U.S. Virgin Islands Emissions Calculations..... | 33-8 |
| 33.3 References..... | 33-8 |

List of Tables

| | |
|---|------|
| Table 33-1: Composting SCCs in the 2020 NEI | 33-1 |
| Table 33-2: Annual Waste (million tons) generated and recovered in the US in 2015 | 33-2 |
| Table 33-3: State-level food waste composting (tons)..... | 33-3 |
| Table 33-4: Ranges and midpoints for data withheld from state and county business patterns..... | 33-5 |
| Table 33-5: 2016 County Business Pattern for NAICS 562212 in Arizona | 33-5 |
| Table 33-6: Sample calculations for VOC emissions from greenwaste composting in Apache County, AZ | 33-7 |

33 Waste Disposal – Composting

There are five sections in this documentation that discuss nonpoint inventory Waste Disposal. This section discusses Composting, the second section (34) discusses Open Burning – Land Clearing Debris, the third section (35) discusses Open Burning – Residential Household Waste, the fourth section (36) discusses Open Burning – Yard Waste, and the fifth section (37) discusses Publicly-Owned Treatment Works (POTWs). The reason these sources are broken up within this EIS sector is because the EPA methodologies for estimating the emissions are different.

33.1 Sector Descriptions and Overview

Greenwaste composting includes the diversion of yard waste, food waste, and other biogenic waste from landfills to composting facilities. Estimates of emissions of volatile organic compounds (VOC), ammonia (NH₃), and three hazardous air pollutants (HAPs), acetaldehyde; methanol; and naphthalene, from greenwaste composting are based on the amount of food and yard waste composted. Composting of biogenic waste is currently not included in emissions estimates for this category as activity data on this waste type is not available.

Note that this source category does not include the composting of biosolids from wastewater treatment plants or manure management facilities. There are separate SCCs for biosolids (2680001000) and for a mixture of greenwaste and biosolids (2680002000). EPA is not currently estimating emissions for these SCCs. If S/L/Ts report any emissions for the mixture SCC, emissions from the greenwaste portion of that mixture may be duplicative of some or all of the EPA emissions estimates described here. Note also that this source category estimates emissions from composting facilities but does not estimate emissions from backyard composting.

Table 33-1 shows, for composting, the SCCs covered by the EPA estimates and by the State/Local and Tribal agencies that submitted data. The SCC level 3 and 4 SCC descriptions are also provided. The SCC level 1 and leading level 2 descriptions is “Waste Disposal, Treatment, and Recovery; Composting:” for all SCCs.

Table 33-1: Composting SCCs in the 2020 NEI

| SCC | Description | EPA | S/L/T |
|------------|--|-----|-------|
| 2680001000 | 100% Biosolids (e.g., sewage sludge, manure, mixtures of these matls); All Processes | | X |
| 2680002000 | Mixed Waste (e.g., a 50:50 mixture of biosolids and green wastes); All Processes | | X |
| 2680003000 | 100% Green Waste (e.g., residential or municipal yard wastes); All Processes | X | X |

A list of agencies that submitted composting emissions is provided in Section 6.2.3.

33.2 EPA-developed emissions

The calculations for estimating the emissions from greenwaste composting involve first estimating the amount of food and yard waste composted in each county. The amount of state-level food waste composted is available from the EPA report *Food Waste Management in the United States, 2014* [ref 1]. The amount of state-level yard waste composted is estimated by calculating the per-capita amount of

yard waste composted using national data from the EPA report *Advancing Sustainable Materials Management: 2015 Fact Sheet* [ref 2] and multiplying that by the state population. The state-level yard and food waste are summed together and distributed to the counties based on the proportion of employment at solid waste landfills. The total amount of greenwaste composted is multiplied by emissions factors for VOC and NH3 to estimate emissions of these pollutants from greenwaste composting.

33.2.1 Activity Data

The activity data for this source category is the amount of food and yard waste composted, which is estimated using data from two EPA reports: the national-level amount of yard waste composted comes from *Advancing Sustainable Materials Management: 2015 Fact Sheet* and the state-level amount of food waste composted comes from *Food Waste Management in the United States, 2014* [ref 1, ref 2]. Table 33-2 shows the total national-level amount of yard waste generated and recovered for composting.

Table 33-2: Annual Waste (million tons) generated and recovered in the US in 2015

| Material | Waste Generated | Waste Recovered |
|----------------|-----------------|-----------------|
| Yard trimmings | 34.72 | 21.29 |

The values from Table 33-2 are used with the U.S. population in 2020 of 336 million people [ref 3] to determine per-capita values of food and yard waste recovered for composting.

$$PC_{yard,US} = \frac{W_{yard,US}}{P_{US}} \quad (1)$$

Where:

- $PC_{yard,US}$ = Per-capita yard waste recovered for composting in the US, in tons per person per year
- $W_{yard,US}$ = Total annual yard waste recovered in the US, in tons/year
- P_{US} = US population

This calculation results in per-capita values of approximately 0.066 tons per person per year of yard waste recovered for composting. Please note that EPA data on composting does not include backyard composting.

The per-capita yard waste values from equation 1 are multiplied by the population of each state to estimate the state-level amount of yard waste recovered for composting.

$$W_{yard,s} = PC_{yard,US} \times P_s \quad (2)$$

Where:

- $W_{yard,s}$ = Annual yard waste recovered for composting in state s , in tons
- $PC_{yard,US}$ = Per-capita yard waste recovered for composting in the US, in tons per person per year

P_s = Population of state s

EPA reports the amount of food waste composted at the state level in the report *Food Waste Management in the United States, 2015* [Table 3 in ref 1]. These values are shown in Table 33-3. EPA collected these data from state environmental websites and contacts with state agencies. The data year for each state is listed and represents the latest data available. The data were not altered from the original reference for use in this methodology.

Table 33-3: State-level food waste composting (tons)

| State | Food Composted | Data Year | State | Food Composted | Data Year |
|---------------|----------------|-----------|----------------|------------------|-----------|
| California | 715,119 | 2012 | Nevada | 35,869 | 2014 |
| Colorado | 29,130 | 2013 | New Hampshire | 110 | 2012 |
| Connecticut | 4,644 | 2013 | New Jersey | 28,634 | 2012 |
| Delaware | 17,626 | 2013 | New York | 44,405 | 2013 |
| Florida | 158,711 | 2014 | North Carolina | 38,014 | 2014 |
| Georgia | 8,021 | 2014 | Ohio | 81,450 | 2014 |
| Hawaii | 39,287 | 2014 | Oregon | 50,143 | 2013 |
| Indiana | 13,525 | 2013 | Pennsylvania | 56,851 | 2013 |
| Iowa | 4,334 | 2010 | Rhode Island | 150 | 2014 |
| Kansas | 1,127 | 2010 | South Carolina | 4,277 | 2014 |
| Maine | 1,658 | 2010 | Tennessee | 1,500 | 2013 |
| Maryland | 69,643 | 2014 | Texas | 188 | 2012 |
| Massachusetts | 2,753 | 2014 | Vermont | 14,738 | 2013 |
| Michigan | 8,700 | 2013 | Virginia | 2,454 | 2014 |
| Minnesota | 46,751 | 2013 | Washington | 65,221 | 2013 |
| Mississippi | 242 | 2013 | Wisconsin | 8,677 | 2013 |
| Missouri | 16,000 | 2014 | Total | 1,569,952 | |

The state-level amount of total greenwaste composted is the sum of the state-level food and yard waste composted.

$$W_{GW,s} = W_{yard,s} + W_{food,s} \quad (3)$$

Where:

$W_{GW,s}$ = Annual total greenwaste recovered for composting in state s , in tons

$W_{yard,s}$ = Annual yard waste recovered for composting in state s , in tons

$W_{food,s}$ = Annual food waste recovered for composting in state s , in tons, from Table 33-3

The process used to distribute the state-level amount of greenwaste composted to the counties is discussed in next section.

33.2.2 Allocation Procedure

Comprehensive data on the county locations of composting facilities is not available. As a result, the analysis assumes that greenwaste composting facilities are co-located with solid waste landfills.

State-level food greenwaste composting activity (from equation 3) is allocated to the county-level using employment at solid waste landfills (NAICS code 562212). Specifically, state-level estimates of greenwaste collected for composting are multiplied by the ratio of county- to state- level number of employees at landfills.

$$EmpFrac_c = \frac{Emp_c}{Emp_s} \quad (4)$$

Where:

- $EmpFrac_c$ = The fraction of landfill employees in county c
- Emp_c = The number of landfill employees in county c
- Emp_s = The number of landfill employees in state s

Employment data are from the U.S. Census Bureau's 2016 County Business Patterns (CBP) [ref 4]. Due to concerns with releasing confidential business information, the CBP does not release exact numbers for a given NAICS code if the data can be traced to an individual business. Instead, a series of range codes is used. Many counties and some smaller states have only one solid waste landfill, leading to withheld data in the county and/or state business pattern data. To estimate employment in counties and states with withheld data, the following procedure is used for NAICS code 562212.

To gap-fill withheld state-level employment data:

- a. State-level data for states with known employment in NAICS 562212 are summed to the national level.
- b. The total sum of state-level known employment from step a is subtracted from the national total reported employment for NAICS 562212 in the national-level CBP to determine the employment total for the withheld states.
- c. Each of the withheld states is assigned the midpoint of the range code reported for that state. Table 33-4 lists the range codes and midpoints.
- d. The midpoints for the states with withheld data are summed to the national level.
- e. An adjustment factor is created by dividing the number of withheld employees (calculated in step b of this section) by the sum of the midpoints (step d).
- f. For the states with withheld employment data, the midpoint of the range for that state (step c) is multiplied by the adjustment factor (step e) to calculate the adjusted state-level employment for landfills.

These same steps are then followed to fill in withheld data in the county-level business patterns.

- g. County-level data for counties with known employment are summed by state.
- h. County-level known employment is subtracted from the state total reported in state-level CBP (or, if the state-level data are withheld, from the state total estimated using the procedure discussed above).

- i. Each of the withheld counties is assigned the midpoint of the range code (Table 33-4).
- j. The midpoints for the counties with withheld data are summed to the state level.
- k. An adjustment factor is created by dividing the number of withheld employees (step h) by the sum of the midpoints (step j).
- l. For counties with withheld employment data, the midpoints (step i) are multiplied by the adjustment factor (step k) to calculate the adjusted county-level employment for landfills.

Table 33-4: Ranges and midpoints for data withheld from state and county business patterns

| Employment Code | Ranges | Midpoint |
|-----------------|---------------|----------|
| A | 0-19 | 10 |
| B | 20-99 | 60 |
| C | 100-249 | 175 |
| E | 250-499 | 375 |
| F | 500-999 | 750 |
| G | 1,000-2,499 | 1,750 |
| H | 2,500-4,999 | 3,750 |
| I | 5,000-9,999 | 7,500 |
| J | 10,000-24,999 | 17,500 |
| K | 25,000-49,999 | 37,500 |
| L | 50,000-99,999 | 75,000 |
| M | 100,000+ | |

For example, take the 2016 CBP data for NAICS 562212 (Landfills) in Arizona provided in Table 33-5.

Table 33-5: 2016 County Business Pattern for NAICS 562212 in Arizona

| State FIPS | County FIPS | County Name | NAICS | Employment Code | Employment |
|------------|-------------|-------------|--------|-----------------|------------|
| 04 | 001 | Apache | 562212 | B | withheld |
| 04 | 007 | Gila | 562212 | A | withheld |
| 04 | 012 | La Paz | 562212 | A | withheld |
| 04 | 013 | Maricopa | 562212 | | 296 |
| 04 | 015 | Mohave | 562212 | B | withheld |
| 04 | 017 | Navajo | 562212 | B | withheld |
| 04 | 021 | Pinal | 562212 | | 40 |
| 04 | 023 | Santa Cruz | 562212 | | withheld |
| 04 | 025 | Yavapai | 562212 | A | withheld |
| 04 | 027 | Yuma | 562212 | B | withheld |

Note: Counties in Arizona that do not have employment in solid waste landfills are excluded from this table.

1. The total number of known county-level employees in Arizona is 336.
2. The state-level CBP reports 522 employees for NAICS 562212 in Arizona. This means there are 186 employees total for the 8 counties for which data are withheld.

3. The counties with withheld data are assigned midpoints according to their employment code in Table 33-4. For example, Apache County is given a midpoint of 60 employees (since range code B is 20-99) and Gila County is given a midpoint of 10 employees.
4. The state total of the midpoints for all withheld counties is 270 employees.
5. The adjustment factor is $186/272 = 0.6889$.
6. The adjusted employment for Apache County is $60 \times 0.6889 = 41$. Gila County has an adjusted employment of $10 \times 0.6889 = 7$ employees.

Once county- and state-level employment have been estimated, the ratio of county to state employees (from equation 4) is calculated and multiplied by the state-level greenwaste recovered for composting (from equation 3) to calculate the amount of waste composted in each county.

$$W_{GW,c} = EmpFrac_c \times W_{GW,s} \quad (5)$$

Where:

- $W_{GW,c}$ = Annual total greenwaste composted in county c , in tons
- $W_{GW,s}$ = Annual total greenwaste recovered for composting in state s , in tons
- $EmpFrac_c$ = The fraction of landfill employees in county c

33.2.3 Emissions Factors

Emissions factors for greenwaste composting (SCC=2680003000) are provided in the “Wagon Wheel Emission Factor Compendium” on the [2020 NEI Supporting Data and Summaries site](#). The emissions factors for VOC and ammonia (NH3) are taken from the California Air Resources Board Emissions Inventory Methodology for Composting Facilities [ref 5] and are unaltered from the original reference. The emissions factors for the HAPs (acetaldehyde, methanol, and naphthalene) are taken from Kumar et al [ref 6].

33.2.4 Controls

There are no controls assumed for this category.

33.2.5 Emissions

Emission factors are provided in the “Wagon Wheel Emission Factor Compendium” on the [2020 NEI Supporting Data and Summaries site](#). For VOC and NH3, the emissions are multiplied by a conversion factor to convert from pounds to tons.

$$E_{p,c} = W_{GW,c} \times EF_p \times \frac{1 \text{ ton}}{2000 \text{ lbs.}} \quad (6)$$

Where:

- $E_{p,c}$ = Annual emissions of pollutant p in county c , in tons for VOC and NH3 and lbs. for HAPs

$W_{GW,c}$ = Annual total greenwaste recovered for composting in state s , in tons
 EF_p = Emissions factor for pollutant p , in tons of pollutant per ton of greenwaste composted

Emissions of HAPs are estimated by applying speciation factors (available in the zip file “HAPAugmentation_Nonpoint_28jan2023”, on the [2020 NEI Supplemental data FTP site](#)) to annual VOC emissions. For HAPS, no conversion factor is needed, and the emissions are reported in tons.

$$E_{h,c} = E_{VOC,c} \times SF_h \quad (7)$$

Where:

$E_{h,c}$ = Annual emissions of HAP h in county c , in tons per year
 $E_{VOC,c}$ = Annual VOC emissions in county c , in tons
 SF_h = Speciation factor for HAP h , in tons of HAP emissions per ton of VOC emissions

33.2.6 Sample Calculations

Table 33-6 lists sample calculations to determine the VOC emissions from composting of greenwaste in Apache County, Arizona.

Table 33-6: Sample calculations for VOC emissions from greenwaste composting in Apache County, AZ

| Eq. # | Equation | Values for Apache County, AZ | Result |
|-------|---|---|---|
| 1 | $PC_{yard,US} = \frac{W_{yard,US}}{P_{US}}$ | $\frac{21.08 \text{ million tons yard waste}}{329 \text{ million people}}$ | 0.064 tons yard waste per person per year |
| 2 | $W_{yard,s} = PC_{yard,US} \times P_s$ | $0.064 \text{ tons yard waste per person} \times 7,016,270 \text{ people in AZ}$ | 449,041 tons yard waste composted in AZ |
| 3 | $W_{GW,s} = W_{yard,s} + W_{food,s}$ | $449,041 \text{ tons yard waste} + 0 \text{ tons food waste}$ | 443,520 tons greenwaste composted in AZ |
| 4 | $EmpFrac_c = \frac{Emp_c}{Emp_s}$ | $\frac{41 \text{ employees in Apache County}}{522 \text{ employees in AZ}}$ | 0.079 fraction of solid waste employees in Apache County, AZ |
| 5 | $W_{GW,c} = EmpFrac_c \times W_{GW,s}$ | $0.079 \text{ fraction} \times 443,520 \text{ tons greenwaste composted}$ | 35,038 tons greenwaste composted in Apache County, AZ |
| 6 | $E_{p,c} = W_{GW,c} \times EF_p \times \frac{1 \text{ ton}}{2000 \text{ lbs.}}$ | $35,038 \text{ tons greenwaste} \times 4.67 \text{ lbs. VOC per ton greenwaste} \times \frac{1 \text{ ton}}{2000 \text{ lbs.}}$ | 82 tons VOC emissions from greenwaste composting in Apache County, AZ |

33.2.7 Improvements/Changes in the 2020 NEI

There are no significant changes from the methodology used to calculate the 2020 NEI emissions.

33.2.8 Puerto Rico and U.S. Virgin Islands Emissions Calculations

Emissions from Puerto Rico are calculated using the same method described above. For the U.S. Virgin Islands, emissions are calculated using 2020 population data [ref 7], since 2014 Census Data does not exist for the U.S. Virgin Islands.

33.3 References

1. U.S. EPA. 2016. [Food Waste Management in the United States, 2014](#). Office of Resource Conservation and Recovery.
2. U.S. EPA. 2018. [Advancing Sustainable Materials Management: Facts and Figures Report](#), 2015 Facts and Figures Sheet, Table1. Generation, Recovery, and Discards of Products in MSW, 2015.
3. U.S. Census Bureau. [2020 Total Population](#), American Community Survey.
4. U.S. Census Bureau. [2020 County Business Patterns](#).
5. California Air Resources Board. 2015. [ARB Emissions Inventory Methodology for Composting Facilities](#). Table A-4. Emission factor data taken from [Draft Staff Report on Proposed Amended Rule 1133.1 \(chipping and grinding activities\) and Proposed Rule 1133.3](#) (emissions reductions from greenwaste composting operations), Table III-3.
6. Kumar, A., C.P. Alaimo, R. Horowitz, F.M. Mitloehner, M.J. Kleeman, and P.G. Green. 2011. Volatile organic compound emissions from green waste composting: Characterization and ozone formation. *Atmospheric Environment*, 45:1841-1848.
7. U.S. Census Bureau. 2020. [County Population Totals](#).

United States
Environmental Protection
Agency

Office of Air Quality Planning and Standards
Air Quality Assessment Division
Research Triangle Park, NC

Publication No. EPA-454/R-23-001gg
March 2023
