

Risk Evaluation for 1,2-Dichloroethane

Supplemental Information File:

Benchmark Dose Modeling Results for 1,2-Dichloroethane

CASRN 107-06-2




TABLE OF CONTENTS

| | | |
|-------------|---|-----------|
| 1 | BENCHMARK DOSE MODELING RESULTS FOR 1,2-DICHLOROETHANE | 21 |
| 1.1 | Non-cancer Endpoints | 22 |
| 1.1.1 | Inhalation Data..... | 22 |
| 1.1.1.1 | Acute..... | 22 |
| 1.1.1.1.1 | Mortality..... | 22 |
| 1.1.1.1.1.1 | Mortality in Male B6C3F1 Mice – 4-Hour Inhalation Exposure | 22 |
| 1.1.1.1.2 | Respiratory Effects..... | 27 |
| 1.1.1.1.2.1 | Degeneration with Necrosis of the Olfactory Mucosa in Male F344 Rats – 4-Hour Inhalation Exposure | 27 |
| 1.1.1.1.2.2 | Degeneration with Necrosis of the Olfactory Mucosa in Female F344 Rats – 4-Hour Inhalation Exposure..... | 31 |
| 1.1.1.1.2.3 | Degeneration with Necrosis of the Olfactory Mucosa in Male and Female F344 Rats (Combined) – 4-Hour Inhalation Exposure | 34 |
| 1.1.1.1.2.4 | Degeneration with Necrosis of the Olfactory Mucosa in Male F344 Rats – 8-Hour Inhalation Exposure | 38 |
| 1.1.1.1.2.5 | Degeneration with Necrosis of the Olfactory Mucosa in Female F344 Rats – 8-Hour Inhalation Exposure..... | 41 |
| 1.1.1.1.2.6 | Degeneration with Necrosis of the Olfactory Mucosa in Male and Female F344 Rats (Combined) – 8-Hour Inhalation Exposure | 44 |
| 1.1.1.1.2.7 | Regeneration of the Olfactory Mucosa in Male F344 Rats – 4-Hour Inhalation Exposure | 47 |
| 1.1.1.1.2.8 | Regeneration of the Olfactory Mucosa in Female F344 Rats – 4-Hour Inhalation Exposure | 51 |
| 1.1.1.1.2.9 | Regeneration of the Olfactory Mucosa in Male and Female F344 Rats (Combined) – 4-Hour Inhalation Exposure | 54 |
| 1.1.1.1.3 | Hepatic Effects..... | 57 |
| 1.1.1.1.4 | Renal Effects | 57 |
| 1.1.1.1.4.1 | Relative Kidney Weight (Kidney Weight/100 g Body Weight) in Male B6C3F1 Mice – 4-Hour Inhalation Exposure..... | 58 |
| 1.1.1.2 | Short-term/Intermediate | 61 |
| 1.1.1.2.1 | Mortality..... | 61 |
| 1.1.1.2.1.1 | Mortality in Male Rats – 30-Day Inhalation Exposure..... | 61 |
| 1.1.1.2.1.2 | Mortality in Female Rabbits – Inhalation Exposure on GD 6 to 18 | 65 |
| 1.1.1.2.2 | Body Weight effects..... | 68 |
| 1.1.1.2.2.1 | Body Weight in Male Mice – 28-Day Inhalation Exposure | 68 |
| 1.1.1.2.2.2 | Body Weight Gain in Male Rats – 30-Day Inhalation Exposure..... | 72 |
| 1.1.1.2.3 | Hepatic Effects:..... | 75 |
| 1.1.1.2.3.1 | Relative Liver Weight in Male Mice – 28-Day Inhalation Exposure | 75 |
| 1.1.1.2.3.2 | Serum ALT in Male Mice – 28-Day Inhalation Exposure | 78 |
| 1.1.1.2.3.3 | Serum AST in Male Mice – 28-Day Inhalation Exposure..... | 81 |
| 1.1.1.2.4 | Reproductive Effects..... | 85 |
| 1.1.1.2.4.1 | Sperm Concentration in Male Mice – 4-Week Inhalation Exposure..... | 85 |
| 1.1.1.2.4.2 | Diameter of Seminiferous Tubules in Male Mice – 4-Week Inhalation Exposure | 89 |
| 1.1.1.2.4.3 | Height of Germinal Epithelium in Male Mice – 4-Week Inhalation Exposure | 92 |
| 1.1.1.2.4.4 | Number of Apoptotic Cells in the Testis in Male Mice – 4-Week Inhalation Exposure | 95 |

| | | |
|-------------|---|-----|
| 1.1.1.3 | Chronic | 98 |
| 1.1.1.3.1 | Hepatic Effects | 98 |
| 1.1.1.3.1.1 | LDH Levels in Female Sprague-Dawley Rats – 12-Month Inhalation Exposure | 98 |
| 1.1.1.3.2 | Renal Effects | 101 |
| 1.1.1.3.2.1 | Blood Urea Nitrogen (BUN) Levels in Male Sprague-Dawley Rats – 12-Month Inhalation Exposure | 101 |
| 1.1.1.3.2.2 | Blood Urea Nitrogen (BUN) in Female Sprague-Dawley Rats – 12-Month Inhalation Exposure | 104 |
| 1.1.1.3.2.3 | Calcium Levels in Male Sprague-Dawley Rats – 12-Month Inhalation Exposure | 107 |
| 1.1.1.3.2.4 | Serum Potassium Levels in Male Sprague-Dawley Rats – 12-Month Inhalation Exposure | 111 |
| 1.1.1.3.2.5 | Serum Potassium Levels in Female Sprague-Dawley Rats – 12-Month Inhalation Exposure | 115 |
| 1.1.1.4 | Developmental | 119 |
| 1.1.1.4.1 | Developmental Effects | 119 |
| 1.1.1.4.1.1 | Body Weight in Male Weanling F1 _B Rats | 119 |
| 1.1.2 | Oral Data | 123 |
| 1.1.2.1 | Acute | 123 |
| 1.1.2.1.1 | Mortality | 123 |
| 1.1.2.1.1.1 | Mortality in Male B6C3F1 Mice – Single Oral Gavage | 123 |
| 1.1.2.1.2 | Hepatic Effects | 126 |
| 1.1.2.1.2.1 | Relative Liver Weight in Male B6C3F1 Mice – Single Oral Gavage | 127 |
| 1.1.2.1.3 | Renal Effects | 132 |
| 1.1.2.1.3.1 | Relative Kidney Weight in Male B6C3F1 Mice – Single Oral Gavage | 132 |
| 1.1.2.1.3.2 | Blood Urea Nitrogen in Male B6C3F1 Mice – Single Oral Gavage | 135 |
| 1.1.2.2 | Short-term/Intermediate | 140 |
| 1.1.2.2.1 | Mortality | 140 |
| 1.1.2.2.1.1 | Mortality in F344 Male Rats – 13-Week Gavage | 140 |
| 1.1.2.2.1.2 | Mortality in F344 Female Rats – 13-Week Gavage | 145 |
| 1.1.2.2.2 | Body Weight Effects | 151 |
| 1.1.2.2.2.1 | Maternal Absolute Weight Gain in Female Rats – Oral Gavage, GD 6 to 20 | 151 |
| 1.1.2.2.3 | Renal Effects | 155 |
| 1.1.2.2.3.1 | Absolute Kidney Weight in F344 Male Rats – 13-Week Gavage | 155 |
| 1.1.2.2.3.2 | Relative Kidney Weight in F344 Male Rats – 13-Week Gavage | 158 |
| 1.1.2.2.4 | Immune Effects | 161 |
| 1.1.2.2.4.1 | Leukocyte Count in CD-1 Male Mice – 14-day Gavage Study | 162 |
| 1.1.2.2.4.2 | Thymus Necrosis in F344 Male Rats – One Time per Day, 5 days per Week | 166 |
| 1.1.2.2.4.3 | Thymus Necrosis in F344 Female Rats – 13-Week Gavage | 169 |
| 1.2 | Cancer Endpoints | 172 |
| 1.2.1 | Rat Data | 173 |
| 1.2.1.1 | Tumor Incidence in Male Rats | 173 |
| 1.2.1.1.1 | Subcutaneous Fibromas in Male Rats | 173 |
| 1.2.1.1.2 | Mammary Gland Fibroadenomas in Male Rats | 175 |
| 1.2.1.1.3 | Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats | 178 |
| 1.2.1.1.4 | Peritoneal Mesothelioma in Male Rats | 181 |
| 1.2.1.1.5 | Combined Mammary Gland, Subcutaneous, and Peritoneal Tumors in Male Rats | 184 |

| | | |
|------------------------|--|------------|
| 1.2.1.2 | Tumor Incidence in Female Rats..... | 186 |
| 1.2.1.2.1 | Subcutaneous Fibromas in Female Rats | 186 |
| 1.2.1.2.2 | Mammary Gland Adenomas in Female Rats | 188 |
| 1.2.1.2.3 | Mammary Gland Fibroadenomas in Female Rats..... | 191 |
| 1.2.1.2.4 | Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats..... | 194 |
| 1.2.1.2.5 | Mammary Gland Adenocarcinomas in Female Rats | 197 |
| 1.2.1.2.6 | Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats | 200 |
| 1.2.1.2.7 | Combined Mammary Gland and Subcutaneous Tumors in Female Rats | 203 |
| 1.2.2 | Mouse Data..... | 205 |
| 1.2.2.1 | Bronchiolo-Alveolar Adenomas in Female Mice..... | 205 |
| 1.2.2.2 | Bronchiolo-Alveolar Carcinomas in Female Mice..... | 207 |
| 1.2.2.3 | Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice | 210 |
| 1.2.2.4 | Endometrial Stromal Polyps in Female Mice..... | 213 |
| 1.2.2.5 | Mammary Gland Adenocarcinomas in Female Mice..... | 216 |
| 1.2.2.6 | Hepatocellular Adenomas in Female Mice | 219 |
| 1.2.2.7 | Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice | 222 |
| 1.2.2.8 | Combined Lung, Uterine, Mammary Gland, and Liver Tumors in Female Mice..... | 225 |
| 1.2.2.9 | Combined Lung, Mammary Gland, and Liver Tumors in Female Mice (Alternate Analysis of Combined Tumors with Uterine Polyps Excluded) | 227 |
| REFERENCES..... | | 230 |

LIST OF TABLES

| | | |
|------------|---|----|
| Table 1-1. | Incidence of Mortality in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 22 |
| Table 1-2. | BMD Modeling Results for Increased Incidence of Mortality in Male Mice Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 24 |
| Table 1-3. | Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 28 |
| Table 1-4. | BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 29 |
| Table 1-5. | Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 32 |
| Table 1-6. | BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 32 |
| Table 1-7. | Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 35 |
| Table 1-8. | BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 36 |
| Table 1-9. | Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 39 |

| | |
|---|----|
| Table 1-10. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 39 |
| Table 1-11. Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 42 |
| Table 1-12. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 42 |
| Table 1-13. Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (combined) and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 45 |
| Table 1-14. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 45 |
| Table 1-15. Incidence of Regeneration of the Olfactory Mucosa in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 48 |
| Table 1-16. BMD Modeling Results for Regeneration of the Olfactory Mucosa in Male Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 49 |
| Table 1-17. Incidence of Regeneration of the Olfactory Mucosa in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 52 |
| Table 1-18. BMD Modeling Results for Regeneration of the Olfactory Mucosa in Female Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 52 |
| Table 1-19. Incidence of Regeneration of the Olfactory Mucosa in Male and Female Rats (Combined) and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane | 55 |
| Table 1-20. BMD Modeling Results for Regeneration of the Olfactory Mucosa in Male and Female Rats (Combined) Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane ^a | 55 |
| Table 1-21. Increased Relative Kidney Weight in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Inhalation Exposure Study | 58 |
| Table 1-22. Summary of BMD Modeling Results for Increased Relative Kidney Weight in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane (Constant Variance) ^a | 58 |
| Table 1-23. Incidence of Mortality in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Inhalation Exposure Study | 61 |
| Table 1-24. Summary of BMD Modeling Results for Incidence of Mortality in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane ^a | 62 |
| Table 1-25. Incidence of Mortality in Female Rabbits and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Inhalation Exposure Study .. | 66 |
| Table 1-26. Summary of BMD Modeling Results for Incidence of Mortality in Female Rabbits Following Inhalation Exposure to 1,2-Dichloroethane Using BMR of 10%ER or 5%ER ^a | 66 |
| Table 1-27. Decreased Body Weight of Male Mice at Week 4 and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane in a 28-Day Inhalation Exposure Study | 69 |
| Table 1-28. Summary of BMD Modeling Results for Decreased Body Weight in Male Mice at Week 4 Following a 28-Day Inhalation Exposure to 1,2-Dichloroethane (Nonconstant Variance) ^a | 69 |
| Table 1-29. Decreased Body Weight Gain of Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 30-Day Inhalation Exposure Study | 72 |

| | |
|--|-----|
| Table 1-30. Summary of BMD Modeling Results for Decreased Body Weight Gain of Male Rats Following Inhalation Exposure to 1,2-Dichloroethane for 30 Days (Constant Variance) ^a | 72 |
| Table 1-31. Relative Liver Weight in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 28-Day Inhalation Exposure Study. | 75 |
| Table 1-32. Summary of BMD Modeling Results for Increased Relative Liver Weight in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 28 Days (Constant Variance) ^a | 76 |
| Table 1-33. Serum ALT in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 28-Day Inhalation Exposure Study | 79 |
| Table 1-34. Summary of BMD Modeling Results for Increased Serum ALT in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 28 Days (Nonconstant Variance) ^a | 79 |
| Table 1-35. Serum AST in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 28-Day Inhalation Exposure Study | 82 |
| Table 1-36. Summary of BMD Modeling Results for Increased Serum AST in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 28 Days (Nonconstant Variance) ^a | 82 |
| Table 1-37. Sperm Concentration in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 4-Week Inhalation Exposure Study | 85 |
| Table 1-38. Summary of BMD Modeling Results for Decreased Sperm Concentration in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 4 Weeks (Constant Variance) ^a | 86 |
| Table 1-39. Diameter of Seminiferous Tubules in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 4-Week Inhalation Exposure Study | 89 |
| Table 1-40. Summary of BMD Modeling Results for Decreased Diameter of Seminiferous Tubules in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 4 Weeks (Constant Variance) ^a | 89 |
| Table 1-41. Height of Germinal Epithelium in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 4-Week Inhalation Exposure Study | 92 |
| Table 1-42. Summary of BMD Modeling Results for Decreased Height of Germinal Epithelium in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 4 Weeks (Constant Variance) ^a | 92 |
| Table 1-43. Number of Apoptotic Cells in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 4-Week Inhalation Exposure Study | 95 |
| Table 1-44. Summary of BMD Modeling Results for Increased Number of Apoptotic Cells in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 4 Weeks (Constant Variance) ^a | 95 |
| Table 1-45. Increased LDH Levels in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study | 98 |
| Table 1-46. Summary of BMD Modeling Results for Increased LDH Levels in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Constant Variance) ^a | 99 |
| Table 1-47. BUN Levels in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study | 101 |
| Table 1-48. Summary of BMD Modeling Results for Increased BUN Levels in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Constant Variance) ^a | 102 |

| | |
|--|-----|
| Table 1-49. BUN Levels in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study | 104 |
| Table 1-50. Summary of BMD Modeling Results for Increased BUN Levels in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Nonconstant Variance) ^a | 104 |
| Table 1-51. Serum Calcium Levels in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study | 107 |
| Table 1-52. Summary of BMD Modeling Results for Decreased Serum Calcium Levels in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Constant Variance) ^a | 108 |
| Table 1-53. Serum Potassium Levels in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study | 111 |
| Table 1-54. Summary of BMD Modeling Results for Increased Serum Potassium Levels in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Constant Variance) ^a | 111 |
| Table 1-55. Serum Potassium Levels in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study | 115 |
| Table 1-56. Summary of BMD Modeling Results for Increased Serum Potassium Levels in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months ^a | 116 |
| Table 1-57. Body Weight of Selected F1b Male Weanling Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a One-Generation Developmental Study ^a | 119 |
| Table 1-58. Summary of BMD Modeling Results for Decreased Body Weight of Selected F1b Male Weanling Rats Following Inhalation Exposure to 1,2-Dichloroethane in a One-Generation Reproduction Study (Constant Variance) ^a | 120 |
| Table 1-59. Incidence of Mortality in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Acute Oral Exposure Study..... | 124 |
| Table 1-60. Summary of BMD Modeling Results for Increased Incidence of Mortality in Male Mice Following a Single Oral Exposure to 1,2-Dichloroethane Using BMR of 10%ER or 20%ER ^a | 124 |
| Table 1-61. Relative Liver Weight in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Acute Oral Exposure Study..... | 127 |
| Table 1-62. Summary of BMD Modeling Results for Increased Relative Liver Weight in Male Mice Following a Single Oral Exposure to 1,2-Dichloroethane (Nonconstant Variance) ^a | 128 |
| Table 1-63. Increased Relative Kidney Weight in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Oral Exposure Study | 132 |
| Table 1-64. Summary of BMD Modeling Results for Increased Relative Kidney Weight in Male Mice Following a Single Oral Exposure to 1,2-Dichloroethane (Constant Variance) ^a | 133 |
| Table 1-65. Blood Urea Nitrogen in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Oral Exposure Study | 135 |
| Table 1-66. Summary of BMD Modeling Results for Increased Blood Urea Nitrogen in Male Mice Following a Single Oral Exposure to 1,2-Dichloroethane (Nonconstant Variance) ^a | 136 |
| Table 1-67. Incidence of Mortality in Male Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study..... | 140 |
| Table 1-68. Summary of BMD Modeling Results for Increased Incidence of Mortality in Male Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks ^a | 141 |

| | |
|--|-----|
| Table 1-69. Incidence of Mortality in Female Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study..... | 146 |
| Table 1-70. Summary of BMD Modeling Results for Increased Incidence of Mortality in Female Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks ^a | 147 |
| Table 1-71. Maternal Absolute Weight Gain in Female Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Oral Gestational Exposure Study | 152 |
| Table 1-72. Summary of BMD Modeling Results for Decreased Maternal Absolute Weight Gain in Female Rats Following Oral Exposure to 1,2-Dichloroethane on GD 6 to 20 (Nonconstant Variance) ^a | 152 |
| Table 1-73. Absolute Kidney Weight in Male Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study..... | 155 |
| Table 1-74. Summary of BMD Modeling Results for Increased Absolute Kidney Weight in Male Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks (Nonconstant Variance) ^a | 155 |
| Table 1-75. Relative Kidney Weight in Male Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study..... | 158 |
| Table 1-76. Summary of BMD Modeling Results for Increased Relative Kidney Weight in Male Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks (Constant Variance) ^a ... | 159 |
| Table 1-77. Leukocytes in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 14-Day Oral Exposure Study..... | 162 |
| Table 1-78. Summary of BMD Modeling Results for Decreased Leukocytes in Male Mice Following Oral Exposure to 1,2-Dichloroethane for 14 Days (Nonconstant Variance) ^a | 162 |
| Table 1-79. Incidence of Thymus Necrosis in Male Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study | 166 |
| Table 1-80. Summary of BMD Modeling Results for Increased Incidence of Thymus Necrosis in Male Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks ^a | 167 |
| Table 1-81. Increased Incidence of Thymus Necrosis in Female Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study | 170 |
| Table 1-82. Summary of BMD Modeling Results for Increased Incidence of Thymus Necrosis in Female Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks ^a | 170 |
| Table 1-83. Increased Incidence of Subcutaneous Fibromas in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 173 |
| Table 1-84. Summary of BMD Modeling Results for Increased Incidence of Subcutaneous Fibromas in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 173 |
| Table 1-85. Increased Incidence of Mammary Gland Fibroadenomas in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 176 |
| Table 1-86. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Fibroadenomas in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 176 |
| Table 1-87. Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay..... | 179 |

| | |
|---|-----|
| Table 1-88. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 179 |
| Table 1-89. Increased Incidence of Peritoneal Mesothelioma in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 182 |
| Table 1-90. Summary of BMD Modeling Results for Increased Incidence of Peritoneal Mesothelioma in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 182 |
| Table 1-91. Summary of BMD Multi-Tumor (MS Combo) Modeling Results for Increased Incidence of Subcutaneous Fibromas, Mammary Gland Adenomas and Fibroadenomas, and Peritoneal Mesothelioma (Combined) in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay | 185 |
| Table 1-92. Increased Incidence of Subcutaneous Fibromas in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 186 |
| Table 1-93. Summary of BMD Modeling Results for Increased Incidence of Subcutaneous Fibromas in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 187 |
| Table 1-94. Increased Incidence of Mammary Gland Adenomas in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 189 |
| Table 1-95. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenomas in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 189 |
| Table 1-96. Increased Incidence of Mammary Gland Fibroadenomas in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 192 |
| Table 1-97. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Fibroadenomas in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 192 |
| Table 1-98. Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay..... | 195 |
| Table 1-99. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 195 |
| Table 1-100. Increased Incidence of Mammary Gland Adenocarcinomas in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 198 |
| Table 1-101. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenocarcinomas in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 198 |
| Table 1-102. Increased Incidence of Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 201 |

| | |
|---|-----|
| Table 1-103. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a . | 201 |
| Table 1-104. Summary of BMD Multi-Tumor (MS Combo) Modeling Results for Increased Incidence of Subcutaneous Fibromas and Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay..... | 204 |
| Table 1-105. Increased Incidence of Bronchiolo-Alveolar Adenomas in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 205 |
| Table 1-106. Summary of BMD Modeling Results for Increased Incidence of Bronchiolo-Alveolar Adenomas in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 206 |
| Table 1-107. Increased Incidence of Bronchiolo-Alveolar Carcinomas in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-year Chronic Bioassay..... | 208 |
| Table 1-108. Summary of BMD Modeling Results for Increased Incidence of Bronchiolo-Alveolar Carcinomas in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 208 |
| Table 1-109. Increased Incidence of Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay..... | 211 |
| Table 1-110. Summary of BMD Modeling Results for Increased Incidence of Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 211 |
| Table 1-111. Increased Incidence of Endometrial Stromal Polyps in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 214 |
| Table 1-112. Summary of BMD Modeling Results for Increased Incidence of Endometrial Stromal Polyps in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 214 |
| Table 1-113. Increased Incidence of Mammary Gland Adenocarcinomas in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay..... | 217 |
| Table 1-114. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenocarcinomas in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 217 |
| Table 1-115. Increased Incidence of Hepatocellular Adenomas in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay | 220 |
| Table 1-116. Summary of BMD Modeling Results for Increased Incidence of Hepatocellular Adenomas in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 220 |
| Table 1-117. Increased Incidence of Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay..... | 223 |
| Table 1-118. Summary of BMD Modeling Results for Increased Incidence of Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay ^a | 223 |

| | |
|---|-----|
| Table 1-119. Summary of BMD Multi-Tumor (MS Combo) Modeling Results for Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Uterine Endometrial Stromal Polyps, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay..... | 226 |
| Table 1-120. Summary of BMD Multi-Tumor (MS Combo) Modeling Results for Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay | 228 |

LIST OF FIGURES

| | |
|---|----|
| Figure 1-1. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for Four Hours and BMR of 10%ER | 25 |
| Figure 1-2. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for Four Hours and BMR of 1%ER | 25 |
| Figure 1-3. Details Regarding the Selected Model (Multistage 1-Degree) for Mortality in Male Mice Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane | 26 |
| Figure 1-4. Details Regarding the Selected Model (Multistage 2-Degree) for Mortality in Male Mice Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane | 27 |
| Figure 1-5. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER..... | 30 |
| Figure 1-6. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane | 31 |
| Figure 1-7. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for Four Hours and BMR of 10%ER | 33 |
| Figure 1-8. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane | 34 |
| Figure 1-9. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER..... | 37 |
| Figure 1-10. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane..... | 38 |
| Figure 1-11. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 8 Hours and BMR of 10%ER..... | 40 |
| Figure 1-12. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane | 41 |
| Figure 1-13. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 8 Hours and BMR of 10%ER..... | 43 |

| | |
|---|----|
| Figure 1-14. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane | 44 |
| Figure 1-15. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Exposed to 1,2-Dichloroethane Via Inhalation for 8 Hours and BMR of 10%ER | 46 |
| Figure 1-16. Details Regarding the Selected Model (Multistage 3-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane..... | 47 |
| Figure 1-17. Plot of Response by Concentration with Fitted Curve for the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER | 50 |
| Figure 1-18. Details Regarding the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Male Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane .. | 51 |
| Figure 1-19. Plot of Response by Concentration with Fitted Curve for the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER | 53 |
| Figure 1-20. Details Regarding the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Female Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane..... | 54 |
| Figure 1-21. Plot of Response by Concentration with Fitted Curve for the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Male and Female Rats (Combined) Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER..... | 56 |
| Figure 1-22. Details Regarding the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Male and Female Rats (Combined) Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane | 57 |
| Figure 1-23. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 1SD (Constant Variance) | 59 |
| Figure 1-24. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%RD (Constant Variance) .. | 59 |
| Figure 1-25. Details Regarding the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours..... | 60 |
| Figure 1-26. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days (Seven Hours per Day, 5 days per Week) and BMR of 10%ER | 63 |
| Figure 1-27. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days (Seven Hours per Day, 5 days per Week) and BMR of 5%ER | 63 |
| Figure 1-28. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days (7n Hours per Day, 5 Days per Week) and BMR of 1%ER | 64 |
| Figure 1-29. Details Regarding the Selected Model (Multistage 3-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane 0Via Inhalation for 30 Days..... | 65 |

| | |
|---|----|
| Figure 1-30. Plot of Response by Dose with Fitted Curve for the Selected Model (Log-Logistic) for Mortality in Female Rabbits Exposed to 1,2-Dichloroethane Via Inhalation on GD 6 to 18 (7 Hours per Day) and BMR of 10%ER..... | 67 |
| Figure 1-31. Plot of Response by Dose with Fitted Curve for the Selected Model (Log-Logistic) for Mortality in Female Rabbits Exposed to 1,2-Dichloroethane Via Inhalation on GD 6 to 18 (7 Hours per Day) and BMR of 5%ER..... | 67 |
| Figure 1-32. Details Regarding the Selected Model (Log-Logistic) for Mortality in Female Rabbits Exposed to 1,2-Dichloroethane Via Inhalation on GD 6 to 18 (7 Hours per Day)..... | 68 |
| Figure 1-33. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Body Weight in Male Mice at Week 4 Following a 28-Day Inhalation Exposure to 1,2-Dichloroethane and a BMR of 1SD (Nonconstant Variance)..... | 70 |
| Figure 1-34. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Body Weight in Male Mice at Week 4 Following a 28-Day Inhalation Exposure to 1,2-Dichloroethane and a BMR of 10%RD (Nonconstant Variance)..... | 70 |
| Figure 1-35. Details Regarding the Selected Model (Exponential 3) for Decreased Body Weight in Male Mice at Week 4 Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days..... | 71 |
| Figure 1-36. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Body Weight Gain in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days and BMR of 1SD (Constant Variance)..... | 73 |
| Figure 1-37. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Body Weight Gain in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days and BMR of 10%RD (Constant Variance)..... | 73 |
| Figure 1-38. Details Regarding the Selected Model (Polynomial 2-Degree) for Body Weight Gain in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days..... | 74 |
| Figure 1-39. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days and BMR of 1SD (Constant Variance)..... | 76 |
| Figure 1-40. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days and BMR of 10%RD (Constant Variance)..... | 77 |
| Figure 1-41. Details Regarding the Selected Model (Linear) for Relative Liver Weight Increases in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days..... | 78 |
| Figure 1-42. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Serum ALT in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days and BMR of 1SD (Nonconstant Variance)..... | 80 |
| Figure 1-43. Details Regarding the Selected Model (Linear) for Increased Serum ALT in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days..... | 81 |
| Figure 1-44. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Serum AST in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days and BMR of 1SD (Nonconstant Variance)..... | 83 |
| Figure 1-45. Details Regarding the Selected Model (Linear) for Increased Serum AST in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days..... | 84 |
| Figure 1-46. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Sperm Concentration in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 1SD (Constant Variance)..... | 87 |
| Figure 1-47. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Sperm Concentration in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 5%RD (Constant Variance)..... | 87 |

| | |
|--|-----|
| Figure 1-48. Details Regarding the Selected Model (Exponential 3) for Decreased Sperm Concentration in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks..... | 88 |
| Figure 1-49. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Decreased Diameter of Seminiferous Tubules in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 1SD (Constant Variance)..... | 90 |
| Figure 1-50. Details Regarding the Selected Model (Linear) for Decreased Diameter of Seminiferous Tubules in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks..... | 91 |
| Figure 1-51. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Height of Germinal Epithelium in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 1SD (Constant Variance)..... | 93 |
| Figure 1-52. Details Regarding the Selected Model (Exponential 3) for Decreased Height of Germinal Epithelium in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks..... | 94 |
| Figure 1-53. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Number of Apoptotic Cells in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 1SD (Constant Variance)..... | 96 |
| Figure 1-54. Details Regarding the Selected Model (Linear) for Increased Number of Apoptotic Cells in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks..... | 98 |
| Figure 1-55. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 5) for Increased LDH Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance) | 99 |
| Figure 1-56. Details Regarding the Selected Model (Exponential 5) for Increased LDH Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months..... | 101 |
| Figure 1-57. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 3-Degree) for Increased BUN Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance)..... | 102 |
| Figure 1-58. Details Regarding the Selected Model (Polynomial 3-Degree) for Increased BUN Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months | 103 |
| Figure 1-59. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased BUN Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Nonconstant Variance) | 105 |
| Figure 1-60. Details Regarding the Selected Model (Polynomial 2-Degree) for Increased BUN Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months..... | 106 |
| Figure 1-61. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Decreased Serum Calcium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance)..... | 108 |
| Figure 1-62. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Decreased Serum Calcium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 10%RD (Constant Variance)..... | 109 |
| Figure 1-63. Details Regarding the Selected Model (Hill) for Increased Serum Calcium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months | 110 |
| Figure 1-64. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Increased Serum Potassium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance) | 112 |
| Figure 1-65. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Increased Serum Potassium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 10%RD (Constant Variance) | 113 |

| | |
|---|-----|
| Figure 1-66. Details Regarding the Selected Model (Hill) for Increased Serum Potassium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months | 115 |
| Figure 1-67. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Increased Serum Potassium Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance)..... | 117 |
| Figure 1-68. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Increased Serum Potassium Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 10%RD (Constant Variance) | 117 |
| Figure 1-69. Details Regarding the Selected Model (Hill) for Increased Serum Potassium Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months..... | 118 |
| Figure 1-70. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Decreased Body Weight of Selected F1B Male Weanling Rats Exposed to 1,2-Dichloroethane Via Inhalation in a One-Generation Reproduction Study and BMR of 1SD (Constant Variance) | 121 |
| Figure 1-71. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Decreased Body Weight of Selected F1B Male Weanling Rats Exposed to 1,2-Dichloroethane Via Inhalation in a One-Generation Reproduction Study and BMR of 5%RD (Constant Variance) | 121 |
| Figure 1-72. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Decreased Body Weight of Selected F1B Male Weanling Rats Exposed to 1,2-Dichloroethane Via Inhalation in a One-Generation Reproduction Study and BMR of 10%RD (Constant Variance) | 122 |
| Figure 1-73. Details Regarding the Selected Model (Linear) for Decreased Body Weight of Selected F1B Male Weanling Rats Exposed to 1,2-Dichloroethane Via Inhalation in a One-Generation Reproduction Study..... | 123 |
| Figure 1-74. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Incidence of Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 10%ER | 125 |
| Figure 1-75. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Incidence of Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 20%ER | 125 |
| Figure 1-76. Details Regarding the Selected Model (Multistage 3-Degree) for Increased Incidence of Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study..... | 126 |
| Figure 1-77. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 5) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 1SD (Nonconstant Variance) ... | 129 |
| Figure 1-78. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 3-Degree) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 10%RD (Nonconstant Variance) | 129 |
| Figure 1-79. Details Regarding the Selected Model (Exponential 5) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study with a BMR of 1SD | 130 |
| Figure 1-80. Details Regarding the Selected Model (Polynomial 3-Degree) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study with a BMR of 10%RD | 131 |
| Figure 1-81. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2- | |

| | |
|--|-----|
| Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 1SD (Constant Variance) | 133 |
| Figure 1-82. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2- Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 10%RD (Constant Variance) | 134 |
| Figure 1-83. Details Regarding the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study | 135 |
| Figure 1-84. Plot of Response by Dose with Fitted Curve for the Selected Model (Power) for Increased Blood Urea Nitrogen in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 1SD (Nonconstant Variance)..... | 137 |
| Figure 1-85. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Blood Urea Nitrogen in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 10%RD (Nonconstant Variance)..... | 137 |
| Figure 1-86. Details Regarding the Selected Model (Power) for Increased BUN Levels in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study (BMR of 1SD) | 138 |
| Figure 1-87. Details Regarding the Selected Model (Linear) for Increased BUN Levels in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study (BMR of 10%RD) | 139 |
| Figure 1-88. Plot of Response by Dose with Fitted Curve for the Selected Model (Logistic Model) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%ER | 142 |
| Figure 1-89. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 5%ER..... | 143 |
| Figure 1-90. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 1%ER..... | 143 |
| Figure 1-91. Details Regarding the Selected Model (Logistic) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks | 144 |
| Figure 1-92. Details Regarding the Selected Model (Multistage 2-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks | 145 |
| Figure 1-93. Plot of Response by Dose with Fitted Curve for the Selected Model (Weibull) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%ER | 148 |
| Figure 1-94. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 5%ER | 148 |
| Figure 1-95. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 1%ER | 149 |
| Figure 1-96. Details Regarding the Selected Model (Weibull) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks..... | 150 |
| Figure 1-97. Details Regarding the Selected Model (Multistage 2-Degree) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks | 151 |

| | |
|---|-----|
| Figure 1-98. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Decreased Maternal Absolute Weight Gain in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage GD 6 to 20 and BMR of 1SD (Nonconstant Variance)..... | 153 |
| Figure 1-99. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Decreased Maternal Absolute Weight Gain in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage GD 6 to 20 and BMR of 10%RD (Nonconstant Variance)..... | 153 |
| Figure 1-100. Details Regarding the Selected Model (Hill) for Decreased Maternal Absolute Weight Gain in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage GD 6 to 20..... | 154 |
| Figure 1-101. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 5) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 1SD (Nonconstant Variance)..... | 156 |
| Figure 1-102. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 5) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%RD (Nonconstant Variance)..... | 157 |
| Figure 1-103. Details Regarding the Selected Model (Exponential 5) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks.. | 158 |
| Figure 1-104. Plot of Response by Dose with Fitted Curve for the Selected Model (Power) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 1SD (Constant Variance)..... | 160 |
| Figure 1-105. Plot of Response by Dose with Fitted Curve for the Selected Model (Power) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%RD (Constant Variance)..... | 160 |
| Figure 1-106. Details Regarding the Selected Model (Power) for Increased Relative Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks..... | 161 |
| Figure 1-107. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Leukocytes in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage for 14 Days and BMR of 1SD (Nonconstant Variance)..... | 163 |
| Figure 1-108. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Leukocytes in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage for 14 Days and BMR of 10%RD (Nonconstant Variance)..... | 164 |
| Figure 1-109. Details Regarding the Selected Model (Exponential 3) for Decreased Leukocytes in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage for 14 Days..... | 165 |
| Figure 1-110. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Increased Incidence of Thymus Necrosis in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%ER..... | 168 |
| Figure 1-111. Details Regarding the Selected Model (Multistage 3-Degree) for Increased Incidence of Thymus Necrosis in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks..... | 169 |
| Figure 1-112. Plot of Response by Dose with Fitted Curve for the Selected Model (Quantal Linear) for Increased Incidence of Thymus Necrosis in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%ER..... | 171 |
| Figure 1-113. Details Regarding the Selected Model (Quantal Linear) for Increased Incidence of Thymus Necrosis in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks..... | 172 |
| Figure 1-114. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Subcutaneous Fibromas in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 174 |

| | |
|--|-----|
| Figure 1-115. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Subcutaneous Fibromas in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 175 |
| Figure 1-116. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Fibroadenomas in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 177 |
| Figure 1-117. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Fibroadenomas in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 178 |
| Figure 1-118. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 180 |
| Figure 1-119. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 181 |
| Figure 1-120. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Peritoneal Mesothelioma in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 183 |
| Figure 1-121. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Peritoneal Mesothelioma in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 184 |
| Figure 1-122. Plot of Response by Concentration with Fitted Curve for Selected Models used for the Multi-Tumor (MS Combo) Model for Increased Incidence of Subcutaneous Fibromas, Mammary Gland Adenomas and Fibroadenomas, and Peritoneal Mesothelioma (Combined) in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 185 |
| Figure 1-123. Details Regarding the Multi-Tumor (MS Combo) Model for the Increased Incidence of Subcutaneous Fibromas, Mammary Gland Adenomas and Fibroadenomas, and Peritoneal Mesothelioma (Combined) in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 186 |
| Figure 1-124. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Subcutaneous Fibromas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 187 |
| Figure 1-125. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Subcutaneous Fibromas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 188 |
| Figure 1-126. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 190 |
| Figure 1-127. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 191 |
| Figure 1-128. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Fibroadenomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 193 |
| Figure 1-129. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Fibroadenomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 194 |

| | |
|--|-----|
| Figure 1-130. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 196 |
| Figure 1-131. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 197 |
| Figure 1-132. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenocarcinomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 199 |
| Figure 1-133. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenocarcinomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 200 |
| Figure 1-134. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 202 |
| Figure 1-135. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 203 |
| Figure 1-136. Plot of Response by Concentration with Fitted Curve for Selected Models Used for the Multi-Tumor (MS Combo) Model for Increased Incidence of Subcutaneous Fibromas and Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 204 |
| Figure 1-137. Details Regarding the Multi-Tumor (MS Combo) Model for the Increased Incidence of Subcutaneous Fibromas and Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 205 |
| Figure 1-138. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Bronchiolo-Alveolar Adenomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 206 |
| Figure 1-139. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Bronchiolo-Alveolar Adenomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 207 |
| Figure 1-140. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 2-Degree) for the Increased Incidence of Bronchiolo-Alveolar Carcinomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 209 |
| Figure 1-141. Details Regarding the Selected Model (Multistage 2-Degree) for the Increased Incidence of Bronchiolo-Alveolar Carcinomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 210 |
| Figure 1-142. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 2-Degree) for the Increased Incidence of Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 212 |
| Figure 1-143. Details Regarding the Selected Model (Multistage 2-Degree) for the Increased Incidence of Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study) | 213 |

| | |
|--|-----|
| Figure 1-144. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Endometrial Stromal Polyps in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 215 |
| Figure 1-145. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Endometrial Stromal Polyps in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 216 |
| Figure 1-146. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenocarcinomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 218 |
| Figure 1-147. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenocarcinomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 219 |
| Figure 1-148. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Hepatocellular Adenomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 221 |
| Figure 1-149. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Hepatocellular Adenomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 222 |
| Figure 1-150. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 2-Degree) for the Increased Incidence of Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 224 |
| Figure 1-151. Details Regarding the Selected Model (Multistage 2-Degree) for the Increased Incidence of Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 225 |
| Figure 1-152. Plot of Response by Concentration with Fitted Curve for Selected Models used for the Multi-Tumor (MS Combo) Model for Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Uterine Endometrial Stromal Polyps, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 227 |
| Figure 1-153. Details Regarding the Multi-Tumor (MS Combo) Model for the Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Uterine Endometrial Stromal Polyps, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 227 |
| Figure 1-154. Plot of Response by Concentration with Fitted Curve for Selected Models Used for the Multi-Tumor (MS Combo) Model for Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 228 |
| Figure 1-155. Details Regarding the Multi-Tumor (MS Combo) Model for the Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)..... | 229 |

1 BENCHMARK DOSE MODELING RESULTS FOR 1,2-DICHLOROETHANE

EPA performed BMD modeling using EPA's BMD modeling software (BMDS Version 3.3 for continuous and cancer data and Version 3.3.2 for dichotomous non-cancer data) for the health domains that were identified during hazard identification and that received a judgment of "likely" ("evidence indicates that 1,2-dichloroethane exposure likely causes [health effect]") or "suggestive" ("evidence suggests but is not sufficient to conclude that 1,2-dichloroethane exposure causes [health effect]") during evidence integration, including cancer (various tumor types), mortality, body weight effects, respiratory tract effects, renal effects, hepatic effects, immune/hematological effects, and male reproductive effects. Although it was concluded during evidence integration that 1,2-dichloroethane exposure likely causes neurological/behavioral effects, none of the data for this domain were amenable for BMD modeling. EPA conducted BMD modeling in a manner consistent with EPA's *Benchmark Dose (BMD) Technical Guidance* ([U.S. EPA, 2012](#)).

EPA used dichotomous models to fit quantal data (e.g., incidences of tumors) and continuous models to fit continuous data (e.g., body and organ weights), as recommended by EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The BMDs/BMDLs are provided based on a daily exposure (i.e., 7 days per week) for easier comparison across all hazard endpoints and thus, doses were adjusted as needed before BMD modeling. EPA modeled endpoints that had statistically significant pairwise comparisons between individual doses and controls or significant dose-response trends. EPA also considered potential biologically significant changes from controls where possible and/or changes that appeared to exhibit a dose-response relationship upon visual inspection. Multiple health endpoints may have been modeled from each study, depending on the relevance of the data to adverse health outcomes and to identify sensitive health endpoints for each domain.

Although some of the data sets could be fit using models after dropping doses (either one, two, or three of the highest doses), EPA considered only modeling results from full data sets for use in quantifying risk. This document does not present results of modeling exercises in which none of the models in the BMD suite provided an adequate fit to the full data sets. Endpoints were also not considered for BMD modeling if changes were observed only at the highest dose. Studies with LOAELs more than 10 times greater than the most sensitive LOAEL for the health domain were also not considered for BMD modeling. For non-cancer endpoints, if BMD modeling was not possible or when data did not fit the available models, EPA used NOAELs and LOAELs during POD selection for the risk evaluation.

EPA relied on the BMD guidance and other information to choose BMRs appropriate for each endpoint. Although the *BMD Technical Guidance* doesn't recommend default BMRs, it describes how various BMD modeling results compare with NOAEL values, and the guidance does recommend calculating 10% ER for quantal data and one SD for continuous data to compare modeling results across endpoints. EPA also modeled percent RD for certain continuous endpoints. EPA's choice of BMRs for the 1,2-dichloroethane health endpoints is described in more detail in the following sections that present BMD modeling results for each health domain.

When modeling dose-response relationships, the data can be modeled as either ER or additional risk. EPA modeled the data as ER. EPA's *BMD Technical Guidance* defines ER as "a measure of the proportional increase in risk of an adverse effect adjusted for the background incidence of the same effect." Mathematically, ER is equal to $[P(d) - P(0)]/[1 - P(0)]$. P(d) is the probability of the effect at dose d, and P(0) is the probability of risk with no exposure to a hazard ([U.S. EPA, 2012](#)).

1.1 Non-cancer Endpoints

Non-cancer endpoints selected for modeling were based on both dichotomous and continuous measurement data. For dichotomous data, the Gamma, Logistic, Log-Logistic, Log-Probit, Multistage, Probit, Weibull, and Quantal Linear dichotomous models available within the software were fit using the selected BMR. For inhalation data, administered concentrations were modeled in units of mg/m³. Adequacy of model fit was judged based on the χ^2 goodness-of-fit p-value ($p > 0.1$), magnitude of scaled residuals in the vicinity of the BMR, and visual inspection of the model fit. Among all models providing adequate fit, the lowest BMDL was selected if the BMDLs estimated from different models varied > 3 -fold; otherwise, the BMDL from the model with the lowest AIC was selected. For continuous measurement data, the Exponential, Hill, Linear, Polynomial, and Power continuous models available within the software were fit employing the selected BMR(s). An adequate fit was judged based on the chi-square goodness-of-fit p-value ($p > 0.1$), magnitude of the scaled residuals in the vicinity of the BMR, and visual inspection of the model fit. In addition to these three criteria for judging adequacy of model fit, a determination was made as to whether the variance across dose groups was constant. If a constant variance model was deemed appropriate based on the statistical test provided in BMDS (*i.e.*, Test 2; p-value > 0.05 [note: this is a change from previous versions of BMDS, which required variance p-value > 0.10 for adequate fit]), the final BMD results were estimated from a constant variance model. If the test for homogeneity of variance was rejected (p-value < 0.05), the model was run again while modeling the variance as a power function of the mean to account for this nonconstant variance. If this nonconstant variance model also did not adequately fit the data (*i.e.*, Test 3; p-value < 0.05), the data set was considered unsuitable for BMD modeling. Among all models providing adequate fit, the lowest BMDL was selected if the BMDLs estimated from different models varied > 3 -fold; otherwise, the BMDL from the model with the lowest AIC was selected.

1.1.1 Inhalation Data

1.1.1.1 Acute

1.1.1.1.1 Mortality

[Storer et al. \(1984\)](#) provided data showing increased mortality in mice following acute inhalation exposure to 1,2-dichloroethane.

1.1.1.1.1.1 Mortality in Male B6C3F1 Mice – 4-Hour Inhalation Exposure

Increased incidence of mortality was observed in male mice exposed to 1,2-dichloroethane by inhalation for four hours ([Storer et al., 1984](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-1. Dichotomous models were fit to the incidence data. EPA chose a BMR of 10% ER according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)) to compare with other PODs. A BMR of one percent ER was also selected based on the severity of the endpoint.

Table 1-1. Incidence of Mortality in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Concentration (mg/m ³) | Number of Animals | Incidence |
|------------------------------------|-------------------|-----------|
| 0 | 5 | 0 |
| 107 | 5 | 0 |

| Concentration (mg/m ³) | Number of Animals | Incidence |
|---------------------------------------|-------------------|-----------|
| 337 | 5 | 0 |
| 723.2 | 5 | 4 |
| 1313 | 5 | 5 |

The BMD modeling results for incidence of mortality are summarized in Table 1-2. All models provided adequate fit to the data (chi-square p-value > 0.1). Using a BMR of 10% ER, the BMDLs were not sufficiently close (differed by > 3-fold); therefore, the model with the lowest BMDL (Multistage 1-degree) was selected. Using a BMR of one percent ER, the Multistage 1-degree/Quantal Linear model was considered questionable because the BMD and BMDL were 10 times lower than the lowest non-zero concentration. The BMDLs of the remaining models were not sufficiently close; therefore, the BMDS recommended the model with the lowest BMDL (Multistage 2-degree).

This data set is not well suited for BMD modeling; there is a single datapoint (at 80% incidence) with incidence between 0 and 100%; there are no data to inform the shape of the curve at the region of interest, as defined by the BMR. As a result, the different models provide a broad range of BMD and BMDL estimates: an ~6-fold BMDL spread at a BMR of 10% and an ~ 38-fold BMDL spread at a BMR of one percent. For both BMRs, selection of the low end of the range to represent the BMDL is not among the more realistic possibilities and involves extrapolation below the range of observation to generate a BMD estimate well below the lowest tested dose. The much higher spread of BMDLs at a BMR of one percent indicates much higher uncertainty associated with modeling using this BMR, reflecting the greater distance from the lowest observable change in the study, which was only 20% (1/5).

Table 1-2. BMD Modeling Results for Increased Incidence of Mortality in Male Mice Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 1%ER (mg/m ³) | BMDL 1%ER (mg/m ³) | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-----------------|--------------|-------------------------------------|--------------------------------------|--------------------------------------|---------------------------------------|--|
| | p-value | AIC | | | | | |
| Dichotomous Hill | 1.000 | 7.004 | 519 | 130 | 593 | 263 | All models provided adequate fit to the data (chi-square p-value > 0.1). With a BMR of 10% ER applied, the BMDLs were not sufficiently close (differed by > 3-fold); therefore, the Multistage 1-degree model, which has the lowest BMDL was selected. With a BMR of one percent ER applied, the Multistage 1-degree/Quantal Linear model was considered questionable because the BMD and BMDL were 10 times lower than the lowest non-zero concentration. The BMDLs of the remaining models were not sufficiently close; therefore, the model with the lowest BMDL (Multistage 2-degree) was selected. NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |
| Gamma | 0.9994 | 7.140 | 328 | 113 | 438 | 242 | |
| Log-Logistic | 1.000 | 7.004 | 519 | 130 | 593 | 263 | |
| Multistage 3 | 0.9730 | 6.507 | 143 | 13.8 | 313 | 137 | |
| Multistage 2 | 0.7410 | 10.32 | 67.3 | 11.1 | 218 | 103 | |
| Multistage 1 | 0.2538 | 15.59 | 7.25 | 4.11 | 76.0 | 43.1 | |
| Weibull | 0.9998 | 7.083 | 350 | 81 | 491 | 230 | |
| Logistic | 1.000 | 7.005 | 500 | 57.5 | 590 | 261 | |
| Log-Probit | 1.000 | 9.004 | 553 | 157 | 604 | 263 | |
| Probit | 1.000 | 9.004 | 492 | 56.8 | 568 | 246 | |
| Quantal Linear | 0.2538 | 15.59 | 7.25 | 4.11 | 76.0 | 43.1 | |

^a Selected model in bold.

Plots of the Multistage 1-degree and Multistage 2-degree models with BMRs of 10% and one percent ER are shown in Figure 1-1 and Figure 1-2, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown below in Figure 1-3 and Figure 1-4 for the Multistage 1-degree and Multistage 2-degree models, respectively.




Figure 1-1. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for Four Hours and BMR of 10%ER




Figure 1-2. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for Four Hours and BMR of 1%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 75.96529122 | | | | |
| BMDL | 43.13559104 | | | | |
| BMDU | 144.8556487 | | | | |
| AIC | 15.58503443 | | | | |
| P-value | 0.253844445 | | | | |
| D.O.F. | 4 | | | | |
| Chi ² | 5.343368092 | | | | |
| Slope Factor | 0.002318271 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | 0.001386956 | 0.664907502 | -1.3018078 | 1.30458172 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-08 | 0 | 5 | -0.000276 |
| 107 | 0.1379175 | 0.689587501 | 0 | 5 | -0.894376 |
| 337 | 0.373373232 | 1.866866159 | 0 | 5 | -1.726045 |
| 723.2 | 0.633239628 | 3.166198142 | 4 | 5 | 0.7737535 |
| 1313 | 0.838148045 | 4.190740227 | 5 | 5 | 0.9826155 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -2.502012118 | 5 | - | - | NA |
| Fitted Model | -6.792517214 | 1 | 8.58101019 | 4 | 0.0724694 |
| Reduced Model | -16.33545487 | 1 | 27.6668855 | 4 | <0.0001 |

Figure 1-3. Details Regarding the Selected Model (Multistage 1-Degree) for Mortality in Male Mice Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 67.31979018 | | | | |
| BMDL | 11.12084919 | | | | |
| BMDU | 97.41764876 | | | | |
| AIC | 10.32271674 | | | | |
| P-value | 0.740950577 | | | | |
| D.O.F. | 4 | | | | |
| Chi ² | 1.971774077 | | | | |
| Slope Factor | 0.000899212 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 3 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | Bounded | NA | NA | NA | |
| b2 | 2.21766E-06 | 1.668562717 | -3.2703206 | 3.27032508 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-08 | 0 | 5 | -0.000276 |
| 107 | 0.025070411 | 0.125352057 | 0 | 5 | -0.358574 |
| 337 | 0.22264463 | 1.113223149 | 0 | 5 | -1.196689 |
| 723.2 | 0.686475475 | 3.432377376 | 4 | 5 | 0.5471752 |
| 1313 | 0.978141831 | 4.890709155 | 5 | 5 | 0.3342651 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -2.502012118 | 5 | - | - | NA |
| Fitted Model | -4.161358368 | 1 | 3.3186925 | 4 | 0.5059761 |
| Reduced Model | -16.33545487 | 1 | 27.6668855 | 4 | <0.0001 |

Figure 1-4. Details Regarding the Selected Model (Multistage 2-Degree) for Mortality in Male Mice Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2 Respiratory Effects

Incidence data for degeneration with necrosis of the olfactory mucosa following a four- or eight-hour inhalation exposure and regeneration of the olfactory mucosa following a four-hour of exposure were modeled for males, females, and males and females combined ([Dow Chemical, 2006](#)).

1.1.1.1.2.1 Degeneration with Necrosis of the Olfactory Mucosa in Male F344 Rats – 4-Hour Inhalation Exposure

Increased incidence of degeneration with necrosis of the olfactory mucosa was observed in male rats exposed to 1,2-dichloroethane by inhalation for four hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration

adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-3. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA’s *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-3. Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Concentration (mg/m ³) | Number of Animals | Incidence |
|------------------------------------|-------------------|-----------|
| 0 | 5 | 0 |
| 35.3 | 5 | 0 |
| 132.5 | 5 | 3 |
| 410.0 | 5 | 5 |
| 1368.7 | 5 | 5 |

The BMD modeling results for increased incidence of degeneration with necrosis of the olfactory mucosa in male rats are summarized in Table 1-4. All models provided adequate fit to the data (chi-square p-value > 0.1). The BMDL computation for the Weibull model failed because the lower limit included zero. The BMDLs were not sufficiently close (differed by >3-fold). Therefore, EPA chose the model with the lowest BMDL (Multistage 1-degree model).

This data set is not well suited for BMD modeling; there is a single datapoint (at 60% incidence) with incidence between 0 and 100%; there are no data to inform the shape of the curve at the region of interest (~10%). As a result, the different models provide a broad range of BMD and BMDL estimates. Selection of the low end of the range to represent the BMDL is not among the more realistic possibilities

and in this case involves extrapolation below the range of observation to generate a BMD estimate well below the lowest tested dose.

Table 1-4. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-----------------|--------------|--------------------------------------|---------------------------------------|---|
| | p-value | AIC | | | |
| Dichotomous Hill | 1.000 | 10.73 | 93.8 | 24.9 | All models provided adequate fit to the data (chi-square p-value > 0.1); however, the BMDL computation for the Weibull model failed because the lower limit included zero. Of the models with adequate and viable fits, the BMDLs were not sufficiently close (differed by > 3-fold); therefore, EPA chose the model with the lowest BMDL. NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |
| Gamma | 0.9999 | 10.74 | 77.1 | 17.4 | |
| Log-Logistic | 1.000 | 10.73 | 93.8 | 24.9 | |
| Multistage 3 | 0.9999 | 6.900 | 65.1 | 13.7 | |
| Multistage 2 | 0.9876 | 9.349 | 47.3 | 12.6 | |
| Multistage 1 | 0.7953 | 11.76 | 16.1 | 8.33 | |
| Weibull | 0.9560 | 11.30 | 50.2 | 0 | |
| Logistic | 0.9999 | 10.74 | 97.0 | 36.3 | |
| Log-Probit | 1.000 | 10.73 | 103 | 24.5 | |
| Probit | 0.7909 | 12.33 | 55.2 | 27.5 | |
| Quantal Linear | 0.7953 | 11.76 | 16.1 | 8.33 | |

^a Selected model in bold.

A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-5. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-6.




Figure 1-5. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 16.14009333 | | | | |
| BMDL | 8.334284842 | | | | |
| BMDU | 32.92192881 | | | | |
| AIC | 11.757832 | | | | |
| P-value | 0.795259973 | | | | |
| D.O.F. | 4 | | | | |
| Chi ² | 1.674965599 | | | | |
| Slope Factor | 0.01199863 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | 0.006527875 | 3.737480823 | -7.3188 | 7.33185574 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-08 | 0 | 5 | -0.000276 |
| 35.3 | 0.205811145 | 1.029055727 | 0 | 5 | -1.014424 |
| 132.5 | 0.578924623 | 2.894623116 | 3 | 5 | 0.0619369 |
| 410 | 0.931191546 | 4.655957729 | 5 | 5 | 0.1594437 |
| 1368.7 | 0.999868263 | 4.999341314 | 5 | 5 | 0.0002946 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -3.365058335 | 5 | - | - | NA |
| Fitted Model | -4.878915999 | 1 | 3.02771533 | 4 | 0.5531981 |
| Reduced Model | -17.30867418 | 1 | 27.8872317 | 4 | <0.0001 |

Figure 1-6. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2.2 Degeneration with Necrosis of the Olfactory Mucosa in Female F344 Rats – 4-Hour Inhalation Exposure

Increased incidence of degeneration with necrosis of the olfactory mucosa was observed in female rats exposed to 1,2-dichloroethane by inhalation for four hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-5. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA’s *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-5. Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 5 | 0 |
| 35.3 | 5 | 0 |
| 132.5 | 5 | 4 |
| 410.0 | 5 | 5 |
| 1368.7 | 5 | 5 |

The BMD modeling results for increased incidence of degeneration with necrosis of the olfactory mucosa in female rats are summarized in Table 1-6. All models provided adequate fit to the data (chi-square p-value > 0.1). The BMDL computation for the Weibull model failed because the lower limit included zero. The BMDLs were not sufficiently close (differed by > 3-fold). Therefore, EPA chose the model with the lowest BMDL (Multistage 1-degree model).

Table 1-6. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-----------------|--------------|--------------------------------|---------------------------------|---|
| | p-value | AIC | | | |
| Dichotomous Hill | 1.000 | 9.008 | 80.1 | 22.5 | All models provided adequate fit to the data (chi-square p-value > 0.1); however, the BMDL computation for the Weibull model failed because the lower limit included zero. Of the viable models, the BMDLs were not sufficiently close (differed by > 3-fold); therefore, EPA chose the model with the lowest BMDL. |
| Gamma | 0.9975 | 9.087 | 60.3 | 17.5 | |
| Log-Logistic | 1.000 | 9.008 | 80.1 | 22.5 | |
| Multistage 3 | 0.9995 | 5.301 | 54.2 | 12.7 | |
| Multistage 2 | 0.9645 | 8.058 | 36.7 | 10.2 | |
| Multistage 1 | 0.6890 | 10.70 | 12.2 | 6.20 | |
| Weibull | 0.8634 | 10.24 | 35.9 | 0 | |
| Logistic | 0.9994 | 9.039 | 83.8 | 30.2 | |
| Log-Probit | 1.000 | 9.004 | 98.0 | 22.7 | |
| Probit | 0.5357 | 12.24 | 41.4 | 21.1 | |
| Quantal Linear | 0.6890 | 10.70 | 12.2 | 6.20 | NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |

^a Selected model in bold.

This data set is not well suited for BMD modeling; there is a single datapoint (at 80% incidence) with incidence between 0 and 100%; there are no data to inform the shape of the curve at the region of interest (~ 10%). As a result, the different models provide a broad range of BMD and BMDL estimates. Selection of the low end of the range to represent the BMDL is not among the more realistic possibilities

and in this case involves extrapolation below the range of observation to generate a BMD estimate well below the lowest tested dose.

A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-7. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-8.




Figure 1-7. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for Four Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 12.20199573 | | | | |
| BMDL | 6.197861053 | | | | |
| BMDU | 25.08064423 | | | | |
| AIC | 10.69847938 | | | | |
| P-value | 0.689023139 | | | | |
| D.O.F. | 4 | | | | |
| Chi ² | 2.254728302 | | | | |
| Slope Factor | 0.016134599 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | 0.008634695 | 5.028308495 | -9.8466689 | 9.86393833 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-08 | 0 | 5 | -0.000276 |
| 35.3 | 0.262732682 | 1.313663408 | 0 | 5 | -1.334841 |
| 132.5 | 0.68148862 | 3.407443098 | 4 | 5 | 0.5687917 |
| 410 | 0.970993197 | 4.854965983 | 5 | 5 | 0.3864798 |
| 1368.7 | 0.999992632 | 4.999963158 | 5 | 5 | 0.0060698 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -2.502012118 | 5 | - | - | NA |
| Fitted Model | -4.349239691 | 1 | 3.69445515 | 4 | 0.4489329 |
| Reduced Model | -17.14824501 | 1 | 29.2924658 | 4 | <0.0001 |

Figure 1-8. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2.3 Degeneration with Necrosis of the Olfactory Mucosa in Male and Female F344 Rats (Combined) – 4-Hour Inhalation Exposure

Increased incidence of degeneration with necrosis of the olfactory mucosa was observed in male and female rats (combined) exposed to 1,2-dichloroethane by inhalation for four hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-7. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA’s *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-7. Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Concentration (mg/m ³) | Number of Animals | Incidence |
|------------------------------------|-------------------|-----------|
| 0 | 10 | 0 |
| 35.3 | 10 | 0 |
| 132.5 | 10 | 7 |
| 410.0 | 10 | 10 |
| 1368.7 | 10 | 10 |

The BMD modeling results for increased incidence of degeneration with necrosis of the olfactory mucosa in male and female rats (combined) are summarized in Table 1-8. All models provided adequate fit to the data (chi-square p-value > 0.1). The BMDL computation for the Weibull model failed because the lower limit included zero. The BMDLs were not sufficiently close (differed by > 3-fold). Therefore, EPA chose the model with the lowest BMDL (Multistage 1-degree model).

This data set is not well suited for BMD modeling; there is a single datapoint (at 70% incidence) with incidence between 0 and 100%; there are no data to inform the shape of the curve at the region of interest (~ 10%). As a result, the different models provide a broad range of BMD and BMDL estimates. Selection of the low end of the range to represent the BMDL is not among the more realistic possibilities and in this case involves extrapolation below the range of observation to generate a BMD estimate well below the lowest tested dose.

Table 1-8. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-----------------|--------------|--------------------------------------|---------------------------------------|---|
| | p-value | AIC | | | |
| Dichotomous Hill | 1.000 | 16.22 | 87.6 | 34.9 | All models provided adequate fit to the data (chi-square p-value > 0.1); however, the BMDL computation for the Weibull model failed because the lower limit included zero. Of the viable models, The BMDLs were not sufficiently close (differed by > 3-fold); therefore, EPA chose the model with the lowest BMDL. |
| Gamma | 0.9987 | 16.27 | 68.9 | 30.9 | |
| Log-Logistic | 1.000 | 16.22 | 87.6 | 34.9 | |
| Multistage 3 | 0.9988 | 12.66 | 59.5 | 24.6 | |
| Multistage 2 | 0.9280 | 15.82 | 41.8 | 18.8 | |
| Multistage 1 | 0.4440 | 20.68 | 14.1 | 8.76 | |
| Weibull | 0.8079 | 17.90 | 42.7 | 0 | |
| Logistic | 0.9992 | 16.26 | 91.0 | 44.7 | |
| Log-Probit | 1.000 | 16.22 | 95.1 | 33.7 | |
| Probit | 0.3831 | 20.81 | 47.8 | 29.2 | |
| Quantal Linear | 0.4440 | 20.68 | 14.1 | 8.76 | NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |

^a Selected model in bold.

A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-9. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-10.




Figure 1-9. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 14.10695256 | | | | |
| BMDL | 8.757017138 | | | | |
| BMDU | 23.27887944 | | | | |
| AIC | 20.67643472 | | | | |
| P-value | 0.444048554 | | | | |
| D.O.F. | 4 | | | | |
| Chi ² | 3.72812481 | | | | |
| Slope Factor | 0.011419414 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | 0.007468694 | 3.038026138 | -5.9469532 | 5.96189056 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 35.3 | 0.231753727 | 2.317537268 | 0 | 10 | -1.736853 |
| 132.5 | 0.628275403 | 6.282754027 | 7 | 10 | 0.4693345 |
| 410 | 0.953213694 | 9.532136945 | 10 | 10 | 0.7005905 |
| 1368.7 | 0.999963653 | 9.999636531 | 10 | 10 | 0.0190652 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -6.108643021 | 5 | - | - | NA |
| Fitted Model | -9.338217359 | 1 | 6.45914868 | 4 | 0.1673826 |
| Reduced Model | -34.49718792 | 1 | 56.7770898 | 4 | <0.0001 |

Figure 1-10. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2.4 Degeneration with Necrosis of the Olfactory Mucosa in Male F344 Rats – 8-Hour Inhalation Exposure

Increased incidence of degeneration with necrosis of the olfactory mucosa was observed in male rats exposed to 1,2-dichloroethane by inhalation for eight hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-9. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA’s *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-9. Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 5 | 0 |
| 71.3 | 5 | 0 |
| 145.0 | 5 | 1 |
| 210.2 | 5 | 4 |

The BMD modeling results for increased incidence of degeneration with necrosis of the olfactory mucosa in male rats are summarized in Table 1-10. All models provided adequate fit to the data (chi-square p-value > 0.1). The BMDLs were not sufficiently close (differed by > 3-fold). Therefore, EPA chose the model with the lowest BMDL (Multistage 1-degree model).

This data set is not well suited for BMD modeling; there is a single datapoint (at 20% incidence) with incidence between 0 and 80%. As a result, the different models provide a broad range of BMD and BMDL estimates. Selection of the low end of the range to represent the BMDL is not among the more realistic possibilities and, in this case, involves extrapolation below the range of observation to generate a BMD estimate well below the lowest tested dose.

Table 1-10. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-----------------|--------------|--------------------------------|---------------------------------|--|
| | p-value | AIC | | | |
| Dichotomous Hill | 1.000 | 14.01 | 138 | 70.1 | All models provided adequate fit to the data (chi-square p-value > 0.1). The BMDLs were not sufficiently close (differed by > 3-fold); therefore, EPA chose the model with the lowest BMDL. NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |
| Gamma | 0.9992 | 12.03 | 127 | 67.7 | |
| Log-Logistic | 0.9969 | 14.02 | 131 | 69.9 | |
| Multistage 3 | 0.9325 | 11.13 | 94.3 | 37.2 | |
| Multistage 2 | 0.5937 | 14.54 | 70.9 | 26.8 | |
| Multistage 1 | 0.2777 | 17.03 | 33.5 | 17.1 | |
| Weibull | 0.9989 | 12.05 | 128 | 63.8 | |
| Logistic | 0.9729 | 14.10 | 130 | 71.5 | |
| Log-Probit | 0.9999 | 14.01 | 132 | 71.7 | |
| Probit | 0.9926 | 14.03 | 130 | 68.3 | |
| Quantal Linear | 0.2777 | 17.03 | 33.5 | 17.1 | |

^a Selected model in bold.

A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-11. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-12.




Figure 1-11. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 8 Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 33.46390531 | | | | |
| BMDL | 17.11048878 | | | | |
| BMDU | 78.84479045 | | | | |
| AIC | 17.03210085 | | | | |
| P-value | 0.277708981 | | | | |
| D.O.F. | 3 | | | | |
| Chi ² | 3.853633558 | | | | |
| Slope Factor | 0.005844368 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | 0.003148482 | 0.300815071 | -0.5864382 | 0.59273519 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-08 | 0 | 5 | -0.000276 |
| 71.3 | 0.20107387 | 1.005369348 | 0 | 5 | -1.121785 |
| 145 | 0.366521946 | 1.832609729 | 1 | 5 | -0.772753 |
| 210.2 | 0.484083813 | 2.420419067 | 4 | 5 | 1.4135365 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -5.004024235 | 4 | - | - | NA |
| Fitted Model | -7.516050426 | 1 | 5.02405238 | 3 | 0.1700444 |
| Reduced Model | -11.24670289 | 1 | 12.4853573 | 3 | 0.0585611 |

Figure 1-12. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male Rats Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2.5 Degeneration with Necrosis of the Olfactory Mucosa in Female F344 Rats – 8-Hour Inhalation Exposure

Increased incidence of degeneration with necrosis of the olfactory mucosa was observed in female rats exposed to 1,2-dichloroethane by inhalation for eight hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-11. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-11. Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 5 | 0 |
| 71.3 | 5 | 0 |
| 145.0 | 5 | 3 |
| 210.2 | 5 | 5 |

The BMD modeling results for increased incidence of degeneration with necrosis of the olfactory mucosa in male rats are summarized in Table 1-12. All models provided adequate fit to the data (chi-square p-value > 0.1). The BMDL computation for the Weibull model failed because the lower limit included zero. The BMDLs were not sufficiently close (differed by > 3-fold). Therefore, EPA chose the model with the lowest BMDL (Multistage 1-degree model).

This data set is not well suited for BMD modeling; there is a single datapoint (at 60% incidence) with incidence between 0 and 100%; there are no data to inform the shape of the curve at the region of interest (~ 10%). As a result, the different models provide a broad range of BMD and BMDL estimates. Selection of the low end of the range to represent the BMDL is not among the more realistic possibilities and in this case involves extrapolation below the range of observation to generate a BMD estimate well below the lowest tested dose.

Table 1-12. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-----------------|--------------|--------------------------------|---------------------------------|---|
| | p-value | AIC | | | |
| Dichotomous Hill | 0.9999 | 8.738 | 125 | 64.6 | All models provided adequate fit to the data (chi-square p-value > 0.1); however, the BMDL computation for the Weibull model failed because the lower limit included zero. Of the viable models, the BMDLs were not sufficiently close (differed by > 3-fold); therefore, EPA chose the model with the lowest BMDL. |
| Gamma | 0.9835 | 9.036 | 97.6 | 57.9 | |
| Log-Logistic | 0.9999 | 8.738 | 125 | 64.7 | |
| Multistage 3 | 0.9249 | 8.426 | 68.9 | 26.1 | |
| Multistage 2 | 0.5547 | 12.47 | 47.1 | 18.6 | |
| Multistage 1 | 0.1987 | 16.31 | 16.3 | 9.20 | |
| Weibull | 0.9997 | 10.73 | 122 | 0 | |
| Logistic | 1.000 | 8.733 | 124 | 58.8 | |
| Log-Probit | 1.000 | 10.73 | 132 | 64.5 | |
| Probit | 1.000 | 10.73 | 126 | 55.0 | |
| Quantal Linear | 0.1987 | 16.31 | 16.3 | 9.20 | NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |

^a Selected model in bold.

A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-13. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-14.




Figure 1-13. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 8 Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 16.34257648 | | | | |
| BMDL | 9.199716086 | | | | |
| BMDU | 32.02480499 | | | | |
| AIC | 16.31075155 | | | | |
| P-value | 0.198741114 | | | | |
| D.O.F. | 3 | | | | |
| Chi ² | 4.656587683 | | | | |
| Slope Factor | 0.010869901 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | 0.006446995 | 0.508897083 | -0.990973 | 1.00386696 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-08 | 0 | 5 | -0.000276 |
| 71.3 | 0.368508494 | 1.842542471 | 0 | 5 | -1.708146 |
| 145 | 0.607341244 | 3.03670622 | 3 | 5 | -0.033615 |
| 210.2 | 0.742093574 | 3.710467871 | 5 | 5 | 1.3182165 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -3.365058335 | 4 | - | - | NA |
| Fitted Model | -7.155375774 | 1 | 7.58063488 | 3 | 0.0555224 |
| Reduced Model | -13.46023334 | 1 | 20.19035 | 3 | 0.000155 |

Figure 1-14. Details Regarding the Selected Model (Multistage 1-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Female Rats Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2.6 Degeneration with Necrosis of the Olfactory Mucosa in Male and Female F344 Rats (Combined) – 8-Hour Inhalation Exposure

Increased incidence of degeneration with necrosis of the olfactory mucosa was observed in male and female rats (combined) exposed to 1,2-dichloroethane by inhalation for eight hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-13. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-13. Incidence of Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (combined) and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 10 | 0 |
| 71.3 | 10 | 0 |
| 145.0 | 10 | 4 |
| 210.2 | 10 | 9 |

The BMD modeling results for increased incidence of degeneration with necrosis of the olfactory mucosa in male rats are summarized in Table 1-14. All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Multistage 1-degree/Quantal Linear model. The BMDLs of the fit models were sufficiently close (differed by < 3-fold). Therefore, EPA chose the model with the lowest AIC (Multistage 3-degree model).

Table 1-14. BMD Modeling Results for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-----------------|--------------|--------------------------------|---------------------------------|--|
| | p-value | AIC | | | |
| Dichotomous Hill | 1.000 | 23.96 | 131 | 78.1 | All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Multistage 1-degree/Quantal Linear model. The BMDLs of the fit models were sufficiently close (differed by < 3-fold); therefore, EPA chose the model with the lowest AIC. |
| Gamma | 0.9847 | 24.01 | 112 | 75.2 | |
| Log-Logistic | 0.9779 | 24.04 | 114 | 77.5 | |
| Multistage 3 | 0.8911 | 21.80 | 81.4 | 48.9 | |
| Multistage 2 | 0.3612 | 26.88 | 57.8 | 34.3 | |
| Multistage 1 | 0.0570 | 32.87 | 23.1 | 14.8 | |
| Weibull | 0.9664 | 22.40 | 106 | 68.2 | |
| Logistic | 0.8515 | 24.46 | 110 | 72.6 | |
| Log-Probit | 0.9965 | 23.97 | 114 | 77.8 | |
| Probit | 0.9049 | 24.26 | 110 | 70.5 | |
| Quantal Linear | 0.0570 | 32.87 | 23.1 | 14.8 | |

^a Selected model in bold.

A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-15. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-16.




Figure 1-15. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Exposed to 1,2-Dichloroethane Via Inhalation for 8 Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 81.37548237 | | | | |
| BMDL | 48.94695881 | | | | |
| BMDU | 97.14021529 | | | | |
| AIC | 21.80354181 | | | | |
| P-value | 0.891072257 | | | | |
| D.O.F. | 4 | | | | |
| Chi ² | 1.120086799 | | | | |
| Slope Factor | 0.002043028 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | Bounded | NA | NA | NA | |
| b2 | Bounded | NA | NA | NA | |
| b3 | Bounded | NA | NA | NA | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 71.3 | 0.068417522 | 0.68417522 | 0 | 10 | -0.856985 |
| 145 | 0.449030218 | 4.490302179 | 4 | 10 | -0.311718 |
| 210.2 | 0.837310844 | 8.373108437 | 9 | 10 | 0.5371181 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -9.980946404 | 4 | - | - | NA |
| Fitted Model | -10.90177091 | 0 | 1.84164901 | 4 | 0.7648543 |
| Reduced Model | -25.22324114 | 1 | 30.4845895 | 3 | <0.0001 |

Figure 1-16. Details Regarding the Selected Model (Multistage 3-Degree) for Degeneration with Necrosis of the Olfactory Mucosa in Male and Female Rats (Combined) Following an 8-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2.7 Regeneration of the Olfactory Mucosa in Male F344 Rats – 4-Hour Inhalation Exposure

Increased incidence of regeneration of the olfactory mucosa was observed in male rats exposed to 1,2-dichloroethane by inhalation for four hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-15. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-15. Incidence of Regeneration of the Olfactory Mucosa in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 5 | 0 |
| 132.5 | 5 | 4 |
| 410.0 | 5 | 5 |
| 1368.7 | 5 | 5 |

The BMD modeling results for increased incidence of regeneration of the olfactory mucosa in male rats are summarized in Table 1-16. All models provided adequate fit to the data (chi-square p-value > 0.1). The BMDL computations failed for the Dichotomous Hill and Log-Probit models because the lower limit included zero. The Gamma, Log-Logistic, Multistage 3-, 2-, and 1-degree, Weibull, and Quantal Linear models were questionable because the BMDL values were 10 times lower than the lowest non-zero concentration. The BMDLs of the remaining viable models (Logistic and Probit) were sufficiently close (differed by < 3-fold). Therefore, EPA chose the model with the lower AIC (Logistic).

This data set is not well suited for BMD modeling. There is a single datapoint (at 80% incidence) with incidence between 0 and 100%; there are no data to inform the shape of the curve at the region of interest (~ 10%). As a result, the different models provide a broad range of BMD and BMDL estimates, all of which in this case involve extrapolation below the range of observation to generate BMD estimates well below the lowest tested dose.

Table 1-16. BMD Modeling Results for Regeneration of the Olfactory Mucosa in Male Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|------------------|-----------------|--------------|--------------------------------------|---------------------------------------|---|
| | p-value | AIC | | | |
| Dichotomous Hill | 0.9998 | 9.005 | 80.07 | 0 | <p>All models provided adequate fit to the data (chi-square p-value > 0.1); however, the BMDL computations failed for the Dichotomous Hill and Log-Probit models because the lower limit included zero. The Gamma, Log-Logistic, Multistage 3-, 2-, and 1-degree, Weibull, and Quantal Linear models were questionable because the BMDL values were 10 times lower than the lowest non-zero concentration. The BMDLs of the remaining viable models (Logistic and Probit) were sufficiently close (differed by < 3-fold); therefore, EPA chose the model with the lower AIC.</p> <p>NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution.</p> |
| Gamma | 1.000 | 9.004 | 58.72 | 3.712 | |
| Log-Logistic | 0.9998 | 9.005 | 80.07 | 0.3723 | |
| Multistage 3 | 1.000 | 7.004 | 17.27 | 3.712 | |
| Multistage 2 | 1.000 | 7.004 | 33.90 | 3.712 | |
| Multistage 1 | 0.9983 | 7.065 | 8.302 | 3.679 | |
| Weibull | 1.000 | 9.004 | 28.36 | 3.712 | |
| Logistic | 0.9994 | 9.007 | 83.51 | 16.06 | |
| Log-Probit | 1.000 | 9.004 | 81.52 | 0 | |
| Probit | 0.6397 | 10.43 | 30.47 | 14.20 | |
| Quantal Linear | 0.9983 | 7.065 | 8.302 | 3.679 | |

^a Selected model in bold.

A plot of the Logistic model with a BMR of 10% ER is shown in Figure 1-17. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-18.




Figure 1-17. Plot of Response by Concentration with Fitted Curve for the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 83.50781438 | | | | |
| BMDL | 16.05730006 | | | | |
| BMDU | 105.6670162 | | | | |
| AIC | 9.006521039 | | | | |
| P-value | 0.999376363 | | | | |
| D.O.F. | 2 | | | | |
| Chi ² | 0.001247664 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| a | -8.295982925 | 1.11664204 | -10.484561 | -6.1074047 | |
| b | 0.07306203 | 3.16E-07 | 0.07306141 | 0.07306265 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.000249455 | 0.001247275 | 0 | 5 | -0.035321 |
| 132.5 | 0.799750546 | 3.998752732 | 4 | 5 | 0.0013938 |
| 410 | 1 | 4.999999998 | 5 | 5 | 4.428E-05 |
| 1368.7 | 1 | 5 | 5 | 5 | 0 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -2.502012118 | 4 | - | - | NA |
| Fitted Model | -2.503260519 | 2 | 0.0024968 | 2 | 0.9987524 |
| Reduced Model | -12.21728604 | 1 | 19.4305478 | 3 | 0.0002227 |

Figure 1-18. Details Regarding the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Male Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2.8 Regeneration of the Olfactory Mucosa in Female F344 Rats – 4-Hour Inhalation Exposure

Increased incidence of regeneration of the olfactory mucosa was observed in female rats exposed to 1,2-dichloroethane by inhalation for four hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-17. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA’s *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-17. Incidence of Regeneration of the Olfactory Mucosa in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 5 | 0 |
| 132.5 | 5 | 4 |
| 410.0 | 5 | 5 |
| 1368.7 | 5 | 5 |

The BMD modeling results for increased incidence of regeneration of the olfactory mucosa in female rats are summarized in Table 1-18. All models provided adequate fit to the data (chi-square p-value > 0.1). The BMDL computations failed for the Dichotomous Hill and Log-Probit models because the lower limit included zero. The Gamma, Log-Logistic, Multistage 3-, 2-, and 1-degree, Weibull, and Quantal Linear models were considered questionable because the BMDL values were 10 times lower than the lowest non-zero concentration. The BMDLs of the remaining models (Logistic and Probit) were sufficiently close (differed by < 3-fold). Therefore, EPA chose the model with the lower AIC (Logistic).

This data set is not well suited for BMD modeling; there is a single datapoint (at 80% incidence) with incidence between 0 and 100%; there are no data to inform the shape of the curve at the region of interest (~ 10%). As a result, the different models provide a broad range of BMD and BMDL estimates, all of which in this case involve extrapolation below the range of observation to generate BMD estimates well below the lowest tested dose.

Table 1-18. BMD Modeling Results for Regeneration of the Olfactory Mucosa in Female Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|------------------|-----------------|--------------|--------------------------------|---------------------------------|---|
| | p-value | AIC | | | |
| Dichotomous Hill | 0.9998 | 9.005 | 80.07 | 0 | All models provided adequate fit to the data (chi-square p-value > 0.1); however, the BMDL computations failed for the Dichotomous Hill and Log-Probit models because the lower limit included zero. The Gamma, Log-Logistic, Multistage 3-, 2-, and 1-degree, Weibull, and Quantal Linear models were considered questionable because the BMDL values were 10 times lower than the lowest non-zero concentration. The BMDLs of the remaining viable models (Logistic and Probit) were sufficiently |
| Gamma | 1.000 | 9.004 | 58.72 | 3.712 | |
| Log-Logistic | 0.9998 | 9.005 | 80.07 | 0.3723 | |
| Multistage 3 | 1.000 | 7.004 | 17.27 | 3.712 | |
| Multistage 2 | 1.000 | 7.004 | 33.90 | 3.712 | |
| Multistage 1 | 0.9983 | 7.065 | 8.302 | 3.679 | |
| Weibull | 1.000 | 9.004 | 28.36 | 3.712 | |
| Logistic | 0.9994 | 9.007 | 83.51 | 16.06 | |
| Log-Probit | 1.000 | 9.004 | 81.52 | 0 | |
| Probit | 0.6397 | 10.43 | 30.47 | 14.20 | |
| Quantal Linear | 0.9983 | 7.065 | 8.302 | 3.679 | |

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|--------------------------------------|-----------------|-----|--------------------------------------|---------------------------------------|---|
| | p-value | AIC | | | |
| | | | | | close (differed by < 3-fold); therefore, EPA chose the model with the lower AIC. NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |
| ^a Selected model in bold. | | | | | |

A plot of the Logistic model with a BMR of 10% ER is shown in Figure 1-19. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-20.




Figure 1-19. Plot of Response by Concentration with Fitted Curve for the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 83.50781438 | | | | |
| BMDL | 16.05730006 | | | | |
| BMDU | 105.6670162 | | | | |
| AIC | 9.006521039 | | | | |
| P-value | 0.99937543 | | | | |
| D.O.F. | 2 | | | | |
| Chi ² | 0.00124953 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| a | -8.295982925 | 1.11664204 | -10.484561 | -6.1074047 | |
| b | 0.07306203 | 3.16E-07 | 0.07306141 | 0.07306265 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.000249455 | 0.001247275 | 0 | 5 | -0.035321 |
| 132.5 | 0.799750546 | 3.998752732 | 4 | 5 | 0.0013938 |
| 410 | 1 | 4.999999998 | 5 | 5 | 4.428E-05 |
| 1368.7 | 1 | 5 | 5 | 5 | 0 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -2.502012118 | 4 | - | - | NA |
| Fitted Model | -2.503260519 | 2 | 0.0024968 | 2 | 0.9987524 |
| Reduced Model | -12.21728604 | 1 | 19.428051 | 3 | 0.000223 |

Figure 1-20. Details Regarding the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Female Rats Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.2.9 Regeneration of the Olfactory Mucosa in Male and Female F344 Rats (Combined) – 4-Hour Inhalation Exposure

Increased incidence of regeneration of the olfactory mucosa was observed in male and female rats (combined) exposed to 1,2-dichloroethane by inhalation for four hours ([Dow Chemical, 2006](#)). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-19. Dichotomous models were fit to the incidence data.

A BMR of 10% ER was chosen according to EPA’s *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-19. Incidence of Regeneration of the Olfactory Mucosa in Male and Female Rats (Combined) and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 10 | 0 |
| 132.5 | 10 | 8 |
| 410.0 | 10 | 10 |
| 1368.7 | 10 | 10 |

The BMD modeling results for increased incidence of regeneration of the olfactory mucosa in male rats are summarized in Table 1-20. All models provided adequate fit to the data (chi-square p-value > 0.1). The Dichotomous Hill, Gamma, Log-Logistic, Multistage 3-, 2-, and 1-degree, Weibull, Log-Probit, and Quantal Linear models were considered questionable because the BMDL values were 10 times lower than the lowest non-zero concentration. The BMDLs of the remaining models (Logistic and Probit) were sufficiently close (differed by < 3-fold). Therefore, EPA chose the model with the lower AIC (Logistic).

This data set is not well suited for BMD modeling; there is a single datapoint (at 80% incidence) with incidence between 0 and 100%; there are no data to inform the shape of the curve at the region of interest (~ 10%). As a result, the different models provide a broad range of BMD and BMDL estimates, all of which in this case involve extrapolation below the range of observation to generate BMD estimates well below the lowest tested dose.

Table 1-20. BMD Modeling Results for Regeneration of the Olfactory Mucosa in Male and Female Rats (Combined) Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|------------------|-----------------|--------------|--------------------------------|---------------------------------|---|
| | p-value | AIC | | | |
| Dichotomous Hill | 0.9996 | 14.01 | 80.07 | 0.9633 | All models provided adequate fit to the data (chi-square p-value > 0.1); however, the Dichotomous Hill, Gamma, Log-Logistic, Multistage 3-, 2-, and 1-degree, Weibull, Log-Probit, and Quantal Linear models were considered questionable because the BMDL values were 10 times lower than the lowest non-zero concentration. The BMDLs of the remaining viable models (Logistic and Probit) were sufficiently close (differed by < 3-fold); therefore, EPA chose the model with the lower AIC. |
| Gamma | 1.000 | 14.01 | 58.72 | 4.706 | |
| Log-Logistic | 0.9996 | 14.01 | 80.07 | 0.9651 | |
| Multistage 3 | 1.000 | 12.01 | 17.27 | 4.706 | |
| Multistage 2 | 1.000 | 12.01 | 33.90 | 4.706 | |
| Multistage 1 | 0.9954 | 12.13 | 8.302 | 4.645 | |
| Weibull | 1.000 | 14.01 | 28.36 | 4.706 | |
| Logistic | 0.9988 | 14.01 | 83.51 | 23.18 | |
| Log-Probit | 1.000 | 14.01 | 81.52 | 0.005232 | |
| Probit | 0.4092 | 16.86 | 30.47 | 17.47 | |
| Quantal Linear | 0.995 | 12.13 | 8.302 | 4.645 | |

| Model | Goodness of Fit | | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|--------------------------------------|-----------------|-----|--------------------------------------|---------------------------------------|---|
| | p-value | AIC | | | |
| | | | | | NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |
| ^a Selected model in bold. | | | | | |

A plot of the Logistic model with a BMR of 10% ER is shown in Figure 1-21. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-22.




Figure 1-21. Plot of Response by Concentration with Fitted Curve for the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Male and Female Rats (Combined) Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 83.50781438 | | | | |
| BMDL | 23.18348326 | | | | |
| BMDU | 99.55849153 | | | | |
| AIC | 14.01304208 | | | | |
| P-value | 0.99875125 | | | | |
| D.O.F. | 2 | | | | |
| Chi ² | 0.002499061 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| a | -8.295982925 | 0.789585159 | -9.8435414 | -6.7484244 | |
| b | 0.07306203 | 3.16E-07 | 0.07306141 | 0.07306265 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.000249455 | 0.002494549 | 0 | 10 | -0.049952 |
| 132.5 | 0.799750546 | 7.997505465 | 8 | 10 | 0.0019712 |
| 410 | 1 | 9.999999996 | 10 | 10 | 6.262E-05 |
| 1368.7 | 1 | 10 | 10 | 10 | 0 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -5.004024235 | 4 | - | - | NA |
| Fitted Model | -5.006521039 | 2 | 0.00499361 | 2 | 0.9975063 |
| Reduced Model | -24.43457208 | 1 | 38.8561021 | 3 | <0.0001 |

Figure 1-22. Details Regarding the Selected Model (Logistic) for Regeneration of the Olfactory Mucosa in Male and Female Rats (Combined) Following a 4-Hour Inhalation Exposure to 1,2-Dichloroethane

1.1.1.1.3 Hepatic Effects

Only one hepatic endpoint was identified for acute-duration inhalation exposure. EPA modeled serum L-iditol dehydrogenase levels in male mice exposed to 1,2-dichloroethane by inhalation for four hours (Storer et al., 1984). The modeling results are not presented because neither the constant nor nonconstant variance models provided adequate fit to the variance data.

1.1.1.1.4 Renal Effects

For acute inhalation exposure, EPA selected two renal endpoints for quantitative dose-response analysis with BMDS, including relative kidney weights and blood urea nitrogen (BUN) levels in serum following four-hour inhalation exposure in male mice (Storer et al., 1984). For both data sets, only data for the control and two lowest concentrations (0, 107, and 337 mg/m³) were modeled; high mortality (four of five and five of five) precluded collection of relevant data at the two highest tested concentrations (723.2

and 1313 mg/m³, respectively). EPA did not present the BMD modeling results for increased BUN in male mice because no model resulted in an adequate fit.

1.1.1.1.4.1 Relative Kidney Weight (Kidney Weight/100 g Body Weight) in Male B6C3F1 Mice – 4-Hour Inhalation Exposure

Relative kidney weight was significantly increased in male mice exposed to 1,2-dichloroethane by inhalation for four hours (Storer et al., 1984). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-21. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected because EPA considers a 10% change in relative kidney weight to be biologically significant.

Table 1-21. Increased Relative Kidney Weight in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (Kidney Weight/100 g Body Weight) | SD (Kidney Weight/100 g Body Weight) |
|---|-------------------|--|--------------------------------------|
| 0 | 5 | 1.53 | 0.07 |
| 107 | 5 | 1.53 | 0.08 |
| 337 | 5 | 1.71 | 0.09 |

The BMD modeling results for increased relative kidney weight in male mice are summarized in Table 1-22. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, only the Polynomial 2-degree and Linear models provided adequate fit to the means. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Polynomial 2-degree model) was selected.

Table 1-22. Summary of BMD Modeling Results for Increased Relative Kidney Weight in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 10%RD (mg/m ³) | BMDL 10%RD (mg/m ³) | Basis for Model Selection |
|----------------------------|-------------------------|---------------|------------------------------|-------------------------------|--------------------------------|---------------------------------|---|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | NA | -28.40 | 319 | 107 | 334 | 213 | Only the Polynomial 2-degree and Linear models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | -26.40 | 205 | 160 | 221 | 113 | |
| Hill | NA | -26.40 | 299 | 106 | 330 | 318 | |
| Polynomial Degree 2 | 0.6765 | -30.22 | 209 | 102 | 303 | 207 | |
| Power | NA | -28.40 | 319 | 105 | 334 | 211 | |
| Linear | 0.1691 | -28.50 | 134 | 88.5 | 262 | 180 | |

^a Selected model in bold.

Plots of the Polynomial 2-degree model with BMRs of one SD and 10% RD are shown in Figure 1-23 and Figure 1-24, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-25 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-23. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 1SD (Constant Variance)




Figure 1-24. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours and BMR of 10%RD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 208.9772425 | | | | | | | |
| BMDL | 102.2542989 | | | | | | | |
| BMDU | 285.9155546 | | | | | | | |
| AIC | -30.22136297 | | | | | | | |
| Test 4 P-value | 0.676535642 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 1.520909738 | 2.41E-02 | 1.47361189 | 1.56820759 | | | | |
| beta | Bounded | NA | NA | NA | | | | |
| beta2 | 1.65792E-06 | 3.66E-07 | 9.4021E-07 | 2.3756E-06 | | | | |
| alpha | 0.005242334 | 1.00E-05 | 0.00522265 | 0.00526202 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 5 | 1.520909738 | 1.53 | 1.53 | 0.07240397 | 0.07 | 0.07 | 0.280736593 |
| 107 | 5 | 1.539891291 | 1.53 | 1.53 | 0.07240397 | 0.08 | 0.08 | -0.305474959 |
| 337 | 5 | 1.709198317 | 1.71 | 1.71 | 0.07240397 | 0.09 | 0.09 | 0.024758571 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | 18.1977074 | 4 | -28.395415 | | | | | |
| A2 | 18.35417009 | 6 | -24.70834 | | | | | |
| A3 | 18.1977074 | 4 | -28.395415 | | | | | |
| fitted | 18.11068148 | 3 | -30.221363 | | | | | |
| R | 11.65750945 | 2 | -19.315019 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test d.f. | p-value | | | | | |
| 1 | 13.39332127 | 4 | 0.0095056 | | | | | |
| 2 | 0.312925375 | 2 | 0.85516343 | | | | | |
| 3 | 0.312925375 | 2 | 0.85516343 | | | | | |
| 4 | 0.174051828 | 1 | 0.67653564 | | | | | |

Figure 1-25. Details Regarding the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Hours

1.1.1.2 Short-term/Intermediate

1.1.1.2.1 Mortality

Two short-term/intermediate-duration inhalation studies were identified that showed potentially treatment-related incidence of mortality in exposed animals (Igwe et al., 1986; Rao et al., 1980). Though neither data set showed a statistically significant increase, these data were modeled.

1.1.1.2.1.1 Mortality in Male Rats – 30-Day Inhalation Exposure

There was mortality in male rats exposed to 1,2-dichloroethane by inhalation for 30 days (seven hours per day, 5 days per week) (Igwe et al., 1986). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day and 7 days per week. The concentration and response data used for the modeling are presented in Table 1-23. Dichotomous models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). BMRs of 1 and 5% ER were also selected due to severity of the endpoint.

Table 1-23. Incidence of Mortality in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 30 | 0 |
| 129 | 12 | 0 |
| 256 | 12 | 1 |
| 383 | 12 | 2 |

The BMD modeling results for incidence of mortality are summarized in Table 1-24. All models provided adequate fit to the data (chi-square p-value > 0.1). The Dichotomous Hill and Log-Probit models were considered questionable/unusable at all attempted BMRs because the BMD/BMDL ratio was > 20 and the BMDL was > 10 times lower than the lowest non-zero concentration or the BMDL computation failed. At a BMR of 10% ER or 5% ER, the BMDLs of the remaining fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree). At a BMR of one percent ER, there is much greater uncertainty because the BMR is well below anything observable in the data. As a result, there is greater spread in the BMDLs (> 3-fold) and additional models provided questionable results (Multistage 1-degree and Quantal Linear models) due to BMDL > 10 times lower than the lowest non-zero concentration. Because the BMDLs of remaining viable models differed by > 3-fold, the BMDS recommended the model with the lowest BMDL (Log-Logistic); however, the Multistage 3-degree model was a viable alternative and was selected to be consistent with model selection at 10% ER and 5% ER. There was little difference in BMD/BMDL values between the Multistage 3-degree and Log-Logistic models.

Table 1-24. Summary of BMD Modeling Results for Incidence of Mortality in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane^a

| Model | Goodness of Fit (Means) | | BMD 1%ER (mg/m ³) | BMDL 1%ER (mg/m ³) | BMD 5%ER (mg/m ³) | BMDL 5%ER (mg/m ³) | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|--------------|-------------------------------|--------------------------------|-------------------------------|--------------------------------|--------------------------------|---------------------------------|--|
| | p-value | AIC | | | | | | | |
| Dichotomous Hill | 1.000 | 21.70 | 220 | 0.0383 | 244 | 0.0428 | 262 | 1.42 | All models provided adequate fit to the data (chi-square p-value > 0.1). The Dichotomous Hill and Log-Probit models were considered questionable/unusable, however, at all attempted BMRs because the BMD/BMDL ratio was > 20 and the BMDL was > 10 times lower than the lowest non-zero concentration or the BMDL computation failed. At BMR = 10% ER or 5% ER, the BMDLs of the fit models were sufficiently close (differed by < 3-fold); therefore, EPA chose the model with the lowest AIC (Multistage 3-degree). At BMR = one percent ER, additional models provided questionable results (Multistage 1-degree and Quantal Linear models) due to BMDL > 10 times lower than the lowest non-zero concentration. The BMDLs of remaining viable models differed by > 3-fold and the BMDS recommended the model with the lowest BMDL (Log-Logistic); however, the Multistage 3-degree model was a viable alternative and was selected to be consistent with model selection at 10% ER and 5% ER. |
| Gamma | 0.9055 | 31.97 | 145 | 14.9 | 239 | 75.8 | 304 | 156 | |
| Log-Logistic | 0.8976 | 22.00 | 138 | 13.7 | 238 | 71.2 | 305 | 150 | |
| Multistage 3 | 0.9929 | 18.02 | 141 | 14.7 | 243 | 75.1 | 308 | 154 | |
| Multistage 2 | 0.9676 | 20.19 | 92.7 | 14.3 | 209 | 73.1 | 300 | 150 | |
| Multistage 1 | 0.8614 | 20.95 | 29.1 | 12.8 | 149 | 65.4 | 305 | 134 | |
| Weibull | 0.9727 | 20.02 | 136 | 14.7 | 239 | 75.1 | 307 | 154 | |
| Logistic | 0.7460 | 22.41 | 143 | 52.2 | 265 | 170 | 325 | 243 | |
| Log-Probit | 0.9254 | 21.91 | 151 | 0 | 236 | 0 | 300 | 0.0316 | |
| Probit | 0.8109 | 22.21 | 143 | 47.7 | 255 | 158 | 316 | 230 | |
| Quantal Linear | 0.8614 | 20.95 | 29.1 | 12.8 | 149 | 65.4 | 305 | 134 | |

^a Selected model in bold.

Plots of the Multistage 3-degree model with BMRs of 10%, 5%, and 1% ER are shown in Figure 1-26, Figure 1-27, and Figure 1-28, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-29 (BMD and BMDL shown are for BMR of 10% ER; the rest is applicable to all BMRs).




Figure 1-26. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days (Seven Hours per Day, 5 days per Week) and BMR of 10%ER




Figure 1-27. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days (Seven Hours per Day, 5 days per Week) and BMR of 5%ER




Figure 1-28. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days (7n Hours per Day, 5 Days per Week) and BMR of 1%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 308.342534 | | | | |
| BMDL | 154.1950362 | | | | |
| BMDU | 738.5150207 | | | | |
| AIC | 18.0236136 | | | | |
| P-value | 0.992913128 | | | | |
| D.O.F. | 4 | | | | |
| Chi ² | 0.248106117 | | | | |
| Slope Factor | 0.000648529 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 4 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | Bounded | NA | NA | NA | |
| b2 | Bounded | NA | NA | NA | |
| b3 | Bounded | NA | NA | NA | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 4.56899E-07 | 0 | 30 | -0.000676 |
| 129 | 0.007685521 | 0.092226257 | 0 | 12 | -0.304862 |
| 256 | 0.058515379 | 0.702184543 | 1 | 12 | 0.3662815 |
| 383 | 0.18283766 | 2.194051923 | 2 | 12 | -0.144924 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -8.848766303 | 4 | - | - | NA |
| Fitted Model | -9.011806799 | 0 | 0.32608099 | 4 | 0.9880689 |
| Reduced Model | -12.20388835 | 1 | 6.71024408 | 3 | 0.0817297 |

Figure 1-29. Details Regarding the Selected Model (Multistage 3-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane 0Via Inhalation for 30 Days

1.1.1.2.1.2 Mortality in Female Rabbits – Inhalation Exposure on GD 6 to 18

There was mortality in female rabbits exposed to exposed to 1,2-dichloroethane by inhalation on GD 6 to 18 (seven hours per day) (Rao et al., 1980). The measured exposure concentrations (reported in units of ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day. The concentration and response data used for the modeling are presented in Table 1-25. Dichotomous models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012). BMRs of 1 and 5% ER were also selected due to severity of the endpoint.

Table 1-25. Incidence of Mortality in Female Rabbits and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Incidence |
|---|-------------------|-----------|
| 0 | 20 | 0 |
| 118 | 21 | 4 |
| 353 | 19 | 3 |

The BMD modeling results for incidence of mortality are summarized in Table 1-26. All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Logistic, Log-Probit, and Probit models. The Dichotomous Hill model was unusable because the BMDL computation failed (lower limit included zero). With BMRs of 10% or 5% ER applied, the BMDLs of the viable models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Log-Logistic) was selected. With a BMR of one percent ER applied, all models were considered questionable because the BMDLs (and in some cases the BMDs) were 10 times lower than the lowest non-zero concentration; no model was selected for the one percent ER BMR.

Table 1-26. Summary of BMD Modeling Results for Incidence of Mortality in Female Rabbits Following Inhalation Exposure to 1,2-Dichloroethane Using BMR of 10%ER or 5%ER^a

| Model | Goodness of Fit (Means) | | BMD 5%ER (mg/m ³) | BMDL 5%ER (mg/m ³) | BMD 10%ER (mg/m ³) | BMDL 10%ER (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|--------------|-------------------------------|--------------------------------|--------------------------------|---------------------------------|--|
| | p-value | AIC | | | | | |
| Dichotomous Hill | 0.7865 | 41.10 | 2.15E-06 | 0 | 7.17E-06 | 0 | All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Logistic, Log-Probit, and Probit models. The Dichotomous Hill model was unusable because the BMDL computation failed (lower limit included zero). The BMDLs of the fit models were sufficiently close (differed by < 3-fold); therefore, EPA chose the model with the lowest AIC. |
| Gamma | 0.1759 | 41.94 | 65.9 | 34.8 | 135 | 71.6 | |
| Log-Logistic | 0.2591 | 41.51 | 54.3 | 28.1 | 115 | 59.4 | |
| Multistage 2 | 0.1963 | 41.90 | 61.5 | 34.9 | 126 | 71.8 | |
| Multistage 1 | 0.1963 | 41.90 | 61.5 | 34.9 | 126 | 71.8 | |
| Weibull | 0.1963 | 41.90 | 61.5 | 34.9 | 126 | 71.8 | |
| Logistic | 0.0537 | 45.64 | 170 | 97.6 | 286 | 167 | |
| Log-Probit | – | – | – | – | – | – | |
| Probit | 0.0542 | 45.54 | 157 | 89.4 | 271 | 156 | |
| Quantal Linear | 0.1963 | 41.90 | 61.5 | 34.9 | 126 | 71.8 | |

^a Selected model in bold.

Plots of the Log-Logistic model with BMRs of 10% and 5% ER are shown in Figure 1-30 and Figure 1-31, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-32 (BMD and BMDL shown are for BMR of 10% ER; the rest is applicable to all BMRs).




Figure 1-30. Plot of Response by Dose with Fitted Curve for the Selected Model (Log-Logistic) for Mortality in Female Rabbits Exposed to 1,2-Dichloroethane Via Inhalation on GD 6 to 18 (7 Hours per Day) and BMR of 10%ER




Figure 1-31. Plot of Response by Dose with Fitted Curve for the Selected Model (Log-Logistic) for Mortality in Female Rabbits Exposed to 1,2-Dichloroethane Via Inhalation on GD 6 to 18 (7 Hours per Day) and BMR of 5%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 114.6125356 | | | | |
| BMDL | 59.4154249 | | | | |
| BMDU | 760.5776692 | | | | |
| AIC | 41.51198247 | | | | |
| P-value | 0.259131334 | | | | |
| D.O.F. | 2 | | | | |
| Chi ² | 2.700840529 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 3 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| a | -6.938781761 | 0.424700555 | -7.7711796 | -6.106384 | |
| b | Bounded | NA | NA | NA | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 3.046E-07 | 0 | 20 | -0.000552 |
| 118 | 0.102652196 | 2.155696122 | 4 | 21 | 1.3260451 |
| 353 | 0.254963342 | 4.844303497 | 3 | 19 | -0.970796 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -18.51225108 | 3 | - | - | NA |
| Fitted Model | -19.75599124 | 1 | 2.48748031 | 2 | 0.2883039 |
| Reduced Model | -21.61383127 | 1 | 6.20316038 | 2 | 0.0449781 |

Figure 1-32. Details Regarding the Selected Model (Log-Logistic) for Mortality in Female Rabbits Exposed to 1,2-Dichloroethane Via Inhalation on GD 6 to 18 (7 Hours per Day)

1.1.1.2.2 Body Weight effects

Two short-term/intermediate-duration inhalation studies were identified for BMD modeling that showed significant changes in body weight ([Zeng et al., 2018](#)) or body weight gain ([Igwe et al., 1986](#)).

1.1.1.2.2.1 Body Weight in Male Mice – 28-Day Inhalation Exposure

Body weight was significantly decreased at week 4 in male mice exposed to 1,2-dichloroethane by inhalation for 28 days (6 hours per day, 7 days per week) ([Zeng et al., 2018](#)). The measured exposure concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week. The concentration and response data used for the modeling are presented in Table 1-27. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected because EPA considers a 10% change in body weight to be biologically significant.

Table 1-27. Decreased Body Weight of Male Mice at Week 4 and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane in a 28-Day Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (g) | SD (g) |
|---|-------------------|----------|--------|
| 0.068 | 10 | 39.85 | 0.83 |
| 90.895 | 10 | 38.80 | 1.08 |
| 182.78 | 10 | 35.68 | 0.94 |

The BMD modeling results for decreased body weight are summarized in Table 1-28. The constant variance model provided adequate fit to the variance data, but with this model applied, none of the available models provided adequate fit to the means (all except for the Linear model were saturated, with degree of freedom = 0). The nonconstant variance model also provided adequate fit to the variance data. With the nonconstant variance model applied, the Exponential 3, Polynomial 2-degree, and Linear models provided adequate fit to the means. The BMDLs for the fit models were sufficiently close (< 3-fold); therefore, the model with the lowest AIC (Exponential 3) was selected. The Exponential 3 and the Polynomial 2-degree model had identical goodness-of-fit statistics; BMDS recommended the Exponential 3 model.

Table 1-28. Summary of BMD Modeling Results for Decreased Body Weight in Male Mice at Week 4 Following a 28-Day Inhalation Exposure to 1,2-Dichloroethane (Nonconstant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 10%RD (mg/m ³) | BMDL 10%RD (mg/m ³) | Basis for Model Selection |
|----------------------|-------------------------|--------------|------------------------------|-------------------------------|--------------------------------|---------------------------------|--|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.9116 | 87.24 | 84.480 | 57.281 | 178.52 | 162.78 | BMDLs for the fit models (Exponential 3, Polynomial 2-degree, and Linear models) were sufficiently close (< 3-fold); therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | 89.24 | 84.489 | 57.313 | 178.51 | 98.30 | |
| Hill | NA | 91.23 | 87.838 | 55.459 | 161.06 | 111.21 | |
| Polynomial Degree 2 | 0.9116 | 87.24 | 84.336 | 53.975 | 178.64 | 163.92 | |
| Power | NA | 89.24 | 84.437 | 56.727 | 178.64 | 163.15 | |
| Linear | 0.0237 | 92.72 | 44.945 | 35.028 | 175.95 | 148.16 | |

^a Selected model in bold.

Plots of the Exponential 3 model with BMRs of one SD and 10% RD are shown in Figure 1-33 and Figure 1-34, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-35 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-33. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Body Weight in Male Mice at Week 4 Following a 28-Day Inhalation Exposure to 1,2-Dichloroethane and a BMR of 1SD (Nonconstant Variance)




Figure 1-34. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Body Weight in Male Mice at Week 4 Following a 28-Day Inhalation Exposure to 1,2-Dichloroethane and a BMR of 10%RD (Nonconstant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 84.4797913 | | | | | | | |
| BMDL | 57.28149371 | | | | | | | |
| BMDU | 122.8881893 | | | | | | | |
| AIC | 87.24381926 | | | | | | | |
| Test 4 P-value | 0.911550593 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 5 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| a | 39.85000044 | 0.286645626 | 39.2881853 | 40.4118156 | | | | |
| b | 0.001852231 | 4.83E-04 | 0.0009047 | 0.00279976 | | | | |
| d | 2.0335034 | 5.03E-01 | 1.04725263 | 3.01975417 | | | | |
| rho | Bounded | NA | NA | NA | | | | |
| log-alpha | -0.19641642 | 2.58E-01 | -0.7023586 | 0.3095258 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0.068 | 10 | 39.84999997 | 39.85 | 39.85 | 0.90646015 | 0.83 | 0.83 | 9.10707E-08 |
| 90.895 | 10 | 38.80000002 | 38.8 | 38.8 | 0.90646015 | 1.08 | 1.08 | -7.0194E-08 |
| 182.78 | 10 | 35.68000002 | 35.68 | 35.68 | 0.90646015 | 0.94 | 0.94 | -5.25043E-08 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -39.62190963 | 4 | 87.2438193 | | | | | |
| A2 | -39.27530885 | 6 | 90.5506177 | | | | | |
| A3 | -39.61573992 | 5 | 89.2314798 | | | | | |
| fitted | -39.62190963 | 4 | 87.2438193 | | | | | |
| R | -63.20374602 | 2 | 130.407492 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 47.85687434 | 4 | <0.0001 | | | | | |
| 2 | 0.693201554 | 2 | 0.70708756 | | | | | |
| 3 | 0.680862138 | 1 | 0.40929001 | | | | | |
| 4 | 0.012339415 | 1 | 0.91155059 | | | | | |

Figure 1-35. Details Regarding the Selected Model (Exponential 3) for Decreased Body Weight in Male Mice at Week 4 Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days

1.1.1.2.2.2 Body Weight Gain in Male Rats – 30-Day Inhalation Exposure

Body weight gain was significantly decreased in male rats exposed to exposed to 1,2-dichloroethane by inhalation for 30 days (seven hours per day, 5 days per week) (Igwe et al., 1986). The measured exposure concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day and 7 days per week. The concentration and response data used for the modeling are presented in Table 1-29. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected.

Table 1-29. Decreased Body Weight Gain of Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 30-Day Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (g) | SD (g) |
|---|-------------------|----------|--------|
| 0 | 30 | 190 | 37 |
| 129 | 12 | 194 | 20 |
| 256 | 11 | 170 | 32 |
| 383 | 10 | 147 | 24 |

The BMD modeling results for decreased body weight gain are summarized in Table 1-30. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, all models, except for the Hill model, provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Polynomial 2-degree) was selected.

Table 1-30. Summary of BMD Modeling Results for Decreased Body Weight Gain of Male Rats Following Inhalation Exposure to 1,2-Dichloroethane for 30 Days (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 10%RD (ppm) | BMDL 10%RD (ppm) | Basis for Model Selection |
|----------------------------|-------------------------|--------------|------------------------------|-------------------------------|-----------------|------------------|---|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.4178 | 619.1 | 322 | 244 | 262 | 168 | All models, except for the Hill model, provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were < 3-fold different; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | 0.7034 | 618.5 | 265 | 243 | 254 | 177 | |
| Hill | NA | 620.5 | 269 | 239 | 253 | 228 | |
| Polynomial Degree 3 | 0.6343 | 617.3 | 336 | 246 | 282 | 164 | |
| Polynomial Degree 2 | 0.6688 | 617.2 | 317 | 248 | 250 | 166 | |
| Power | 0.3992 | 619.1 | 325 | 305 | 264 | 168 | |
| Linear | 0.1894 | 619.7 | 306 | 209 | 188 | 134 | |

^a Selected model in bold.

Plots of the Polynomial 2-degree model with BMRs of one SD and 10% RD are shown in Figure 1-36 and Figure 1-37, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-38 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-36. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Body Weight Gain in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days and BMR of 1SD (Constant Variance)




Figure 1-37. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Body Weight Gain in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days and BMR of 10%RD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 317.080575 | | | | | | | |
| BMDL | 248.0936048 | | | | | | | |
| BMDU | 417.0049545 | | | | | | | |
| AIC | 617.2047371 | | | | | | | |
| Test 4 P-value | 0.668842162 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 192.1048183 | 4.79868076 | 182.699577 | 201.51006 | | | | |
| beta | Bounded | NA | NA | NA | | | | |
| beta2 | -0.000307689 | 7.39E-05 | -0.0004525 | -0.0001629 | | | | |
| alpha | 956.9809221 | 1.63E+05 | -318857.33 | 320771.297 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 30 | 192.1048183 | 190 | 190 | 30.9351082 | 37 | 37 | -0.37266928 |
| 129 | 12 | 186.984561 | 194 | 194 | 30.9351082 | 20 | 20 | 0.785586185 |
| 256 | 11 | 171.9400936 | 170 | 170 | 30.9351082 | 32 | 32 | -0.208001943 |
| 383 | 10 | 146.9701855 | 147 | 147 | 30.9351082 | 24 | 24 | 0.003047727 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -305.2001614 | 5 | 620.400323 | | | | | |
| A2 | -301.4914854 | 8 | 618.982971 | | | | | |
| A3 | -305.2001614 | 5 | 620.400323 | | | | | |
| fitted | -305.6023685 | 3 | 617.204737 | | | | | |
| R | -313.2489895 | 2 | 630.497979 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 23.51500828 | 6 | 0.00064112 | | | | | |
| 2 | 7.417351949 | 3 | 0.0597205 | | | | | |
| 3 | 7.417351949 | 3 | 0.0597205 | | | | | |
| 4 | 0.804414355 | 2 | 0.66884216 | | | | | |

Figure 1-38. Details Regarding the Selected Model (Polynomial 2-Degree) for Body Weight Gain in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 30 Days

1.1.1.2.3 Hepatic Effects:

EPA modeled both absolute and relative liver weight changes in male mice in a 28-day inhalation study (Zeng et al., 2018). Absolute liver weight in male mice, though not statistically significant, was increased by >10% at both concentrations tested. Relative liver weight in male mice was increased by >10% at both tested concentrations and reached statistical significance at the highest concentration. EPA also modeled statistically significant increases in serum alanine aminotransferase (ALT) and aspartate aminotransferase (AST) in the male mice.

Modeled results for relative liver weight and serum liver enzyme levels from the Zeng et al. (2018) studies are presented. Results for absolute liver weight changes in male mice are not presented because neither the constant nor the nonconstant variance models provided adequate fit to the variance data.

EPA also modeled statistically significant metabolic changes in the liver, including increased liver concentrations of glycogen, triglycerides, and free fatty acids in male mice (Zeng et al., 2018). These results are not shown because, for each of these data sets, none of the models provided adequate fits to the data either assuming constant or nonconstant variance.

1.1.1.2.3.1 Relative Liver Weight in Male Mice – 28-Day Inhalation Exposure

Relative liver weight was significantly increased in male mice exposed to 1,2-dichloroethane by inhalation for 28 days (6 hours per day, 7 days per week) (Zeng et al., 2018). The measured exposure concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week. The concentration and response data used for the modeling are presented in Table 1-31. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected because EPA considers a 10% change in relative liver weight to be biologically significant.

Table 1-31. Relative Liver Weight in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 28-Day Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (Liver/Body Weight Ratio) | SD (Liver/Body Weight Ratio) |
|---|-------------------|--------------------------------|------------------------------|
| 0.068 | 10 | 4.38 | 0.30 |
| 90.895 | 10 | 5.14 | 0.46 |
| 182.78 | 10 | 5.59 | 0.46 |

The BMD modeling results for increased relative liver weight are summarized in Table 1-32. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, all models, except for the Exponential 5 and Hill models, provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected. The Power model converged on the Linear model and these had the lowest AIC; the Linear model was selected as the more parsimonious choice.

Table 1-32. Summary of BMD Modeling Results for Increased Relative Liver Weight in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 28 Days (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 10%RD (mg/m ³) | BMDL 10%RD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|----------------|------------------------------|-------------------------------|--------------------------------|---------------------------------|---|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.2069 | 36.60 | 66.637 | 51.379 | 73.314 | 58.420 | All models, except for the Exponential 5 and Hill models, provided adequate fit to the means (test 4 p-value > 0.1). Among the fit models, BMDLs differed by < 3-fold; the Linear model was selected based on the lowest AIC. |
| Exponential 5 | NA | 37.00 | 40.867 | 20.758 | 46.314 | 22.869 | |
| Hill | NA | 39.00 | 56.026 | 15.337 | 59.981 | 16.927 | |
| Polynomial Degree 2 | 0.3012 | 36.0735 | 60.246 | 45.510 | 66.752 | 51.726 | |
| Power | 0.3013 | 36.0730 | 60.345 | 45.508 | 66.968 | 51.720 | |
| Linear | 0.3013 | 36.0730 | 60.345 | 45.508 | 66.968 | 51.720 | |

^a Selected model in bold.

Plots of the Linear model with BMRs of one SD and 10% RD are shown in Figure 1-39 and Figure 1-40, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-41 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-39. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days and BMR of 1SD (Constant Variance)




Figure 1-40. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days and BMR of 10%RD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 60.34484029 | | | | | | | |
| BMDL | 45.50823313 | | | | | | | |
| BMDU | 88.69790107 | | | | | | | |
| AIC | 36.07303736 | | | | | | | |
| Test 4 P-value | 0.301262139 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 3 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 4.432689666 | 0.115222371 | 4.20685797 | 4.65852137 | | | | |
| beta | 0.006619095 | 9.78E-04 | 0.00470294 | 0.00853525 | | | | |
| alpha | 0.15954289 | 6.57E-03 | 0.14666168 | 0.1724241 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0.068 | 10 | 4.433139765 | 4.38 | 4.38 | 0.3994282 | 0.3 | 0.3 | -0.420708124 |
| 90.895 | 10 | 5.034332331 | 5.14 | 5.14 | 0.3994282 | 0.46 | 0.46 | 0.836572145 |
| 182.78 | 10 | 5.6425279 | 5.59 | 5.59 | 0.3994282 | 0.46 | 0.46 | -0.415863983 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -14.50221924 | 4 | 37.0044385 | | | | | |
| A2 | -13.41744443 | 6 | 38.8348889 | | | | | |
| A3 | -14.50221924 | 4 | 37.0044385 | | | | | |
| fitted | -15.03651868 | 3 | 36.0730374 | | | | | |
| R | -28.94761589 | 2 | 61.8952318 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 31.06034292 | 4 | <0.0001 | | | | | |
| 2 | 2.169549619 | 2 | 0.33797789 | | | | | |
| 3 | 2.169549619 | 2 | 0.33797789 | | | | | |
| 4 | 1.068598887 | 1 | 0.30126214 | | | | | |

Figure 1-41. Details Regarding the Selected Model (Linear) for Relative Liver Weight Increases in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days

1.1.1.2.3.2 Serum ALT in Male Mice – 28-Day Inhalation Exposure

Serum ALT levels were significantly increased in male mice exposed to 1,2-dichloroethane by inhalation for 28 days (6 hours per day, 7 days per week) (Zeng et al., 2018). The measured exposure

concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week. The concentration and response data used for the modeling are presented in Table 1-33. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-33. Serum ALT in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 28-Day Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (U/L) | SD (U/L) |
|---|-------------------|------------|----------|
| 0.068 | 10 | 34.80 | 2.78 |
| 90.895 | 10 | 50.40 | 8.06 |
| 182.78 | 10 | 65.20 | 7.45 |

The BMD modeling results for increased serum ALT are summarized in Table 1-34. The constant variance model did not provide adequate fit to the variance data, but the nonconstant variance model did. With the nonconstant variance model applied, the Polynomial 2-degree, Power, and Linear models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (< 3-fold); therefore, the model with the lowest AIC (Linear Model) was selected.

Table 1-34. Summary of BMD Modeling Results for Increased Serum ALT in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 28 Days (Nonconstant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|-----------------|------------------------------|-------------------------------|---|
| | Test 4 p-value | AIC | | | |
| Exponential 3 | 0.0657 | 198.7 | 27.148 | 18.025 | The Polynomial 2-degree, Power, and Linear models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | 199.3 | 18.907 | 7.8091 | |
| Hill | NA | 199.3 | 31.175 | 7.0150 | |
| Polynomial Degree 2 | 0.4543 | 195.8730 | 18.715 | 12.621 | |
| Power | 0.4535 | 195.8752 | 18.607 | 12.620 | |
| Linear | 0.4543 | 195.8729 | 18.705 | 12.620 | |

^a Selected model in bold.

A plot of the Linear model with a BMR of one SD is shown in Figure 1-42. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-43.




Figure 1-42. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Serum ALT in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days and BMR of 1SD (Nonconstant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 18.70544198 | | | | | | | |
| BMDL | 12.62045044 | | | | | | | |
| BMDU | 30.11786611 | | | | | | | |
| AIC | 195.8729428 | | | | | | | |
| Test 4 P-value | 0.454349562 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 34.80093952 | 0.977490421 | 32.8850935 | 36.7167856 | | | | |
| beta | 0.16873094 | 0.014569002 | 0.14017622 | 0.19728566 | | | | |
| rho | 3.377393896 | 1.243700214 | 0.93978625 | 5.81500154 | | | | |
| alpha | 6.19107E-05 | 1.85E-08 | 6.1874E-05 | 6.1947E-05 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0.068 | 10 | 34.81241322 | 34.8 | 34.8 | 3.15794416 | 2.78 | 2.78 | -0.012430257 |
| 90.895 | 10 | 50.13773829 | 50.4 | 50.4 | 5.84717017 | 8.06 | 8.06 | 0.141836875 |
| 182.78 | 10 | 65.64158069 | 65.2 | 65.2 | 9.21610491 | 7.45 | 7.45 | -0.151517454 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -97.31179356 | 4 | 202.623587 | | | | | |
| A2 | -92.16353343 | 6 | 196.327067 | | | | | |
| A3 | -93.65658221 | 5 | 197.313164 | | | | | |
| fitted | -93.93647138 | 4 | 195.872943 | | | | | |
| R | -121.4712012 | 2 | 246.942402 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 58.61533564 | 4 | <0.0001 | | | | | |
| 2 | 10.29652026 | 2 | 0.0058095 | | | | | |
| 3 | 2.986097556 | 1 | 0.08398233 | | | | | |
| 4 | 0.559778352 | 1 | 0.45434956 | | | | | |

Figure 1-43. Details Regarding the Selected Model (Linear) for Increased Serum ALT in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days

1.1.1.2.3.3 Serum AST in Male Mice – 28-Day Inhalation Exposure

Serum AST levels were significantly increased in male mice exposed to 1,2-dichloroethane by inhalation for 28 days (6 hours per day, 7 days per week) (Zeng et al., 2018). The measured exposure

concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week. The concentration and response data used for the modeling are presented in Table 1-35. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-35. Serum AST in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 28-Day Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (U/L) | SD (U/L) |
|---|-------------------|------------|----------|
| 0.068 | 10 | 137.30 | 9.09 |
| 90.895 | 10 | 182.30 | 8.82 |
| 182.78 | 10 | 231.80 | 18.09 |

The BMD modeling results for increased serum AST are summarized in Table 1-36. The constant variance model did not provide adequate fit to the variance data, but the nonconstant variance model did. With the nonconstant variance model applied, the Exponential 3 and Linear models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (< 3-fold); therefore, the model with the lowest AIC (Linear model) was selected.

Table 1-36. Summary of BMD Modeling Results for Increased Serum AST in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 28 Days (Nonconstant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|--------------|------------------------------|-------------------------------|---|
| | Test 4 p-value | AIC | | | |
| Exponential 3 | 0.3721 | 238.3 | 19.271 | 14.286 | The Exponential 3 and Linear models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | 241.5 | 34.809 | 11.861 | |
| Hill | NA | 241.5 | 28.918 | 11.976 | |
| Polynomial Degree 2 | NA | 239.5 | 16.812 | 11.559 | |
| Power | NA | 239.5 | 18.264 | 11.562 | |
| Linear | 0.5631 | 237.8 | 15.370 | 11.360 | |

^a Selected model in bold.

A plot of the Linear model with a BMR of one SD is shown in Figure 1-44. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-45.




Figure 1-44. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Serum AST in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days and BMR of 1SD (Nonconstant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 15.37039918 | | | | | | | |
| BMDL | 11.35979052 | | | | | | | |
| BMDU | 22.23350808 | | | | | | | |
| AIC | 237.8188306 | | | | | | | |
| Test 4 P-value | 0.563122068 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 136.9827797 | 2.397347299 | 132.284065 | 141.681494 | | | | |
| beta | 0.512404641 | 2.81E-02 | 0.45735724 | 0.56745205 | | | | |
| rho | 2.562173101 | 1.12E+00 | 0.37562471 | 4.7487215 | | | | |
| alpha | 0.000208012 | 2.51E-07 | 0.00020752 | 0.0002085 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0.068 | 10 | 137.0176232 | 137.3 | 137.3 | 7.87843043 | 9.09 | 9.09 | 0.113341577 |
| 90.895 | 10 | 183.5577996 | 182.3 | 182.3 | 11.4586407 | 8.82 | 8.82 | -0.347118969 |
| 182.78 | 10 | 230.6401001 | 231.8 | 231.8 | 15.3521135 | 18.09 | 18.09 | 0.238919918 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -117.3531051 | 4 | 242.70621 | | | | | |
| A2 | -113.783309 | 6 | 239.566618 | | | | | |
| A3 | -114.7422517 | 5 | 239.484503 | | | | | |
| fitted | -114.9094153 | 4 | 237.818831 | | | | | |
| R | -153.5664702 | 2 | 311.13294 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 79.56632236 | 4 | <0.0001 | | | | | |
| 2 | 7.139592092 | 2 | 0.0281616 | | | | | |
| 3 | 1.917885267 | 1 | 0.16608997 | | | | | |
| 4 | 0.334327285 | 1 | 0.56312207 | | | | | |

Figure 1-45. Details Regarding the Selected Model (Linear) for Increased Serum AST in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 28 Days

1.1.1.2.4 Reproductive Effects

Reproductive endpoints from a short-term/intermediate inhalation study were identified for BMD modeling (Zhang et al., 2017). EPA did not present the BMD modeling results for abnormalities in the sperm head, abnormalities in sperm body, or total sperm abnormalities because there were no adequately fit models with either the constant or nonconstant variance models. Although the Hill model provided adequate fit to data for abnormalities in sperm tail (with nonconstant variance applied), the results are not presented because the BMD/BMDL ratio was greater than 10 and the BMDL was more than three times lower than the lowest dose tested.

1.1.1.2.4.1 Sperm Concentration in Male Mice – 4-Week Inhalation Exposure

Sperm concentration was significantly decreased in male mice exposed to 1,2-dichloroethane by inhalation for four weeks (6 hours per day, 7 days per week) (Zhang et al., 2017). The measured exposure concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week. The concentration and response data used for the modeling are presented in Table 1-37. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 5% RD was also selected because EPA considers a 5% change in sperm concentration to be biologically relevant. In some strains of rats and mice, production of normal sperm can be reduced by up to 90% or more without compromising fertility (Working, 1988; Robaire et al., 1984; Meistrich et al., 1982; Aafjes et al., 1980). However, less severe reductions can cause reduced fertility in human males who appear to function closer to the threshold for the number of normal sperm needed to ensure full reproductive competence. This difference between test species and humans suggests that results from a test species may not fully represent toxicity in humans due to chemical exposure.

Table 1-37. Sperm Concentration in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 4-Week Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (M/g) | SD (M/g) |
|---|-------------------|------------|----------|
| 0.075 | 10 | 4.65 | 0.52 |
| 25.675 | 10 | 4.36 | 0.40 |
| 89.010 | 10 | 3.89 | 0.47 |
| 176.75 | 10 | 3.30 | 0.57 |

The BMD modeling results for decreased sperm concentration are summarized in Table 1-38. The constant variance model provided adequate fit to the variance data. With the constant variance model applied, all models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Exponential 3) was selected.

Table 1-38. Summary of BMD Modeling Results for Decreased Sperm Concentration in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 4 Weeks (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 5%RD (mg/m ³) | BMDL 5%RD (mg/m ³) | Basis for Model Selection |
|----------------------|-------------------------|--------------|------------------------------|-------------------------------|-------------------------------|--------------------------------|---|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.9453 | 59.00 | 55.847 | 41.335 | 26.735 | 21.240 | All models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were < 3-fold different; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | 0.7855 | 60.96 | 51.913 | 26.379 | 24.578 | 12.196 | |
| Hill | 0.7929 | 60.95 | 51.506 | 23.774 | 24.276 | 10.475 | |
| Polynomial Degree 3 | 0.8432 | 59.23 | 64.002 | 48.238 | 31.202 | 25.351 | |
| Polynomial Degree 2 | 0.8484 | 59.21 | 62.561 | 48.259 | 30.564 | 25.353 | |
| Power | 0.8494 | 59.21 | 62.986 | 48.252 | 30.748 | 25.351 | |
| Linear | 0.8494 | 59.21 | 62.986 | 48.252 | 30.748 | 25.351 | |

^a Selected model in bold.

Plots of the Exponential 3 model with a BMR of one SD and 5% RD are shown in Figure 1-46 and Figure 1-47, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-48. (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-46. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Sperm Concentration in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 1SD (Constant Variance)




Figure 1-47. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Sperm Concentration in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 5%RD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 55.84663409 | | | | | | | |
| BMDL | 41.33515229 | | | | | | | |
| BMDU | 100.8093304 | | | | | | | |
| AIC | 58.99784667 | | | | | | | |
| Test 4 P-value | 0.945316531 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| a | 4.618972239 | 0.113350836 | 4.39680868 | 4.8411358 | | | | |
| b | 0.001918616 | 2.92E-04 | 0.00134608 | 0.00249115 | | | | |
| d | Bounded | NA | NA | NA | | | | |
| log-alpha | -1.512930902 | 2.24E-01 | -1.9511885 | -1.0746734 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0.075 | 10 | 4.618307635 | 4.65 | 4.65 | 0.46932234 | 0.52 | 0.52 | 0.213542059 |
| 25.675 | 10 | 4.396952797 | 4.36 | 4.36 | 0.46932234 | 0.4 | 0.4 | -0.248986667 |
| 89.01 | 10 | 3.893841526 | 3.89 | 3.89 | 0.46932234 | 0.47 | 0.47 | -0.025884069 |
| 176.75 | 10 | 3.29055696 | 3.3 | 3.3 | 0.46932234 | 0.57 | 0.57 | 0.063626873 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -26.44268788 | 5 | 62.8853758 | | | | | |
| A2 | -25.776744 | 8 | 67.553488 | | | | | |
| A3 | -26.44268788 | 5 | 62.8853758 | | | | | |
| fitted | -26.49892334 | 3 | 58.9978467 | | | | | |
| R | -42.10709522 | 2 | 88.2141904 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 32.66070244 | 6 | <0.0001 | | | | | |
| 2 | 1.331887768 | 3 | 0.7215753 | | | | | |
| 3 | 1.331887768 | 3 | 0.7215753 | | | | | |
| 4 | 0.112470908 | 2 | 0.94531653 | | | | | |

Figure 1-48. Details Regarding the Selected Model (Exponential 3) for Decreased Sperm Concentration in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks

1.1.1.2.4.2 Diameter of Seminiferous Tubules in Male Mice – 4-Week Inhalation Exposure

The diameter of seminiferous tubules was significantly decreased in male mice exposed to 1,2-dichloroethane by inhalation for four weeks (6 hours per day, 7 days per week) (Zhang et al., 2017). The measured exposure concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week. The concentration and response data used for the modeling are presented in Table 1-39. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012).

Table 1-39. Diameter of Seminiferous Tubules in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 4-Week Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (µm) | SD (µm) |
|---|-------------------|-----------|---------|
| 0.075 | 5 | 249 | 29.3 |
| 25.675 | 5 | 236 | 28.2 |
| 89.010 | 5 | 180 | 19.2 |
| 176.75 | 5 | 100 | 11.3 |

The BMD modeling results for decreased diameter of seminiferous tubules are summarized in Table 1-40. The constant variance model provided adequate fit to the variance data. With the constant variance model applied, the Exponential 3, Power, and Linear models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Linear) was selected.

Table 1-40. Summary of BMD Modeling Results for Decreased Diameter of Seminiferous Tubules in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 4 Weeks (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|--------------|------------------------------|-------------------------------|--|
| | Test 4 p-value | AIC | | | |
| Exponential 3 | 0.9424 | 186.0 | 37.039 | 20.640 | The Exponential 3, Power, and Linear models provided adequate fit to the means (test 4 p-value > 0.1). BMDLs differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | 188.0 | 37.039 | 20.640 | |
| Hill | NA | 189.0 | 81.915 | 16.819 | |
| Polynomial Degree 3 | NA | 188.3 | 28.539 | 19.113 | |
| Polynomial Degree 2 | 0.0353 | 190.5 | 64.600 | 63.257 | |
| Power | 0.7415 | 186.1 | 31.371 | 19.242 | |
| Linear | 0.7447 | 184.6 | 24.471 | 18.815 | |

^a Selected model in bold.

A plot of the Linear model with a BMR of one SD is shown in Figure 1-49. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-50.




Figure 1-49. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Decreased Diameter of Seminiferous Tubules in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 1SD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 24.47144499 | | | | | | | |
| BMDL | 18.81509191 | | | | | | | |
| BMDU | 34.10427505 | | | | | | | |
| AIC | 184.6234089 | | | | | | | |
| Test 4 P-value | 0.744678566 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 3 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 253.9223097 | 6.889545669 | 240.419048 | 267.425571 | | | | |
| beta | -0.85996788 | 6.90E-02 | -0.9953008 | -0.724635 | | | | |
| alpha | 442.8775746 | 6.20E+04 | -121124.09 | 122009.843 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0.075 | 5 | 253.8578122 | 249 | 249 | 21.0446567 | 29.3 | 29.3 | -0.51615944 |
| 25.675 | 5 | 231.8426344 | 236 | 236 | 21.0446567 | 28.2 | 28.2 | 0.441734553 |
| 89.01 | 5 | 177.3765688 | 180 | 180 | 21.0446567 | 19.2 | 19.2 | 0.278748693 |
| 176.75 | 5 | 101.922987 | 100 | 100 | 21.0446567 | 11.3 | 11.3 | -0.20432406 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -89.01690182 | 5 | 188.033804 | | | | | |
| A2 | -86.63044764 | 8 | 189.260895 | | | | | |
| A3 | -89.01690182 | 5 | 188.033804 | | | | | |
| fitted | -89.31170443 | 3 | 184.623409 | | | | | |
| R | -111.0087441 | 2 | 226.017488 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 48.75659293 | 6 | <0.0001 | | | | | |
| 2 | 4.772908345 | 3 | 0.18920143 | | | | | |
| 3 | 4.772908345 | 3 | 0.18920143 | | | | | |
| 4 | 0.589605219 | 2 | 0.74467857 | | | | | |

Figure 1-50. Details Regarding the Selected Model (Linear) for Decreased Diameter of Seminiferous Tubules in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks

1.1.1.2.4.3 Height of Germinal Epithelium in Male Mice – 4-Week Inhalation Exposure

The height of germinal epithelium was significantly decreased in male mice exposed to 1,2-dichloroethane by inhalation for four weeks (6 hours per day, 7 days per week) (Zhang et al., 2017). The measured exposure concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week. The concentration and response data used for the modeling are presented in Table 1-41. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012).

Table 1-41. Height of Germinal Epithelium in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 4-Week Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (µm) | SD (µm) |
|---|-------------------|-----------|---------|
| 0.075 | 5 | 100 | 8.2 |
| 25.675 | 5 | 90 | 9.4 |
| 89.010 | 5 | 52 | 8.2 |
| 176.75 | 5 | 30 | 5.9 |

The BMD modeling results for decreased height of germinal epithelium are summarized in Table 1-42. The constant variance model provided adequate fit to the variance data. With the constant variance model applied, only the Exponential 3 model provided adequate fit to the means (test 4 p-value > 0.1); therefore, this model was selected.

Table 1-42. Summary of BMD Modeling Results for Decreased Height of Germinal Epithelium in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 4 Weeks (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | Basis for Model Selection |
|----------------------|-------------------------|--------------|------------------------------|-------------------------------|---|
| | Test 4 p-value | AIC | | | |
| Exponential 3 | 0.1184 | 146.0 | 13.975 | 8.6304 | Only the Exponential 3 model provided adequate fit to the means (test 4 p-value > 0.1); therefore, this model was selected. |
| Exponential 5 | NA | 145.6 | 20.581 | 10.711 | |
| Hill | NA | 145.6 | 20.977 | 11.178 | |
| Polynomial Degree 3 | < 0.0001 | 168.0 | 71.926 | 70.452 | |
| Polynomial Degree 2 | 0.0022 | 152.9 | 22.452 | 17.037 | |
| Power | 0.0010 | 150.8 | 22.172 | 17.118 | |
| Linear | 0.0010 | 150.8 | 22.172 | 17.118 | |

^a Selected model in bold.

A plot of the Exponential 3 model with a BMR of one SD is shown in Figure 1-51. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-52.




Figure 1-51. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Height of Germinal Epithelium in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 1SD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 13.97510463 | | | | | | | |
| BMDL | 8.630430863 | | | | | | | |
| BMDU | 25.05659301 | | | | | | | |
| AIC | 146.0395411 | | | | | | | |
| Test 4 P-value | 0.118373042 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| a | 101.3424129 | 3.217463633 | 95.0363 | 107.648526 | | | | |
| b | 0.007113595 | 5.18E-04 | 0.00609928 | 0.00812791 | | | | |
| d | 1.103577087 | 1.41E-01 | 0.82721977 | 1.3799344 | | | | |
| log-alpha | 4.064099969 | 3.16E-01 | 3.44430945 | 4.68389048 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0.075 | 5 | 101.3176446 | 100 | 100 | 7.62971113 | 8.2 | 8.2 | -0.386167039 |
| 25.675 | 5 | 86.95185553 | 90 | 90 | 7.62971113 | 9.4 | 9.4 | 0.893331102 |
| 89.01 | 5 | 55.40096565 | 52 | 52 | 7.62971113 | 8.2 | 8.2 | -0.996733724 |
| 176.75 | 5 | 27.96627608 | 30 | 30 | 7.62971113 | 5.9 | 5.9 | 0.596031076 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -67.80041229 | 5 | 145.600825 | | | | | |
| A2 | -67.26698689 | 8 | 150.533974 | | | | | |
| A3 | -67.80041229 | 5 | 145.600825 | | | | | |
| fitted | -69.01977056 | 4 | 146.039541 | | | | | |
| R | -95.87257757 | 2 | 195.745155 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 57.21118136 | 6 | <0.0001 | | | | | |
| 2 | 1.066850786 | 3 | 0.78508185 | | | | | |
| 3 | 1.066850786 | 3 | 0.78508185 | | | | | |
| 4 | 2.438716552 | 1 | 0.11837304 | | | | | |

Figure 1-52. Details Regarding the Selected Model (Exponential 3) for Decreased Height of Germinal Epithelium in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks

1.1.1.2.4.4 Number of Apoptotic Cells in the Testis in Male Mice – 4-Week Inhalation Exposure

The number of apoptotic cells in the testis was significantly increased in male mice exposed to 1,2-dichloroethane by inhalation for four weeks (6 hours per day, 7 days per week) (Zhang et al., 2017). The measured exposure concentrations were duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week. The concentration and response data used for the modeling are presented in Table 1-43. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012).

Table 1-43. Number of Apoptotic Cells in Male Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 4-Week Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean | SD |
|---|-------------------|------|-----|
| 0.075 | 5 | 169 | 108 |
| 25.675 | 5 | 207 | 160 |
| 89.010 | 5 | 273 | 198 |
| 176.75 | 5 | 400 | 216 |

The BMD modeling results for increased number of apoptotic cells are summarized in Table 1-44. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, all models, except for the Exponential 5 and Hill models, provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Linear) was selected.

Table 1-44. Summary of BMD Modeling Results for Increased Number of Apoptotic Cells in Male Mice Following Inhalation Exposure to 1,2-Dichloroethane for 4 Weeks (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|----------------|------------------------------|-------------------------------|--|
| | Test 4 p-value | AIC | | | |
| Exponential 3 | 0.9847 | 265.019 | 137.37 | 98.972 | All models, except for the Exponential 5 and Hill models, provided adequate fit to the means (test 4 p-value > 0.1). BMDLs differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | 269.009 | 127.39 | 87.083 | |
| Hill | NA | 269.016 | 127.36 | 23.227 | |
| Polynomial Degree 3 | 0.9605 | 265.068 | 112.45 | 71.178 | |
| Polynomial Degree 2 | 0.9054 | 267.002 | 128.54 | 71.734 | |
| Power | 0.8850 | 267.009 | 127.39 | 71.487 | |
| Linear | 0.9852 | 265.018 | 122.18 | 71.431 | |

^a Selected model in bold.

A plot of the Linear model with a BMR of one SD is shown in Figure 1-53. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-54.




Figure 1-53. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Number of Apoptotic Cells in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks and BMR of 1SD (Constant Variance)

Model Results**Benchmark Dose**

| | |
|----------------|-------------|
| BMD | 122.1802163 |
| BMDL | 71.4305637 |
| BMDU | 417.429177 |
| AIC | 265.0175897 |
| Test 4 P-value | 0.985180162 |
| D.O.F. | 2 |

Model Parameters

| # of Parameters | 3 | | | |
|-----------------|-------------|-------------|------------|------------|
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 168.5794306 | 51.40667943 | 67.8241895 | 269.334672 |
| beta | 1.285315329 | 0.515232602 | 0.27547798 | 2.29515268 |
| alpha | 24661.59453 | 192327488.6 | -376930292 | 376979615 |

Goodness of Fit

| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
|--------|------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| 0.075 | 5 | 168.6758292 | 169 | 169 | 157.040105 | 108 | 108 | 0.004615814 |
| 25.675 | 5 | 201.5799016 | 207 | 207 | 157.040105 | 160 | 160 | 0.077175881 |
| 89.01 | 5 | 282.985348 | 273 | 273 | 157.040105 | 198 | 198 | -0.142179712 |
| 176.75 | 5 | 395.7589149 | 400 | 400 | 157.040105 | 216 | 216 | 0.060388106 |

Likelihoods of Interest

| Model | Log Likelihood* | # of Parameters | AIC |
|--------|-----------------|-----------------|------------|
| A1 | -129.4938641 | 5 | 268.987728 |
| A2 | -128.2515876 | 8 | 272.503175 |
| A3 | -129.4938641 | 5 | 268.987728 |
| fitted | -129.5087949 | 3 | 265.01759 |
| R | -132.2176673 | 2 | 268.435335 |

| Tests of Interest | | | |
|-------------------|--------------------------|---------|------------|
| Test | -2*Log(Likelihood Ratio) | Test df | p-value |
| 1 | 7.9321595 | 6 | 0.24311574 |
| 2 | 2.484553103 | 3 | 0.47808915 |
| 3 | 2.484553103 | 3 | 0.47808915 |
| 4 | 0.029861498 | 2 | 0.98518016 |

Figure 1-54. Details Regarding the Selected Model (Linear) for Increased Number of Apoptotic Cells in Male Mice Exposed to 1,2-Dichloroethane Via Inhalation for 4 Weeks

1.1.1.3 Chronic

1.1.1.3.1 Hepatic Effects

EPA identified hepatic endpoints in a chronic inhalation study for BMD modeling ([IRFMN, 1978](#)). Modeling results are presented for increased serum lactate dehydrogenase (LDH) levels in female rats. Modeled results are not presented for serum LDH levels in male rats or serum ALT or cholesterol levels in male or female rats because neither the constant nor nonconstant variance models provided adequate fit to the variance data or because none of the models provided adequate fits to the means (test 4 p-value < 0.1) assuming either constant or nonconstant variance.

1.1.1.3.1.1 LDH Levels in Female Sprague-Dawley Rats – 12-Month Inhalation Exposure

Serum LDH levels were significantly increased in female rats exposed to 1,2-dichloroethane by inhalation for 12 months (seven hours per day, 5 days per week) ([IRFMN, 1978](#)). The exposure concentrations (reported in ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day and 7 days per week. The concentration and response data used for the modeling are presented in Table 1-45. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-45. Increased LDH Levels in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (mU/mL) | SD (mU/mL) |
|---|-------------------|--------------|------------|
| 0 | 8 | 617.50 | 47.12 |
| 4 | 8 | 682.50 | 78.15 |
| 8.3 | 8 | 700.00 | 81.40 |
| 42 | 8 | 770.00 | 51.28 |
| 126 | 8 | 705.00 | 91.81 |

The BMD modeling results for increased serum LDH levels in female rats are summarized in Table 1-46. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, only the Exponential 5 model provided adequate fit to the means (test 4 p-value > 0.1); therefore, this model was selected.

Table 1-46. Summary of BMD Modeling Results for Increased LDH Levels in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | Basis for Model Selection |
|----------------------|-------------------------|--------------|------------------------------|-------------------------------|---|
| | Test 4 p-value | AIC | | | |
| Exponential 3 | 0.0019 | 471.3 | 210 | 100 | Only the Exponential 5 model provided adequate fit to the means (test 4 p-value > 0.1); therefore, this model was selected. |
| Exponential 5 | 0.1563 | 462.2 | 5.5 | 1.7 | |
| Hill | 0.0007 | 472.8 | 210 | 210 | |
| Polynomial Degree 3 | 0.0020 | 471.2 | 210 | 96 | |
| Polynomial Degree 2 | 0.0020 | 471.2 | 210 | 96 | |
| Power | 0.0020 | 471.2 | 210 | 96 | |
| Linear | 0.0020 | 471.2 | 210 | 96 | |

^a Selected model in bold.

A plot of the Exponential 5 model with a BMR of one SD is shown in Figure 1-55. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-56.




Figure 1-55. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 5) for Increased LDH Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance)

Model Results

| Benchmark Dose | |
|----------------|-------------|
| BMD | 5.542740664 |
| BMDL | 1.66153532 |
| BMDU | Infinity |
| AIC | 462.1616961 |
| Test 4 P-value | 0.156276737 |
| D.O.F. | 2 |

| Model Parameters | | | | |
|------------------|-------------|-------------|------------|------------|
| # of Parameters | 5 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| a | 619.0800415 | 24.60242104 | 570.860182 | 667.299901 |
| b | 0.166171746 | 9.54E-02 | -0.0207616 | 0.35310507 |
| c | 1.18966716 | 4.58E-02 | 1.09996 | 1.27937432 |
| d | Bounded | NA | NA | NA |
| log-alpha | 8.516165337 | 2.24E-01 | 8.07790512 | 8.95442556 |

| Goodness of Fit | | | | | | | | |
|-----------------|------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 8 | 619.0800415 | 617.5 | 617.5 | 70.6743473 | 47.12 | 47.12 | -0.063234151 |
| 4 | 8 | 676.094728 | 682.5 | 682.5 | 70.6743473 | 78.15 | 78.15 | 0.256342587 |
| 8.3 | 8 | 706.9361662 | 700 | 700 | 70.6743473 | 81.4 | 81.4 | -0.277589271 |
| 42 | 8 | 736.3898734 | 770 | 770 | 70.6743473 | 51.28 | 51.28 | 1.345096167 |
| 126 | 8 | 736.4991947 | 705 | 705 | 70.6743473 | 91.81 | 91.81 | -1.260615487 |

| Likelihoods of Interest | | | |
|-------------------------|-----------------|-----------------|------------|
| Model | Log Likelihood* | # of Parameters | AIC |
| A1 | -225.2247211 | 6 | 462.449442 |
| A2 | -222.6287127 | 10 | 465.257425 |
| A3 | -225.2247211 | 6 | 462.449442 |
| fitted | -227.080848 | 4 | 462.161696 |
| R | -233.6433119 | 2 | 471.286624 |

| Tests of Interest | | | |
|-------------------|--------------------------|---------|------------|
| Test | -2*Log(Likelihood Ratio) | Test df | p-value |
| 1 | 22.0291984 | 8 | 0.00486206 |
| 2 | 5.192016838 | 4 | 0.26815665 |
| 3 | 5.192016838 | 4 | 0.26815665 |
| 4 | 3.712253773 | 2 | 0.15627674 |

Figure 1-56. Details Regarding the Selected Model (Exponential 5) for Increased LDH Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months

1.1.1.3.2 Renal Effects

EPA identified renal endpoints in a chronic inhalation study for BMD modeling ([IRFMN, 1978](#)). Modeled results are presented for BUN and serum potassium levels in male and female rats and serum calcium levels in male rats. Modeled results are not presented for serum calcium in female rats or serum uric acid in male or female rats ([IRFMN, 1978](#)) because neither the constant nor nonconstant variance models provided adequate fit to the variance data or because none of the models provided adequate fits to the means (test 4 p-value < 0.1) assuming either constant or nonconstant variance.

1.1.1.3.2.1 Blood Urea Nitrogen (BUN) Levels in Male Sprague-Dawley Rats – 12-Month Inhalation Exposure

Serum BUN levels were significantly increased in male rats exposed to 1,2-dichloroethane by inhalation for 12 months (seven hours per day, 5 days per week) ([IRFMN, 1978](#)). The exposure concentrations (reported in ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day and 7 days per week. The concentration and response data used for the modeling are presented in Table 1-47. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)).

Table 1-47. BUN Levels in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (mg%) | SD (mg%) |
|---|-------------------|------------|----------|
| 0 | 8 | 10.29 | 0.82 |
| 4 | 8 | 11.00 | 1.50 |
| 8.3 | 8 | 10.25 | 1.27 |
| 42 | 8 | 10.63 | 1.41 |
| 126 | 8 | 15.50 | 2.06 |

The BMD modeling results for increased serum BUN levels in male rats are summarized in Table 1-48. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, all models, except for the Linear model, provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Polynomial 3-degree) was selected.

Table 1-48. Summary of BMD Modeling Results for Increased BUN Levels in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | Basis for Model Selection |
|----------------------------|-------------------------|--------------|------------------------------|-------------------------------|--|
| | Test 4 p-value | AIC | | | |
| Exponential 3 | 0.4759 | 148.3 | 89 | 50 | All models, except for the Linear model, provided adequate fit to the means (test 4 p-value > 0.1). BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | 0.2232 | 150.3 | 52 | 43 | |
| Hill | 0.2232 | 150.3 | 68 | 43 | |
| Polynomial Degree 3 | 0.6821 | 146.4 | 82 | 50 | |
| Polynomial Degree 2 | 0.5522 | 147.0 | 67 | 47 | |
| Power | 0.4760 | 148.3 | 87 | 50 | |
| Linear | 0.0473 | 152.8 | 38 | 30 | |

^a Selected model in bold.

A plot of the Polynomial 3-degree model with a BMR of one SD is shown in Figure 1-57. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-58.




Figure 1-57. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 3-Degree) for Increased BUN Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance)

Model Results

| Benchmark Dose | |
|----------------|-------------|
| BMD | 82.38167775 |
| BMDL | 49.68477751 |
| BMDU | 90.9704127 |
| AIC | 146.3563545 |
| Test 4 P-value | 0.682130008 |
| D.O.F. | 3 |

| Model Parameters | | | | |
|------------------|-------------|-------------|------------|------------|
| # of Parameters | 5 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 10.49575203 | 0.249598752 | 10.0065475 | 10.9849566 |
| beta | Bounded | NA | NA | NA |
| beta2 | Bounded | NA | NA | NA |
| beta3 | 2.5017E-06 | 2.79E-07 | 1.9552E-06 | 3.0482E-06 |
| alpha | 1.956385162 | 8.56E-01 | 0.27882579 | 3.63394453 |

| Goodness of Fit | | | | | | | | |
|-----------------|------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 8 | 10.49575203 | 10.29 | 10.29 | 1.39870839 | 0.82 | 0.82 | -0.416065723 |
| 4 | 8 | 10.49591214 | 11 | 11 | 1.39870839 | 1.5 | 1.5 | 1.019351707 |
| 8.3 | 8 | 10.49718247 | 10.25 | 10.25 | 1.39870839 | 1.27 | 1.27 | -0.499845144 |
| 42 | 8 | 10.681098 | 10.63 | 10.63 | 1.39870839 | 1.41 | 1.41 | -0.10332888 |
| 126 | 8 | 15.50009328 | 15.5 | 15.5 | 1.39870839 | 2.06 | 2.06 | -0.000188621 |

| Likelihoods of Interest | | | |
|-------------------------|-----------------|-----------------|------------|
| Model | Log Likelihood* | # of Parameters | AIC |
| A1 | -69.42787324 | 6 | 150.855746 |
| A2 | -66.18552753 | 10 | 152.371055 |
| A3 | -69.42787324 | 6 | 150.855746 |
| fitted | -70.17817725 | 3 | 146.356355 |
| R | -92.22187689 | 2 | 188.443754 |

| Tests of Interest | | | |
|-------------------|--------------------------|---------|------------|
| Test | -2*Log(Likelihood Ratio) | Test df | p-value |
| 1 | 52.07269871 | 8 | <0.0001 |
| 2 | 6.484691423 | 4 | 0.16575751 |
| 3 | 6.484691423 | 4 | 0.16575751 |
| 4 | 1.500608017 | 3 | 0.68213001 |

Figure 1-58. Details Regarding the Selected Model (Polynomial 3-Degree) for Increased BUN Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months

1.1.1.3.2.2 Blood Urea Nitrogen (BUN) in Female Sprague-Dawley Rats – 12-Month Inhalation Exposure

Serum BUN levels were significantly increased in female rats exposed to 1,2-dichloroethane by inhalation for 12 months (seven hours per day, 5 days per week) (IRFMN, 1978). The exposure concentrations (reported in ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day and 7 days per week. The concentration and response data used for the modeling are presented in Table 1-49. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012).

Table 1-49. BUN Levels in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (mg%) | SD (mg%) |
|---|-------------------|------------|----------|
| 0 | 8 | 10.50 | 0.93 |
| 4 | 8 | 10.88 | 0.99 |
| 8.3 | 8 | 10.75 | 1.50 |
| 42 | 8 | 10.72 | 2.04 |
| 126 | 8 | 15.25 | 2.38 |

The BMD modeling results for increased serum BUN levels in female rats are summarized in Table 1-50. The constant variance model did not provide adequate fit to the variance data. With the nonconstant variance model applied, all models provided an adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Polynomial 2-degree) was selected.

Table 1-50. Summary of BMD Modeling Results for Increased BUN Levels in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Nonconstant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | Basis for Model Selection |
|----------------------------|-------------------------|--------------|------------------------------|-------------------------------|--|
| | Test 4 p-value | AIC | | | |
| Exponential 3 | 0.7765 | 155.2 | 74 | 32 | All models provided an adequate fit to the means (test 4 p-value > 0.1). BMDLs differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | 0.4750 | 157.2 | 73 | 64 | |
| Hill | 0.4748 | 157.2 | 73 | 24 | |
| Polynomial Degree 3 | 0.8080 | 155.1 | 76 | 29 | |
| Polynomial Degree 2 | 0.9064 | 153.2 | 67 | 29 | |
| Power | 0.7748 | 155.2 | 73 | 29 | |
| Linear | 0.5460 | 154.8 | 38 | 25 | |

^a Selected model in bold.

A plot of the Polynomial 2-degree model with a BMR of one SD is shown in Figure 1-59. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-60.




Figure 1-59. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased BUN Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Nonconstant Variance)

Model Results

| Benchmark Dose | |
|----------------|-------------|
| BMD | 67.27795923 |
| BMDL | 28.82214651 |
| BMDU | 84.94460652 |
| AIC | 153.2223286 |
| Test 4 P-value | 0.906394034 |
| D.O.F. | 3 |

| Model Parameters | | | | |
|------------------|-------------|-------------|------------|------------|
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| # of Parameters | 5 | | | |
| g | 10.60180857 | 0.240171167 | 10.1310817 | 11.0725354 |
| beta | Bounded | NA | NA | NA |
| beta2 | 0.00028596 | 5.57E-05 | 0.00017677 | 0.00039515 |
| rho | 3.413973354 | 1.87E+00 | -0.2537519 | 7.08169862 |
| alpha | 0.000529032 | 1.28E-06 | 0.00052652 | 0.00053154 |

| Goodness of Fit | | | | | | | | |
|-----------------|------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 8 | 10.60180857 | 10.5 | 10.5 | 1.29434746 | 0.93 | 0.93 | -0.222473594 |
| 4 | 8 | 10.60638393 | 10.88 | 10.88 | 1.29530112 | 0.99 | 0.99 | 0.597469648 |
| 8.3 | 8 | 10.62150836 | 10.75 | 10.75 | 1.29845562 | 1.5 | 1.5 | 0.279893469 |
| 42 | 8 | 11.10624197 | 10.72 | 10.72 | 1.40123221 | 2.04 | 2.04 | -0.779640423 |
| 126 | 8 | 15.14170917 | 15.25 | 15.25 | 2.37839389 | 2.38 | 2.38 | 0.128781329 |

| Likelihoods of Interest | | | |
|-------------------------|-----------------|-----------------|------------|
| Model | Log Likelihood* | # of Parameters | AIC |
| A1 | -74.56612506 | 6 | 161.13225 |
| A2 | -69.31006848 | 10 | 158.620137 |
| A3 | -72.33309109 | 7 | 158.666182 |
| fitted | -72.61116432 | 4 | 153.222329 |
| R | -91.72530575 | 2 | 187.450612 |

| Tests of Interest | | | |
|-------------------|--------------------------|---------|------------|
| Test | -2*Log(Likelihood Ratio) | Test df | p-value |
| 1 | 44.83047455 | 8 | <0.0001 |
| 2 | 10.51211316 | 4 | 0.03263054 |
| 3 | 6.046045235 | 3 | 0.10939139 |
| 4 | 0.556146456 | 3 | 0.90639403 |

Figure 1-60. Details Regarding the Selected Model (Polynomial 2-Degree) for Increased BUN Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months

1.1.1.3.2.3 Calcium Levels in Male Sprague-Dawley Rats – 12-Month Inhalation Exposure

Serum calcium levels were significantly decreased in male rats exposed to 1,2-dichloroethane by inhalation for 12 months (seven hours per day, 5 days per week) ([IRFMN, 1978](#)). The exposure concentrations (reported in ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day and 7 days per week. The concentration and response data used for the modeling are presented in Table 1-51. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). A BMR of 10% RD was also selected because EPA considers a 10% change in serum calcium levels to be biologically relevant.

Table 1-51. Serum Calcium Levels in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (mg%) | SD (mg%) |
|---|-------------------|------------|----------|
| 0 | 8 | 9.98 | 0.37 |
| 4 | 8 | 9.63 | 0.31 |
| 8.3 | 8 | 9.48 | 0.31 |
| 42 | 8 | 8.95 | 0.14 |
| 126 | 8 | 8.73 | 0.37 |

The BMD modeling results for decreased serum calcium levels in male rats are summarized in Table 1-52. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, the Exponential 5 and Hill models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Hill model) was selected.

Table 1-52. Summary of BMD Modeling Results for Decreased Serum Calcium Levels in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 10%RD (mg/m ³) | BMDL 10%RD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|--------------|------------------------------|-------------------------------|--------------------------------|---------------------------------|---|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.0004 | 39.30 | 42 | 32 | 110 | 91 | The Exponential 5 and Hill models provided adequate fit to the means (test 4 p-value > 0.1). BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | 0.5273 | 24.18 | 5.2 | 2.6 | 34 | 16 | |
| Hill | 0.9146 | 23.08 | 3.8 | 2.0 | 38 | 18 | |
| Polynomial Degree 3 | 0.0003 | 39.91 | 44 | 34 | 120 | 94 | |
| Polynomial Degree 2 | 0.0003 | 39.91 | 44 | 34 | 120 | 94 | |
| Power | 0.0003 | 39.91 | 44 | 34 | 120 | 94 | |
| Linear | 0.0003 | 39.91 | 44 | 34 | 120 | 94 | |

^a Selected model in bold.

Plots of the Hill model with BMRs of one SD and 10% RD are shown in Figure 1-61 and Figure 1-62, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-63 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-61. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Decreased Serum Calcium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance)




Figure 1-62. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Decreased Serum Calcium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 10%RD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 3.793853645 | | | | | | | |
| BMDL | 1.973102694 | | | | | | | |
| BMDU | 9.002210197 | | | | | | | |
| AIC | 23.07602981 | | | | | | | |
| Test 4 P-value | 0.916438367 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 5 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 9.968378751 | 9.97E-02 | 9.77293889 | 10.1638186 | | | | |
| v | -1.367698827 | 1.45E-01 | -1.6527488 | -1.0826488 | | | | |
| k | 13.96705849 | 6.12E+00 | 1.9724673 | 25.9616497 | | | | |
| n | Bounded | NA | NA | NA | | | | |
| alpha | 0.08535158 | 1.63E-03 | 0.0821589 | 0.08854426 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 8 | 9.968378751 | 9.98 | 9.98 | 0.29214993 | 0.37 | 0.37 | 0.112510234 |
| 4 | 8 | 9.663888435 | 9.63 | 9.63 | 0.29214993 | 0.31 | 0.31 | -0.328088289 |
| 8.3 | 8 | 9.458571842 | 9.48 | 9.48 | 0.29214993 | 0.31 | 0.31 | 0.207455072 |
| 42 | 8 | 8.942000873 | 8.95 | 8.95 | 0.29214993 | 0.14 | 0.14 | 0.077442935 |
| 126 | 8 | 8.737160105 | 8.73 | 8.73 | 0.29214993 | 0.37 | 0.37 | -0.069320011 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -7.450754442 | 6 | 26.9015089 | | | | | |
| A2 | -3.711046548 | 10 | 27.4220931 | | | | | |
| A3 | -7.450754442 | 6 | 26.9015089 | | | | | |
| fitted | -7.538014904 | 4 | 23.0760298 | | | | | |
| R | -32.1569544 | 2 | 68.3139088 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 56.89181569 | 8 | <0.0001 | | | | | |
| 2 | 7.479415788 | 4 | 0.1126204 | | | | | |
| 3 | 7.479415788 | 4 | 0.1126204 | | | | | |
| 4 | 0.174520924 | 2 | 0.91643837 | | | | | |

Figure 1-63. Details Regarding the Selected Model (Hill) for Increased Serum Calcium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months

1.1.1.3.2.4 Serum Potassium Levels in Male Sprague-Dawley Rats – 12-Month Inhalation Exposure

Serum potassium levels were significantly increased in male rats exposed to 1,2-dichloroethane by inhalation for 12 months (seven hours per day, 5 days per week) (IRFMN, 1978). The exposure concentrations (reported in ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day and 7 days per week. The concentration and response data used for the modeling are presented in Table 1-53. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected because EPA considers a 10% change in serum potassium levels to be biologically relevant.

Table 1-53. Serum Potassium Levels in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (mg%) | SD (mg%) |
|---|-------------------|------------|----------|
| 0 | 8 | 4.91 | 0.57 |
| 4 | 8 | 5.34 | 0.48 |
| 8.3 | 8 | 6.44 | 0.71 |
| 42 | 8 | 6.08 | 0.71 |
| 126 | 8 | 6.26 | 0.59 |

The BMD modeling results for increased serum potassium levels in male rats are summarized in Table 1-54. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, the Exponential 5 and Hill models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Hill) was selected.

Table 1-54. Summary of BMD Modeling Results for Increased Serum Potassium Levels in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 10%RD (mg/m ³) | BMDL 10%RD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|--------------|------------------------------|-------------------------------|--------------------------------|---------------------------------|---|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | - | - | - | - | - | - | The Exponential 5 and Hill models provided adequate fit to the means (test 4 p-value > 0.1). BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | 0.2176 | 81.23 | 4.1 | 3.0 | 4.0 | 3.8 | |
| Hill | 0.4676 | 79.23 | 4.1 | 3.6 | 4.0 | 3.8 | |
| Polynomial Degree 3 | < 0.0001 | 97.90 | 119 | 71 | 87 | 51 | |
| Polynomial Degree 2 | < 0.0001 | 97.90 | 119 | 71 | 87 | 51 | |
| Power | < 0.0001 | 97.90 | 119 | 71 | 87 | 51 | |

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 10%RD (mg/m ³) | BMDL 10%RD (mg/m ³) | Basis for Model Selection |
|--------|-------------------------|-------|------------------------------|-------------------------------|--------------------------------|---------------------------------|---------------------------|
| | Test 4 p-value | AIC | | | | | |
| Linear | < 0.0001 | 97.90 | 119 | 71 | 87 | 51 | |

^a Selected model in bold.

Plots of the Hill model with BMRs of one SD and 10% RD are shown in Figure 1-64 and Figure 1-65, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-66 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-64. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Increased Serum Potassium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance)




Figure 1-65. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Increased Serum Potassium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 10%RD (Constant Variance)

Model Results

| Benchmark Dose | |
|----------------|-------------|
| BMD | 4.113998297 |
| BMDL | 3.556136042 |
| BMDU | 6.037422863 |
| AIC | 79.23435408 |
| Test 4 P-value | 0.467568201 |
| D.O.F. | 2 |

| Model Parameters | | | | |
|------------------|-------------|-------------|------------|------------|
| # of Parameters | 5 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 4.909998383 | 0.208411252 | 4.50151983 | 5.31847693 |
| v | 1.350002887 | 0.240652145 | 0.87833335 | 1.82167243 |
| k | 4.172638129 | 0.201800737 | 3.77711595 | 4.56816031 |
| n | Bounded | NA | NA | NA |
| alpha | 0.347490412 | 0.026999944 | 0.29457149 | 0.40040933 |

| Goodness of Fit | | | | | | | | |
|-----------------|------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 8 | 4.909998383 | 4.91 | 4.91 | 0.58948317 | 0.57 | 0.57 | 7.76078E-06 |
| 4 | 8 | 5.340003496 | 5.34 | 5.34 | 0.58948317 | 0.48 | 0.48 | -1.67735E-05 |
| 8.3 | 8 | 6.25999559 | 6.44 | 6.44 | 0.58948317 | 0.71 | 0.71 | 0.863687681 |
| 42 | 8 | 6.26000127 | 6.08 | 6.08 | 0.58948317 | 0.71 | 0.71 | -0.863672616 |
| 126 | 8 | 6.26000127 | 6.26 | 6.26 | 0.58948317 | 0.59 | 0.59 | -6.09342E-06 |

| Likelihoods of Interest | | | |
|-------------------------|-----------------|-----------------|------------|
| Model | Log Likelihood* | # of Parameters | AIC |
| A1 | -34.85696698 | 6 | 81.713934 |
| A2 | -34.01730185 | 10 | 88.0346037 |
| A3 | -34.85696698 | 6 | 81.713934 |
| fitted | -35.61717704 | 4 | 79.2343541 |
| R | -48.90125201 | 2 | 101.802504 |

| Tests of Interest | | | |
|-------------------|--------------------------|---------|------------|
| Test | -2*Log(Likelihood Ratio) | Test df | p-value |
| 1 | 29.76790032 | 8 | 0.0002323 |
| 2 | 1.679330262 | 4 | 0.79446879 |
| 3 | 1.679330262 | 4 | 0.79446879 |
| 4 | 1.520420112 | 2 | 0.4675682 |

Figure 1-66. Details Regarding the Selected Model (Hill) for Increased Serum Potassium Levels in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months

1.1.1.3.2.5 Serum Potassium Levels in Female Sprague-Dawley Rats – 12-Month Inhalation Exposure

Serum potassium levels were significantly increased in female rats exposed to 1,2-dichloroethane by inhalation for 12 months (seven hours per day, 5 days per week) (IRFMN, 1978). The exposure concentrations (reported in ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day and 7 days per week. The concentration and response data used for the modeling are presented in Table 1-55. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected because EPA considers a 10% change in serum potassium levels to be biologically relevant.

Table 1-55. Serum Potassium Levels in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 12-Month Inhalation Exposure Study

| Adjusted Concentration (mg/m ³) | Number of Animals | Mean (mg%) | SD (mg%) |
|---|-------------------|------------|----------|
| 0 | 8 | 4.97 | 0.76 |
| 4 | 8 | 5.61 | 1.10 |
| 8.3 | 8 | 6.64 | 0.74 |
| 42 | 8 | 6.19 | 0.57 |
| 126 | 8 | 6.10 | 0.71 |

The BMD modeling results for increased serum potassium levels in female rats are summarized in Table 1-56. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, the Exponential 5 and Hill models provided adequate fit to the means (test 4 p-value > 0.1). At a BMR of one SD, the BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Hill) was selected. At a BMR of 10% RD, the BMDLs for the fit models were not sufficiently close (differed by > 3-fold); therefore, BMDS recommended the model with the lowest BMDL (Exponential 5). The Hill model was selected, however, because it has a lower AIC, has an estimated BMD/BMDL ratio within 3-fold, BMDL within 3-fold of lowest non-zero concentration, and is consistent with the selection for the one SD BMR.

Table 1-56. Summary of BMD Modeling Results for Increased Serum Potassium Levels in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane for 12 Months^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 10%RD (mg/m ³) | BMDL 10%RD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|--------------|------------------------------|-------------------------------|--------------------------------|---------------------------------|--|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | - | - | - | - | - | - | The Exponential 5 and Hill models provided adequate fit to the means (test 4 p-value > 0.1). At a BMR of one SD, BMDLs for the fit models differed by < 3-fold; therefore, the model with the lowest AIC was selected (Hill). At a BMR of 10% RD, the BMDLs for the fit models differed by > 3-fold; therefore, BMDS recommended the model with the lowest BMDL (Exponential 5). The Hill model was selected, however, because it has a lower AIC, has a BMD/BMDL ratio within 3-fold, BMDL within 3-fold of lowest non-zero concentration, and is consistent with the selection for the one SD BMR. |
| Exponential 5 | 0.1254 | 102.2 | 4.2 | 1.9 | 3.7 | 1.0 | |
| Hill | 0.3091 | 100.2 | 4.1 | 3.2 | 3.9 | 3.6 | |
| Polynomial Degree 3 | 0.0007 | 112.8 | 250 | 100 | 160 | 63 | |
| Polynomial Degree 2 | 0.0007 | 112.8 | 250 | 100 | 160 | 63 | |
| Power | 0.0007 | 112.8 | 250 | 100 | 150 | 63 | |
| Linear | 0.0007 | 112.8 | 250 | 100 | 150 | 63 | |

^a Selected model in bold.

Plots of the Hill model with BMRs of one SD and 10% RD are shown in Figure 1-67 and Figure 1-68, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-69 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-67. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Increased Serum Potassium Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 1SD (Constant Variance)




Figure 1-68. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Increased Serum Potassium Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months and BMR of 10%RD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 4.085130438 | | | | | | | |
| BMDL | 3.209977644 | | | | | | | |
| BMDU | 5.384570286 | | | | | | | |
| AIC | 100.2205383 | | | | | | | |
| Test 4 P-value | 0.309094341 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 5 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 4.969999172 | 0.270915253 | 4.43901503 | 5.50098332 | | | | |
| v | 1.340001086 | 0.312822339 | 0.72688056 | 1.95312161 | | | | |
| k | 4.01996137 | 0.210192831 | 3.60799099 | 4.43193175 | | | | |
| n | Bounded | NA | NA | NA | | | | |
| alpha | 0.587215363 | 7.71E-02 | 0.43611298 | 0.73831774 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 8 | 4.969999172 | 4.97 | 4.97 | 0.76629979 | 0.76 | 0.76 | 3.05499E-06 |
| 4 | 8 | 5.610002828 | 5.61 | 5.61 | 0.76629979 | 1.1 | 1.1 | -1.04376E-05 |
| 8.3 | 8 | 6.309997376 | 6.64 | 6.64 | 0.76629979 | 0.74 | 0.74 | 1.218045977 |
| 42 | 8 | 6.310000258 | 6.19 | 6.19 | 0.76629979 | 0.57 | 0.57 | -0.442923242 |
| 126 | 8 | 6.310000258 | 6.1 | 6.1 | 0.76629979 | 0.71 | 0.71 | -0.775114958 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -44.93616041 | 6 | 101.872321 | | | | | |
| A2 | -43.00818559 | 10 | 106.016371 | | | | | |
| A3 | -44.93616041 | 6 | 101.872321 | | | | | |
| fitted | -46.11026914 | 4 | 100.220538 | | | | | |
| R | -54.15242891 | 2 | 112.304858 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 22.28848664 | 8 | 0.00440857 | | | | | |
| 2 | 3.855949636 | 4 | 0.42585182 | | | | | |
| 3 | 3.855949636 | 4 | 0.42585182 | | | | | |
| 4 | 2.348217473 | 2 | 0.30909434 | | | | | |

Figure 1-69. Details Regarding the Selected Model (Hill) for Increased Serum Potassium Levels in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation for 12 Months

1.1.1.4 Developmental

1.1.1.4.1 Developmental Effects

Male pup weight at weaning (~21 days of age) was nonsignificantly decreased by ≥ 5 percent at the highest tested concentration in a one-generation reproduction study of 1,2-dichloroethane inhalation in rats when the data were limited to a very small subset of available pups (five pups from different F1b litters selected for organ weight measurements) (Rao et al., 1980). There was no decrease relative to controls in the corresponding selected F1b female pups. Overall, pup body weight data from all pups of both sexes in F1a and F1b litters recorded from birth through weaning were reported by the study authors not to show any differences from controls (data for whole group were not presented in the study).

1.1.1.4.1.1 Body Weight in Male Weanling F1_B Rats

The data for body weight at weaning of the selected male F1b pups were modeled (Rao et al., 1980). The parental exposure concentrations (reported in ppm) were converted to units of mg/m³ and duration adjusted to estimate an equivalent time-weighted average (TWA) inhalation concentration for parental animals (exposure was 6 hours per day, 5 days per week for 60 days prior to mating and then 7 days per week for 116 days, except that maternal exposure was stopped to allow for delivery and rearing of the young from GD 21 to postnatal day 4). The concentration and response data used for the modeling are presented in Table 1-57. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). BMRs of 5% and 10% RD were also selected because EPA considers these BMRs to be biologically relevant for pup body weight change in a reproduction study.

Table 1-57. Body Weight of Selected F1b Male Weanling Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a One-Generation Developmental Study^a

| TWA Concentration (mg/m ³) | Number of Animals | Mean (g) | SD (g) |
|--|-------------------|----------|--------|
| 0 | 5 | 42 | 9 |
| 23 | 5 | 42 | 6 |
| 68 | 5 | 40 | 5 |
| 137 | 5 | 36 | 6 |

^a Weanling body weight data presented are only for the small subset limited to male weanlings from F1B litters that were selected for organ weight measurements.

The BMD modeling results for body weight of selected F1b male rats at weaning are summarized in Table 1-58. The test for significant difference in responses and variances failed (test 1 p-value > 0.05), indicating that there is no clear dose-response present in the data. This means the additional modeling results presented here are suspect and should be interpreted with caution. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, all models, except for the Exponential 5 and Hill models, provided adequate fit to the means (test 4 p-value < 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Linear model) was selected.

Table 1-58. Summary of BMD Modeling Results for Decreased Body Weight of Selected F1b Male Weanling Rats Following Inhalation Exposure to 1,2-Dichloroethane in a One-Generation Reproduction Study (Constant Variance)^a

| Model | Goodness of Fit (Means) | | BMD 1SD (mg/m ³) | BMDL 1SD (mg/m ³) | BMD 5%RD (mg/m ³) | BMDL 5%RD (mg/m ³) | BMD 10%RD (mg/m ³) | BMDL 10%RD (mg/m ³) | Basis for Model Selection |
|---------------------|-------------------------|--------------|------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|---------------------------------|--|
| | Test 4 p-value | AIC | | | | | | | |
| Exponential 3 | 0.9195 | 136.2 | 130 | 110 | 69 | 23 | 110 | 46 | All models, except for the Exponential 5 and Hill models, provided adequate fit to the means (test 4 p-value > 0.1). BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. NOTE: This data set (small subset of the available data) is not representative of larger results in this study and BMD modeling no showed no clear dose-response present in these selected data. Although BMDs and BMDLs are presented here, they are suspect and should be interpreted with caution. |
| Exponential 5 | NA | 138.2 | 95 | 35 | 69 | 2 | 78 | 10 | |
| Hill | NA | 138.2 | 130 | 24 | 69 | 16 | 82 | 21 | |
| Polynomial Degree 3 | 0.8615 | 136.2 | 130 | 67 | 70 | 25 | 110 | 51 | |
| Polynomial Degree 2 | 0.8932 | 136.2 | 130 | 67 | 71 | 25 | 110 | 51 | |
| Power | 0.9144 | 136.2 | 130 | 67 | 69 | 25 | 110 | 51 | |
| Linear | 0.9366 | 134.3 | 130 | 67 | 46 | 25 | 93 | 50 | |

^a Selected model in bold.

Plots of the Linear model with BMRs of one SD, 5% RD, and 10% RD are shown in Figure 1-70, Figure 1-71, and Figure 1-72, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-73 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to all BMRs).




Figure 1-70. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Decreased Body Weight of Selected F1B Male Weanling Rats Exposed to 1,2-Dichloroethane Via Inhalation in a One-Generation Reproduction Study and BMR of 1SD (Constant Variance)




Figure 1-71. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Decreased Body Weight of Selected F1B Male Weanling Rats Exposed to 1,2-Dichloroethane Via Inhalation in a One-Generation Reproduction Study and BMR of 5%RD (Constant Variance)




Figure 1-72. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Decreased Body Weight of Selected F1B Male Weanling Rats Exposed to 1,2-Dichloroethane Via Inhalation in a One-Generation Reproduction Study and BMR of 10%RD (Constant Variance)

| Model Results | | | | | | | | |
|-------------------------|--------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 130.2879275 | | | | | | | |
| BMDL | 66.67084085 | | | | | | | |
| BMDU | 2727.30983 | | | | | | | |
| AIC | 134.3353986 | | | | | | | |
| Test 4 P-value | 0.936625044 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 3 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 42.61889064 | 1.980363222 | 38.73745 | 46.5003313 | | | | |
| beta | -0.045945453 | 2.56E-02 | -0.096131 | 0.00424011 | | | | |
| alpha | 35.83384612 | 4.06E+02 | -760.01881 | 831.686503 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 5 | 42.61889064 | 42 | 42 | 5.98613783 | 9 | 9 | -0.231181035 |
| 23 | 5 | 41.56214522 | 42 | 42 | 5.98613783 | 6 | 6 | 0.163556716 |
| 68 | 5 | 39.49459984 | 40 | 40 | 5.98613783 | 5 | 5 | 0.188787687 |
| 137 | 5 | 36.32436359 | 36 | 36 | 5.98613783 | 6 | 6 | -0.121163103 |

| Likelihoods of Interest | | | |
|-------------------------|-----------------|-----------------|------------|
| Model | Log Likelihood* | # of Parameters | AIC |
| A1 | -64.10222704 | 5 | 138.204454 |
| A2 | -63.09824229 | 8 | 142.196485 |
| A3 | -64.10222704 | 5 | 138.204454 |
| fitted | -64.16769929 | 3 | 134.335399 |
| R | -65.65977234 | 2 | 135.319545 |

| Tests of Interest | | | |
|-------------------|--------------------------|---------|------------|
| Test | -2*Log(Likelihood Ratio) | Test df | p-value |
| 1 | 5.123060089 | 6 | 0.52812954 |
| 2 | 2.007969501 | 3 | 0.57075425 |
| 3 | 2.007969501 | 3 | 0.57075425 |
| 4 | 0.130944487 | 2 | 0.93662504 |

Figure 1-73. Details Regarding the Selected Model (Linear) for Decreased Body Weight of Selected F1B Male Weanling Rats Exposed to 1,2-Dichloroethane Via Inhalation in a One-Generation Reproduction Study

1.1.2 Oral Data

1.1.2.1 Acute

1.1.2.1.1 Mortality

[Storer et al. \(1984\)](#) provided data showing increased mortality in male mice following an acute oral exposure to 1,2-dichloroethane.

1.1.2.1.1.1 Mortality in Male B6C3F1 Mice – Single Oral Gavage

There was an increased incidence of mortality in male mice exposed to 1,2-dichloroethane following a single oral gavage exposure in an acute toxicity study by [Storer et al. \(1984\)](#). The dose and response data used for the modeling are presented in Table 1-59. Dichotomous models were fit to the dose-response data.

A BMR of 10% ER was chosen according to *BMD Technical Guidance* ([U.S. EPA, 2012](#)). A BMR of one percent ER was selected due to severity of the endpoint and a BMR of 20 percent ER was selected because it is near the low end of the observable range in the study.

Table 1-59. Incidence of Mortality in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Acute Oral Exposure Study

| Dose (mg/kg) | Number of Animals | Incidence |
|--------------|-------------------|-----------|
| 0 | 5 | 0 |
| 200 | 5 | 0 |
| 300 | 5 | 0 |
| 400 | 5 | 2 |
| 500 | 5 | 4 |
| 600 | 5 | 4 |

The BMD modeling results for increased incidence of mortality in male rats are summarized in Table 1-60. With the BMRs of 10 and 20 percent applied, all models provided adequate fit to the data (chi-square p-value > 0.1). Despite the overall adequate fit, the Multistage 1-degree model was not considered further because this model provided poor fit in the lower portion of the dose-response curve, with scaled residuals of -1.3 and -1.6 at the two lowest doses. The BMDLs for the remaining models differed by < 3-fold and were considered sufficiently close; therefore, the model with the lowest AIC (Multistage 3-degree) was selected. Model outputs could not be generated by the BMDS using a BMR of one percent ER.

Table 1-60. Summary of BMD Modeling Results for Increased Incidence of Mortality in Male Mice Following a Single Oral Exposure to 1,2-Dichloroethane Using BMR of 10%ER or 20%ER^a

| Model | Goodness of Fit | | BMD 10%ER (mg/kg) | BMDL 10%ER (mg/kg) | BMD 20%ER (mg/kg) | BMDL 20%ER (mg/kg) | Basis for Model Selection |
|---------------------|-----------------|--------------|-------------------|--------------------|-------------------|--------------------|---|
| | P-value | AIC | | | | | |
| Dichotomous Hill | 0.9999 | 20.79 | 360 | 257 | 377 | 303 | All models provided adequate fit to the data (chi-square p-value > 0.1). Despite the overall adequate fit, the Multistage 1-degree model was not considered further because this model provided poor fit in the lower portion of the dose-response curve, with scaled residuals of -1.3 and -1.6 at the two lowest doses. The BMDLs for the remaining models differed by less than 3-fold and were considered sufficiently close; therefore, EPA chose the model with the lowest AIC (Multistage 3-degree). |
| Gamma | 0.8556 | 22.30 | 323 | 216 | 363 | 273 | |
| Log-Logistic | 0.8800 | 22.15 | 325 | 222 | 364 | 277 | |
| Multistage 3 | 0.9027 | 20.21 | 239 | 129 | 307 | 224 | |
| Multistage 2 | 0.5828 | 24.55 | 174 | 90.9 | 254 | 170 | |
| Multistage 1 | 0.2148 | 28.84 | 74.6 | 45.6 | 158 | 96.5 | |
| Weibull | 0.8616 | 21.17 | 301 | 184 | 356 | 254 | |
| Logistic | 0.7493 | 22.94 | 316 | 199 | 366 | 275 | |
| Log-Probit | 0.8880 | 22.07 | 326 | 227 | 363 | 277 | |
| Probit | 0.7659 | 22.86 | 318 | 195 | 365 | 270 | |
| Quantal Linear | 0.2148 | 28.84 | 74.6 | 45.6 | 158 | 96.5 | |

^a Selected model in bold.

Plots of the Multistage 3-degree model with BMRs of 10% ER and 20 percent ER are shown in Figure 1-74 and Figure 1-75, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-76 (BMD and BMDL shown are for BMR of 10% RD; the rest is applicable to both BMRs).




Figure 1-74. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Incidence of Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 10%ER




Figure 1-75. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Incidence of Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 20%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 238.9043391 | | | | |
| BMDL | 129.2249981 | | | | |
| BMDU | 291.4737157 | | | | |
| AIC | 20.20936618 | | | | |
| P-value | 0.902688778 | | | | |
| D.O.F. | 6 | | | | |
| Chi ² | 2.177325909 | | | | |
| Slope Factor | 0.000773844 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 4 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | Bounded | NA | NA | NA | |
| b2 | Bounded | NA | NA | NA | |
| b3 | Bounded | NA | NA | NA | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-08 | 0 | 5 | -0.000276 |
| 200 | 0.059943491 | 0.299717454 | 0 | 5 | -0.56465 |
| 300 | 0.188301704 | 0.941508522 | 0 | 5 | -1.076998 |
| 400 | 0.390137775 | 1.950688873 | 2 | 5 | 0.0452101 |
| 500 | 0.619345703 | 3.096728513 | 4 | 5 | 0.8319579 |
| 600 | 0.811566932 | 4.057834662 | 4 | 5 | -0.06614 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -8.36908257 | 6 | - | - | NA |
| Fitted Model | -10.10468309 | 0 | 3.47120104 | 6 | 0.7477972 |
| Reduced Model | -19.09542505 | 1 | 21.452685 | 5 | 0.0006651 |

Figure 1-76. Details Regarding the Selected Model (Multistage 3-Degree) for Increased Incidence of Mortality in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study

1.1.2.1.2 Hepatic Effects

EPA identified hepatic endpoints in an acute oral gavage study in male mice for BMD modeling ([Storer et al., 1984](#)). Modeled results are presented for relative liver weight. Modeled results are not presented for serum ALT or LDH because neither the constant nor nonconstant variance models provided adequate fit to the variance data or because none of the models provided adequate fits to the means (test 4 p-value < 0.1) and viable results assuming either constant or nonconstant variance.

1.1.2.1.2.1 Relative Liver Weight in Male B6C3F1 Mice – Single Oral Gavage

Relative liver weights were significantly increased in male mice exposed to 1,2-dichloroethane following a single oral gavage exposure in an acute toxicity study by [Storer et al. \(1984\)](#). The dose and response data used for the modeling are presented in Table 1-61. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). A BMR of 10% RD was also selected because EPA considers a 10% change in relative liver weight to be biologically significant.

Table 1-61. Relative Liver Weight in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Acute Oral Exposure Study

| Dose (mg/kg) | Number of Animals | Mean (Liver Weight/100 g Body Weight) | SD (Liver Weight/100 g Body Weight) |
|--------------|-------------------|---------------------------------------|-------------------------------------|
| 0 | 5 | 4.26 | 0.14 |
| 200 | 5 | 4.21 | 0.27 |
| 300 | 5 | 4.47 | 0.20 |
| 400 | 3 | 5.10 | 0.58 |

The BMD modeling results for increased relative liver weight in male mice are summarized in Table 1-62. The constant variance model did not provide adequate fit to the variance data, but the nonconstant variance model did. With the nonconstant variance model applied and using a BMR of one SD, the Exponential 3, Exponential 5, and Power models provided an adequate fit to the means (test 4 p-value > 0.1). BMD computations failed for the Polynomial 2- and 3-degree and Linear models. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Exponential 5). When a BMR of 10% RD was applied, the same models plus the Polynomial 3-degree provided an adequate fit to the means (test 4 p-value > 0.1). The BMDLs among these models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Polynomial 3-degree).

Table 1-62. Summary of BMD Modeling Results for Increased Relative Liver Weight in Male Mice Following a Single Oral Exposure to 1,2-Dichloroethane (Nonconstant Variance)^a

| Model | Goodness of Fit | | BMD 1SD (mg/kg) | BMDL 1SD (mg/kg) | BMD 10%R D (mg/kg) | BMDL 10%R D (mg/kg) | Basis for Model Selection |
|----------------------------|-------------------|--------------|-----------------------|------------------------|-----------------------------|------------------------------|---|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.7631 | 7.354 | 290 | 220 | 345 | 300 | Using a BMR of one SD, the Exponential 3, Exponential 5, and Power models provided an adequate fit to the means (test 4 p-value > 0.1). BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC (Exponential 5). Using a BMR of 10% RD, the same models plus the Polynomial 3-degree provided an adequate fit to the means (test 4 p-value > 0.1). BMDLs among these models differed by < 3-fold; therefore, the EPA chose the model with the lowest AIC (Polynomial 3-degree). |
| Exponential 5 | 0.9858 | 7.263 | 297 | 272 | 315 | 298 | |
| Hill | NA | 9.263 | 297 | 278 | 319 | 310 | |
| Polynomial Degree 3 | 0.6308 | 4.991 | - | - | 326 | 285 | |
| Polynomial Degree 2 | 0.0965 | 9.939 | - | - | 313 | 259 | |
| Power | 0.7732 | 7.346 | 290 | 252 | 345 | 300 | |
| Linear | 0.0073 | 15.10 | - | - | 307 | 209 | |

^a Selected model in bold.

A Plot of the Exponential 5 model with a BMR of one SD is shown in Figure 1-77 and a plot of the Polynomial 3-degree model with a BMR of 10% RD is shown in Figure 1-78. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood for the Exponential 5 model with BMR of one SD are shown in Figure 1-79 and additional modeling details for the Polynomial 3-degree model with BMR of 10% RD are shown in Figure 1-80.




Figure 1-77. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 5) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 1SD (Nonconstant Variance)




Figure 1-78. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 3-Degree) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 10%RD (Nonconstant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 297.3046064 | | | | | | | |
| BMDL | 271.7913769 | | | | | | | |
| BMDU | 334.4904423 | | | | | | | |
| AIC | 7.263445078 | | | | | | | |
| Test 4 P-value | 0.985787525 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 6 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| a | 4.239995767 | 5.76E-02 | 4.12701127 | 4.35298026 | | | | |
| b | 0.003099096 | 1.10E-04 | 0.00288322 | 0.00331498 | | | | |
| c | 1.209416148 | 5.05E-02 | 1.11038106 | 1.30845124 | | | | |
| d | Bounded | NA | NA | NA | | | | |
| rho | 9.32012481 | 5.00E+00 | -0.4827733 | 19.123023 | | | | |
| log-alpha | -16.87383537 | 7.44E+00 | -31.448574 | -2.2990966 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 5 | 4.239995767 | 4.26 | 4.26 | 0.18174197 | 0.14 | 0.14 | 0.246122706 |
| 200 | 5 | 4.240157615 | 4.21 | 4.21 | 0.1817743 | 0.27 | 0.27 | -0.370979159 |
| 300 | 5 | 4.449696613 | 4.47 | 4.47 | 0.22759008 | 0.2 | 0.2 | 0.199480371 |
| 400 | 3 | 5.127919347 | 5.1 | 5.1 | 0.44082218 | 0.58 | 0.58 | -0.109698943 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -1.189417326 | 5 | 12.3788347 | | | | | |
| A2 | 2.79948267 | 8 | 10.4010347 | | | | | |
| A3 | 1.368436124 | 6 | 9.26312775 | | | | | |
| fitted | 1.368277461 | 5 | 7.26344508 | | | | | |
| R | -9.233982277 | 2 | 22.4679646 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 24.06692989 | 6 | 0.00050766 | | | | | |
| 2 | 7.977799991 | 3 | 0.04647275 | | | | | |
| 3 | 2.862093091 | 2 | 0.23905861 | | | | | |
| 4 | 0.000317326 | 1 | 0.98578752 | | | | | |

Figure 1-79. Details Regarding the Selected Model (Exponential 5) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study with a BMR of 1SD

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 325.7980347 | | | | | | | |
| BMDL | 284.922869 | | | | | | | |
| BMDU | 377.6371666 | | | | | | | |
| AIC | 4.990841016 | | | | | | | |
| Test 4 P-value | 0.630790181 | | | | | | | |
| D.O.F. | 3 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 6 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 4.190234558 | 0.058404416 | 4.07576401 | 4.30470511 | | | | |
| beta | Bounded | NA | NA | NA | | | | |
| beta2 | Bounded | NA | NA | NA | | | | |
| beta3 | Bounded | NA | NA | NA | | | | |
| rho | 10.15179628 | 0.240897127 | 9.67964658 | 10.623946 | | | | |
| alpha | 1.523E-08 | 1.83E-20 | 1.523E-08 | 1.523E-08 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 5 | 4.190234558 | 4.26 | 4.26 | 0.177733 | 0.14 | 0.14 | 0.877722612 |
| 200 | 5 | 4.287170082 | 4.21 | 4.21 | 0.1996107 | 0.27 | 0.27 | -0.864470458 |
| 300 | 5 | 4.517391953 | 4.47 | 4.47 | 0.26031146 | 0.2 | 0.2 | -0.407095508 |
| 400 | 3 | 4.965718753 | 5.1 | 5.1 | 0.42080801 | 0.58 | 0.58 | 0.552703217 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -1.189417326 | 5 | 12.3788347 | | | | | |
| A2 | 2.79948267 | 8 | 10.4010347 | | | | | |
| A3 | 1.368436124 | 6 | 9.26312775 | | | | | |
| fitted | 0.504579492 | 3 | 4.99084102 | | | | | |
| R | -9.233982277 | 2 | 22.4679646 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 24.06692989 | 6 | 0.00050766 | | | | | |
| 2 | 7.977799991 | 3 | 0.04647275 | | | | | |
| 3 | 2.862093091 | 2 | 0.23905861 | | | | | |
| 4 | 1.727713264 | 3 | 0.63079018 | | | | | |

Figure 1-80. Details Regarding the Selected Model (Polynomial 3-Degree) for Increased Relative Liver Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study with a BMR of 10%RD

1.1.2.1.3 Renal Effects

For acute oral exposure, EPA selected two renal endpoints for quantitative dose-response analysis with BMDS, including relative kidney weights and BUN levels following a single oral gavage exposure in male mice (Storer et al., 1984). For both data sets, only data for the control and three lowest doses (0, 200, 300, and 400 mg/kg) were modeled due to high mortality (four of five) at the two highest tested doses (500 and 600 mg/kg). EPA modeled relative kidney weight change because a statistically and biologically significant change was identified. Though not statistically significant, EPA modeled BUN in male mice from the acute oral Storer et al. (1984) study because a dose-related trend was evident in the data.

1.1.2.1.3.1 Relative Kidney Weight in Male B6C3F1 Mice – Single Oral Gavage

Relative kidney weight was significantly increased in male mice exposed to 1,2-dichloroethane following a single oral gavage exposure in an acute toxicity study by Storer et al. (1984). The dose and response data used for the modeling are presented in Table 1-63. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected because EPA considers a 10% change in relative kidney weight to be biologically significant.

Table 1-63. Increased Relative Kidney Weight in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Oral Exposure Study

| Dose (mg/kg) | Number of Animals | Mean (Kidney Weight/100 g Body Weight) | SD (Kidney Weight/100 g Body Weight) |
|--------------|-------------------|--|--------------------------------------|
| 0 | 5 | 1.50 | 0.09 |
| 200 | 5 | 1.58 | 0.19 |
| 300 | 5 | 1.69 | 0.09 |
| 400 | 3 | 1.75 | 0.08 |

The BMD modeling results for increased relative kidney weight in male mice are summarized in Table 1-64. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, all models except Exponential 5 and Hill provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Polynomial 2-degree) was selected.

Table 1-64. Summary of BMD Modeling Results for Increased Relative Kidney Weight in Male Mice Following a Single Oral Exposure to 1,2-Dichloroethane (Constant Variance)^a

| Model | Goodness of Fit | | BMD 1SD (mg/kg) | BMDL 1SD (mg/kg) | BMD 10%RD (mg/kg) | BMDL 10%RD (mg/kg) | Basis for Model Selection |
|----------------------------|-----------------|---------------|-----------------------|------------------------|-------------------------|--------------------------|--|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.6040 | -19.77 | 221 | 124 | 271 | 162 | All models except Exponential 5 and Hill provided adequate fit to the means (test 4 p-value > 0.1). BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | -18.04 | 228 | 94.9 | 262 | 128 | |
| Hill | NA | -18.04 | 226 | 89.1 | 260 | 124 | |
| Polynomial Degree 3 | 0.8476 | -21.71 | 217 | 117 | 270 | 153 | |
| Polynomial Degree 2 | 0.8641 | -21.74 | 219 | 117 | 270 | 153 | |
| Power | 0.6148 | -19.78 | 222 | 118 | 271 | 154 | |
| Linear | 0.7377 | -21.43 | 179 | 115 | 235 | 150 | |

^a Selected model in bold.

Plots of the Polynomial 2-degree model with BMRs of one SD and 10% RD are shown in Figure 1-81 and Figure 1-82, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-83 (BMD and BMDL shown are for BMR of 10% RD; the rest is applicable to both BMRs).




Figure 1-81. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 1SD (Constant Variance)




Figure 1-82. Plot of Response by Dose with Fitted Curve for the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 10%RD (Constant Variance)

| Model Results | | | | | | | | |
|-------------------------|--------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 219.1614151 | | | | | | | |
| BMDL | 117.214416 | | | | | | | |
| BMDU | 380.6486941 | | | | | | | |
| AIC | -21.74414133 | | | | | | | |
| Test 4 P-value | 0.8640863 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 1.496455497 | 4.99E-02 | 1.39871499 | 1.594196 | | | | |
| beta | 0.000323664 | 5.94E-04 | -0.0008398 | 0.00148709 | | | | |
| beta2 | Bounded | NA | NA | NA | | | | |
| alpha | 0.012545321 | 5.25E-05 | 0.01244245 | 0.01264819 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 5 | 1.496455497 | 1.5 | 1.5 | 0.1120059 | 0.09 | 0.09 | 0.070761898 |
| 200 | 5 | 1.59539172 | 1.58 | 1.58 | 0.1120059 | 0.19 | 0.19 | -0.307277867 |
| 300 | 5 | 1.67051238 | 1.69 | 1.69 | 0.1120059 | 0.09 | 0.09 | 0.389047757 |
| 400 | 3 | 1.762734739 | 1.75 | 1.75 | 0.1120059 | 0.08 | 0.08 | -0.196929058 |

| Likelihoods of Interest | | | |
|-------------------------|-----------------|-----------------|------------|
| Model | Log Likelihood* | # of Parameters | AIC |
| A1 | 14.0181533 | 5 | -18.036307 |
| A2 | 16.70117875 | 8 | -17.402358 |
| A3 | 14.0181533 | 5 | -18.036307 |
| fitted | 13.87207067 | 3 | -21.744141 |
| R | 9.252043047 | 2 | -14.504086 |

| Tests of Interest | | | |
|-------------------|--------------------------|---------|------------|
| Test | -2*Log(Likelihood Ratio) | Test df | p-value |
| 1 | 14.89827141 | 6 | 0.02106286 |
| 2 | 5.366050911 | 3 | 0.14687341 |
| 3 | 5.366050911 | 3 | 0.14687341 |
| 4 | 0.292165262 | 2 | 0.8640863 |

Figure 1-83. Details Regarding the Selected Model (Polynomial 2-Degree) for Increased Relative Kidney Weight in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study

1.1.2.1.3.2 Blood Urea Nitrogen in Male B6C3F1 Mice – Single Oral Gavage

BUN levels appeared to show a dose-related increasing trend (not statistically significant) in male mice exposed to 1,2-dichloroethane following a single oral gavage exposure in an acute toxicity study by [Storer et al. \(1984\)](#). The doses and response data used for the modeling are presented in Table 1-65. Continuous models were fit to the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* ([U.S. EPA, 2012](#)). A BMR of 10% RD was also selected.

Table 1-65. Blood Urea Nitrogen in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Oral Exposure Study

| Dose (mg/kg) | Number of Animals | Mean (mg urea nitrogen/100 mL serum) | SD (mg urea nitrogen/100 mL serum) |
|--------------|-------------------|--------------------------------------|------------------------------------|
| 0 | 5 | 13.4 | 2.3 |
| 200 | 5 | 16.6 | 15.1 |
| 300 | 5 | 18.5 | 10.3 |
| 400 | 3 | 26.2 | 21.8 |

The BMD modeling results for increased BUN levels in male mice are summarized in Table 1-66. The constant variance model did not provide adequate fit to the variance data, but the nonconstant variance model did. With the nonconstant variance model applied and using a BMR of one SD, only the Power model provided adequate fit to the means (test 4 p-value > 0.1) and a viable result; this model was selected for the one SD BMR. When a BMR of 10% RD was applied, all models, except for the Exponential 3 model, provided an adequate fit to the means (test 4 p-value > 0.1); however, the BMDL computation failed for the Hill model and BMDs and BMDLs for the Exponential 5 model were 10 times lower than the lowest non-zero dose and, therefore, not considered viable. The BMDLs for the

remaining fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected for the 10% RD BMR (Linear).

Table 1-66. Summary of BMD Modeling Results for Increased Blood Urea Nitrogen in Male Mice Following a Single Oral Exposure to 1,2-Dichloroethane (Nonconstant Variance)^a

| Model | Goodness of Fit | | BMD 1SD (mg/kg) | BMDL 1SD (mg/kg) | BMD 10%RD (mg/kg) | BMDL 10%RD (mg/kg) | Basis for Model Selection |
|------------------------|-------------------|----------------|-----------------------|------------------------|-------------------------|--------------------------|---|
| | Test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.0642 | 138.0 | 156 | 70.9 | 68.0 | 37.7 | Using a BMR of one SD, only the Power model provided adequate fit to the means (test 4 p-value > 0.1) and a viable result; this model was selected for the one SD BMR. With a BMR of 10% RD, all models, except for the Exponential 3 model, provided an adequate fit to the means (test 4 p-value > 0.1); however, the Hill and Exponential 5 models were not viable. The BMDLs for the remaining viable models differed by < 3-fold; therefore, the model with the lowest AIC was selected for the 10% RD BMR (Linear). |
| Exponential 5 | 0.3474 | 134.6 | 35.4 | 0.814 | 19.1 | 0.712 | |
| Hill | 0.1560 | 136.5 | 24.7 | 0 | 12.7 | 0 | |
| Polynomial Degree 3 | 0.1170 | 136.824 | - | - | 54.3 | 25.8 | |
| Polynomial Degree 2 | 0.1168 | 136.827 | - | - | 53.9 | 25.8 | |
| Power | 0.1072 | 136.999 | 106 | 42.8 | 46.6 | 25.3 | |
| Linear | 0.1171 | 136.822 | - | - | 54.9 | 25.8 | |

^a Selected model in bold.

A plot of the Power model with a BMR of one SD is shown in Figure 1-84 and a plot of the Linear model with a BMR of 10% RD is shown in Figure 1-85. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood for the Power model with BMR of one SD are shown in Figure 1-86 and additional modeling details for the Linear model with BMR of 10% RD are shown in Figure 1-87.




Figure 1-84. Plot of Response by Dose with Fitted Curve for the Selected Model (Power) for Increased Blood Urea Nitrogen in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 1SD (Nonconstant Variance)




Figure 1-85. Plot of Response by Dose with Fitted Curve for the Selected Model (Linear) for Increased Blood Urea Nitrogen in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study and BMR of 10%RD (Nonconstant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 106.3910977 | | | | | | | |
| BMDL | 42.83193912 | | | | | | | |
| BMDU | 108.5933934 | | | | | | | |
| AIC | 136.9992953 | | | | | | | |
| Test 4 P-value | 0.107170237 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 5 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 12.93552585 | 1.295836951 | 10.3957321 | 15.4753196 | | | | |
| v | 0.02778794 | 1.24E-02 | 0.00350199 | 0.05207389 | | | | |
| n | Bounded | NA | NA | NA | | | | |
| rho | 6.354251024 | 2.97E+00 | 0.54176928 | 12.1667328 | | | | |
| alpha | 7.533E-07 | 4.75E-12 | 7.5329E-07 | 7.5331E-07 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 5 | 12.93552585 | 13.4 | 13.4 | 2.95638946 | 2.3 | 2.3 | 0.351305461 |
| 200 | 5 | 18.49311389 | 16.6 | 16.6 | 9.2030968 | 15.1 | 15.1 | -0.459968144 |
| 300 | 5 | 21.27190791 | 18.5 | 18.5 | 14.3579284 | 10.3 | 10.3 | -0.431690029 |
| 400 | 3 | 24.05070193 | 26.2 | 26.2 | 21.2080278 | 21.8 | 21.8 | 0.175532279 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -69.2249334 | 5 | 148.449867 | | | | | |
| A2 | -61.90358802 | 8 | 139.807176 | | | | | |
| A3 | -62.26631094 | 6 | 136.532622 | | | | | |
| fitted | -64.49964765 | 4 | 136.999295 | | | | | |
| R | -70.38699059 | 2 | 144.773981 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 16.96680515 | 6 | 0.00940602 | | | | | |
| 2 | 14.64269077 | 3 | 0.00214891 | | | | | |
| 3 | 0.725445834 | 2 | 0.6957792 | | | | | |
| 4 | 4.466673423 | 2 | 0.10717024 | | | | | |

Figure 1-86. Details Regarding the Selected Model (Power) for Increased BUN Levels in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study (BMR of 1SD)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 54.86685634 | | | | | | | |
| BMDL | 25.80392856 | | | | | | | |
| BMDU | 112.6650676 | | | | | | | |
| AIC | 136.8221271 | | | | | | | |
| Test 4 P-value | 0.117097004 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 13.17889964 | 1.067007319 | 11.0876037 | 15.2701956 | | | | |
| beta | 0.024019782 | 8.14E-03 | 0.0080685 | 0.03997107 | | | | |
| rho | 7.756854984 | 3.18E-01 | 7.1330916 | 8.38061837 | | | | |
| alpha | 1.523E-08 | 1.81E-20 | 1.523E-08 | 1.523E-08 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 5 | 13.17889964 | 13.4 | 13.4 | 2.72094491 | 2.3 | 2.3 | 0.181699903 |
| 200 | 5 | 17.98285606 | 16.6 | 16.6 | 9.08297293 | 15.1 | 15.1 | -0.340434809 |
| 300 | 5 | 20.38483427 | 18.5 | 18.5 | 14.7707533 | 10.3 | 10.3 | -0.285335316 |
| 400 | 3 | 22.78681248 | 26.2 | 26.2 | 22.7523877 | 21.8 | 21.8 | 0.259832695 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -69.2249334 | 5 | 148.449867 | | | | | |
| A2 | -61.90358802 | 8 | 139.807176 | | | | | |
| A3 | -62.26631094 | 6 | 136.532622 | | | | | |
| fitted | -64.41106353 | 4 | 136.822127 | | | | | |
| R | -70.38699059 | 2 | 144.773981 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 16.96680515 | 6 | 0.00940602 | | | | | |
| 2 | 14.64269077 | 3 | 0.00214891 | | | | | |
| 3 | 0.725445834 | 2 | 0.6957792 | | | | | |
| 4 | 4.289505196 | 2 | 0.117097 | | | | | |

Figure 1-87. Details Regarding the Selected Model (Linear) for Increased BUN Levels in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage in an Acute Toxicity Study (BMR of 10%RD)

1.1.2.2 Short-term/Intermediate

1.1.2.2.1 Mortality

[NTP \(1991\)](#) provided data showing increased mortality in rats following intermediate oral exposure to 1,2-dichloroethane.

1.1.2.2.1.1 Mortality in F344 Male Rats – 13-Week Gavage

There was an increased incidence of mortality in male rats exposed to 1,2-dichloroethane by gavage for 13 weeks (5 days per week) ([NTP, 1991](#)). The administered doses were duration adjusted to estimate an equivalent oral dose for animals exposed for 7 days per week. The dose and response data used for the modeling are presented in Table 1-67. Dichotomous models were used to fit dose-response data.

A BMR of 10% ER was chosen according to *BMD Technical Guidance* ([U.S. EPA, 2012](#)). BMRs of five and one percent ER were also selected due to severity of the endpoint.

Table 1-67. Incidence of Mortality in Male Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study

| Adjusted Dose (mg/kg-day) | Number of Animals | Incidence |
|---------------------------|-------------------|-----------|
| 0 | 10 | 0 |
| 21 | 10 | 0 |
| 43 | 10 | 0 |
| 86 | 10 | 0 |
| 171 | 10 | 10 |
| 343 | 10 | 10 |

The BMD modeling results for mortality in male rats are summarized in Table 1-68. All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Multistage 1-degree/Quantal Linear model. The BMD computation failed for the Weibull model because the lower limit included zero; therefore, this model was unusable. With a BMR of 10% ER applied, the BMDLs for the remaining models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Logistic) was selected. With BMRs of five and one percent ER applied (below the observable range in the study), model uncertainty increased and the BMDLs of the fit models were not sufficiently close (differed by > 3-fold); therefore, the BMDS recommended the model with the lowest BMDL (Multistage 2-degree).

This data set is not well suited for BMD modeling; there are no data points with incidence between 0 and 100 percent, and no data to inform the shape of the curve at the region of interest, as defined by the BMR. As a result, the different models provide a broad range of BMD and BMDL estimates. For the five and one percent ER BMRs, selection of the low end of the range to represent the BMDL is not among the more realistic possibilities and involves extrapolation below the observable range in the study (1/10 = 10%) to generate BMD estimates well below the lowest tested dose where any mortality was observed.

Table 1-68. Summary of BMD Modeling Results for Increased Incidence of Mortality in Male Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks^a

| Model | Goodness of Fit (Means) | | BMD 1%ER (mg/kg-day) | BMDL 1%ER (mg/kg-day) | BMD 5%ER (mg/kg-day) | BMDL 5%ER (mg/kg-day) | BMD 10%ER (mg/kg-day) | BMDL 10%ER (mg/kg-day) | Basis for Model Selection |
|---------------------|-------------------------|--------------|----------------------|-----------------------|----------------------|-----------------------|-----------------------|------------------------|--|
| | p-value | AIC | | | | | | | |
| Dichotomous Hill | 0.9997 | 4.099 | 95 | 64 | 100 | 78 | 110 | 84 | <p>All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Multistage 1-degree/Quantal Linear model. The BMD computation failed for the Weibull model. With a BMR of 10% ER applied, BMDLs for the remaining models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC (Logistic). With BMRs of five and one percent ER applied, BMDLs of the fit models differed by > 3-fold; therefore, EPA chose the model with the lowest BMDL (Multistage 2-degree) for this BMR.</p> <p>NOTE: This data set is not well suited for BMD modeling and the results for BMR = 5% ER and one percent ER are highly unrealistic.</p> |
| Gamma | 0.9010 | 5.080 | 66 | 51 | 80 | 65 | 88 | 73 | |
| Log-Logistic | 0.9997 | 4.099 | 95 | 64 | 100 | 78 | 110 | 84 | |
| Multistage 3 | 0.5659 | 8.699 | 29 | 9.0 | 50 | 34 | 63 | 50 | |
| Multistage 2 | 0.1217 | 17.01 | 14 | 6.0 | 32 | 22 | 47 | 35 | |
| Multistage 1 | 0.0028 | 31.61 | 1.7 | 1.2 | 8.7 | 5.9 | 18 | 12 | |
| Weibull | 0.9101 | 4.941 | 57 | 0 | 78 | 0 | 90 | 0 | |
| Logistic | 1.000 | 2.098 | 94 | 51 | 100 | 74 | 110 | 83 | |
| Log-Probit | 1.000 | 4.000 | 110 | 68 | 110 | 79 | 120 | 84 | |
| Probit | 0.9922 | 4.518 | 84 | 54 | 97 | 72 | 100 | 81 | |
| Quantal Linear | 0.0028 | 31.61 | 1.7 | 1.2 | 8.7 | 5.9 | 18 | 12 | |

^a Selected model in bold.

A plot of the Logistic model with a BMR of 10% ER is shown in Figure 1-88 and plots of the Multistage (2-degree) model for BMRs of 5% ER and 5% ER are shown in Figure 1-89 and Figure 1-90, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood for the Logistic model with BMR of 10% ER are shown in Figure 1-91; additional modeling details for the Multistage 2-degree model are shown in Figure 1-92 (BMD and BMDL shown are for BMR of 5% ER; the rest is applicable to both BMRs).




Figure 1-88. Plot of Response by Dose with Fitted Curve for the Selected Model (Logistic Model) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%ER




Figure 1-89. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 5%ER




Figure 1-90. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 1%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 110.6991871 | | | | |
| BMDL | 83.20918801 | | | | |
| BMDU | 135.7943266 | | | | |
| AIC | 2.098271204 | | | | |
| P-value | 0.999971933 | | | | |
| D.O.F. | 5 | | | | |
| Chi ² | 0.049202422 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| a | Bounded | NA | NA | NA | |
| b | 0.142754215 | 3.73E-02 | 0.06967734 | 0.21583109 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 21 | 3.05242E-07 | 3.05242E-06 | 0 | 10 | -0.001747 |
| 43 | 7.0564E-06 | 7.0564E-05 | 0 | 10 | -0.0084 |
| 86 | 0.003258783 | 0.032587826 | 0 | 10 | -0.180816 |
| 171 | 0.998359268 | 9.983592681 | 10 | 10 | 0.1281963 |
| 343 | 1 | 10 | 10 | 10 | 5.972E-07 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | 4.44089E-16 | 6 | - | - | NA |
| Fitted Model | -0.049135602 | 1 | 0.0982712 | 5 | 0.9998445 |
| Reduced Model | -38.1908501 | 1 | 76.283429 | 5 | <0.0001 |

Figure 1-91. Details Regarding the Selected Model (Logistic) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 32.47950141 | | | | |
| BMDL | 21.58183101 | | | | |
| BMDU | 42.58707132 | | | | |
| AIC | 17.00744744 | | | | |
| P-value | 0.121722096 | | | | |
| D.O.F. | 5 | | | | |
| Chi ² | 8.698255892 | | | | |
| Slope Factor | 0.002316764 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 3 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | Bounded | NA | NA | NA | |
| b2 | 4.8623E-05 | 1.850501141 | -3.626867 | 3.62696424 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 21 | 0.021214505 | 0.212145047 | 0 | 10 | -0.465557 |
| 43 | 0.085981052 | 0.859810517 | 0 | 10 | -0.969893 |
| 86 | 0.302055617 | 3.020556174 | 0 | 10 | -2.080334 |
| 171 | 0.758717227 | 7.587172269 | 10 | 10 | 1.783295 |
| 343 | 0.996721763 | 9.967217631 | 10 | 10 | 0.1813565 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | 4.44089E-16 | 6 | - | - | NA |
| Fitted Model | -7.503723719 | 1 | 15.0074474 | 5 | 0.0103306 |
| Reduced Model | -38.1908501 | 1 | 61.3742528 | 5 | <0.0001 |

Figure 1-92. Details Regarding the Selected Model (Multistage 2-Degree) for Mortality in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks

1.1.2.2.1.2 Mortality in F344 Female Rats – 13-Week Gavage

There was an increased incidence of mortality in female rats exposed to 1,2-dichloroethane by gavage for 13 weeks (5 days per week) (NTP, 1991). The administered doses were duration adjusted to estimate an equivalent oral dose for animals exposed for 7 days per week. The dose and response data used for the modeling are presented in Table 1-69. Dichotomous models were used to fit dose-response data.

A BMR of 10% ER was chosen according to *BMD Technical Guidance* (U.S. EPA, 2012). BMRs of five and one percent ER were also selected due to severity of the endpoint.

Table 1-69. Incidence of Mortality in Female Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study

| Adjusted Dose (mg/kg-day) | Number of Animals | Incidence |
|---------------------------|-------------------|-----------|
| 0 | 10 | 0 |
| 13 | 10 | 0 |
| 26 | 10 | 0 |
| 54 | 10 | 0 |
| 107 | 10 | 0 |
| 214 | 10 | 9 |

The BMD modeling results for increased mortality in female rats are summarized in Table 1-70. All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Multistage 1-degree/Quantal Linear model. With a BMR of 10% ER applied, the BMDLs were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Weibull) was selected. With BMRs of five and one percent ER applied (below the observable range in the study), model uncertainty increased and the BMDLs of the fit models were not sufficiently close (differed by > 3-fold); therefore, the BMDS recommended the model with the lowest BMDL (Multistage 2-degree).

This data set is not well suited for BMD modeling; there are no data points with incidence between 0 and 90 percent, and there are no data to inform the shape of the curve at the region of interest, as defined by the BMR. As a result, the different models provide a broad range of BMD and BMDL estimates. For the five and one percent ER BMRs, selection of the low end of the range to represent the BMDL is not among the more realistic possibilities and involves extrapolation below the range of observation in the study (1/10 = 10%) to generate a BMD estimate well below the lowest tested dose where any mortality was observed.

Table 1-70. Summary of BMD Modeling Results for Increased Incidence of Mortality in Female Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks^a

| Model | Goodness of Fit (Means) | | BMD 1%RD (mg/kg-day) | BMDL 1%RD (mg/kg-day) | BMD 5%RD (mg/kg-day) | BMDL 5%RD (mg/kg-day) | BMD 10%RD (mg/kg-day) | BMDL 10%RD (mg/kg-day) | Basis for Model Selection |
|---------------------|-------------------------|--------------|----------------------|-----------------------|----------------------|-----------------------|-----------------------|------------------------|--|
| | p-value | AIC | | | | | | | |
| Dichotomous Hill | 1.000 | 8.502 | 150 | 67 | 160 | 90 | 168 | 103 | <p>All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Multistage 1-degree/Quantal Linear model. With a BMR of 10% ER applied, BMDLs differed by < 3-fold; therefore, EPA chose the model with the lowest AIC (Weibull). With BMRs of five and one percent ER applied, BMDLs of the fit models differed by > 3-fold; therefore, the BMDS recommended the model with the lowest BMDL (Multistage 2-degree).</p> <p>NOTE: This data set is not well suited for BMD modeling and the results for BMR = 5% ER and one percent ER are highly unrealistic.</p> |
| Gamma | 0.9903 | 9.462 | 91 | 62 | 110 | 83 | 121 | 95.9 | |
| Log-Logistic | 1.000 | 8.502 | 150 | 67 | 160 | 90 | 168 | 103 | |
| Multistage 3 | 0.7864 | 11.83 | 40 | 12 | 69 | 46 | 87.4 | 68.3 | |
| Multistage 2 | 0.2605 | 19.79 | 20 | 8.3 | 46 | 30 | 65.5 | 48.3 | |
| Multistage 1 | 0.0146 | 28.73 | 3.4 | 2.1 | 18 | 10 | 36.0 | 21.6 | |
| Weibull | 1.000 | 8.502 | 160 | 58 | 170 | 86 | 180 | 102 | |
| Logistic | 1.000 | 8.509 | 140 | 48 | 160 | 87 | 167 | 105 | |
| Log-Probit | 1.000 | 10.50 | 170 | 75 | 180 | 92 | 185 | 103 | |
| Probit | 1.000 | 10.50 | 160 | 56 | 170 | 87 | 174 | 103 | |
| Quantal Linear | 0.0146 | 28.73 | 3.4 | 2.1 | 18 | 10 | 36.0 | 21.6 | |

^a Selected model in bold.

A plot of the Weibull model with a BMR of 10% ER is shown in Figure 1-93 and plots of the Multistage (2-degree) model with a BMR of 5% ER and one percent ER are shown in Figure 1-94 and Figure 1-95, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood for the Weibull model with BMR of 10% ER are shown in Figure 1-96; additional modeling details for the Multistage 2-degree model are shown in Figure 1-97 (BMD and BMDL shown are for BMR of 5% ER; the rest is applicable to both BMRs).




Figure 1-93. Plot of Response by Dose with Fitted Curve for the Selected Model (Weibull) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%ER




Figure 1-94. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 5%ER




Figure 1-95. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 2-Degree) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 1%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 180.2998504 | | | | |
| BMDL | 102.2067187 | | | | |
| BMDU | 187.4610082 | | | | |
| AIC | 8.501836664 | | | | |
| P-value | 1 | | | | |
| D.O.F. | 5 | | | | |
| Chi ² | 8.8597E-05 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 3 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| a | Bounded | NA | NA | NA | |
| b | 2.59873E-42 | 0.948650386 | -1.8593206 | 1.8593206 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 13 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 26 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 54 | 1.52696E-08 | 1.52696E-07 | 0 | 10 | -0.000391 |
| 107 | 8.79873E-06 | 8.79873E-05 | 0 | 10 | -0.00938 |
| 214 | 0.899996567 | 8.999965671 | 9 | 10 | 3.618E-05 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -3.250829734 | 6 | - | - | NA |
| Fitted Model | -3.250918332 | 1 | 0.0001772 | 5 | 1 |
| Reduced Model | -25.36254527 | 1 | 44.2232539 | 5 | <0.0001 |

Figure 1-96. Details Regarding the Selected Model (Weibull) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 45.67835414 | | | | |
| BMDL | 30.17422595 | | | | |
| BMDU | 62.50104948 | | | | |
| AIC | 18.78831282 | | | | |
| P-value | 0.260531164 | | | | |
| D.O.F. | 5 | | | | |
| Chi ² | 6.500319411 | | | | |
| Slope Factor | 0.001657043 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 3 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | Bounded | NA | NA | NA | |
| b2 | 2.45833E-05 | 0.395402699 | -0.7749505 | 0.77499964 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 13 | 0.00414597 | 0.041459704 | 0 | 10 | -0.20404 |
| 26 | 0.016480987 | 0.164809869 | 0 | 10 | -0.409355 |
| 54 | 0.069175797 | 0.691757972 | 0 | 10 | -0.862071 |
| 107 | 0.245314319 | 2.453143191 | 0 | 10 | -1.802928 |
| 214 | 0.675612239 | 6.756122394 | 9 | 10 | 1.5157172 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -3.250829734 | 6 | - | - | NA |
| Fitted Model | -8.394156409 | 1 | 10.2866533 | 5 | 0.0675087 |
| Reduced Model | -25.36254527 | 1 | 44.2234311 | 5 | <0.0001 |

Figure 1-97. Details Regarding the Selected Model (Multistage 2-Degree) for Mortality in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks

1.1.2.2.2 Body Weight Effects

One short-term/intermediate-duration inhalation study was identified for BMD modeling that showed significant changes in maternal absolute body weight gain in mice ([Payan et al., 1995](#)).

1.1.2.2.2.1 Maternal Absolute Weight Gain in Female Rats – Oral Gavage, GD 6 to 20

Absolute body weight gain was significantly increased in maternal rats exposed to 1,2-dichloroethane by gavage on GD 6 to 20 ([Payan et al., 1995](#)). Doses were reported in mmol/kg/day and converted to units of mg/kg-day. The dose and response data used for the modeling are presented in Table 1-71. Continuous models were used to fit dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected.

Table 1-71. Maternal Absolute Weight Gain in Female Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from an Oral Gestational Exposure Study

| Dose (mg/kg-day) | Number of Animals | Mean (g) | SD (g) |
|------------------|-------------------|----------|--------|
| 0 | 26 | 43 | 10 |
| 119 | 26 | 39 | 15 |
| 158 | 25 | 33 | 20 |
| 198 | 26 | 30 | 20 |
| 238 | 26 | 22 | 25 |

The BMD modeling results for decreased maternal absolute weight gain in female rats are summarized in Table 1-72. The constant variance model did not provide adequate fit to the variance data, but the nonconstant variance model did. With the nonconstant variance model applied, only the Hill model provided an adequate fit to the means (test 4 p-value > 0.1); therefore, this model was selected.

Table 1-72. Summary of BMD Modeling Results for Decreased Maternal Absolute Weight Gain in Female Rats Following Oral Exposure to 1,2-Dichloroethane on GD 6 to 20 (Nonconstant Variance)^a

| Model | Goodness of Fit | | BMD 1SD (mg/kg-day) | BMDL 1SD (mg/kg-day) | BMD 10%RD (mg/kg-day) | BMDL 10%RD (mg/kg-day) | Basis for Model Selection |
|---------------------|-----------------|-------------|---------------------|----------------------|-----------------------|------------------------|---|
| | test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.0002 | 1125 | 224 | 176 | 118 | 52.7 | Only the Hill model provided an adequate fit to the means (test 4 p-value > 0.1); therefore, this model was selected. |
| Exponential 5 | < 0.0001 | 1127 | 224 | 176 | 118 | 52.7 | |
| Hill | 0.3438 | 1109 | 154 | 111 | 99.1 | 41.8 | |
| Polynomial Degree 3 | 0.0005 | 1123 | 224 | 178 | 113 | 48.5 | |
| Polynomial Degree 2 | 0.0005 | 1123 | 223 | 178 | 108 | 48.4 | |
| Power | 0.0002 | 1125 | 224 | 177 | 114 | 48.5 | |
| Linear | 0.0002 | 1125 | 221 | 158 | 54.2 | 42.6 | |

^a Selected model in bold.

Plots of the Hill model with BMRs of one SD and 10% RD are shown in Figure 1-98 and Figure 1-99, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-100 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-98. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Decreased Maternal Absolute Weight Gain in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage GD 6 to 20 and BMR of 1SD (Nonconstant Variance)




Figure 1-99. Plot of Response by Dose with Fitted Curve for the Selected Model (Hill) for Decreased Maternal Absolute Weight Gain in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage GD 6 to 20 and BMR of 10%RD (Nonconstant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 154.0869579 | | | | | | | |
| BMDL | 110.7483035 | | | | | | | |
| BMDU | 189.773709 | | | | | | | |
| AIC | 1109.497668 | | | | | | | |
| Test 4 P-value | 0.343844647 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 6 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 43.68213907 | 1.959630152 | 39.8413345 | 47.5229436 | | | | |
| v | -22.53459618 | 19.65018267 | -61.048247 | 15.9790544 | | | | |
| k | 160.4215626 | 105.8714667 | -47.082701 | 367.925826 | | | | |
| n | 2.957753882 | 3.483418283 | -3.8696205 | 9.78512831 | | | | |
| rho | -3.482162919 | 1.383320231 | -6.1934208 | -0.7709051 | | | | |
| alpha | 57829944.68 | 1.66E+16 | -3.252E+16 | 3.2521E+16 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 26 | 43.68213907 | 43 | 43 | 10.5967743 | 10 | 10 | -0.328235774 |
| 119 | 26 | 37.09148674 | 39 | 39 | 14.0877651 | 15 | 15 | 0.690779997 |
| 158 | 25 | 32.66824306 | 33 | 33 | 17.5735729 | 20 | 20 | 0.094390862 |
| 198 | 26 | 29.01694655 | 30 | 30 | 21.6013361 | 20 | 20 | 0.232050864 |
| 238 | 26 | 26.49829695 | 22 | 22 | 25.3010171 | 25 | 25 | -0.906560548 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -558.2556686 | 6 | 1128.51134 | | | | | |
| A2 | -547.2429538 | 10 | 1114.48591 | | | | | |
| A3 | -548.3008121 | 7 | 1110.60162 | | | | | |
| fitted | -548.7488342 | 6 | 1109.49767 | | | | | |
| R | -567.776198 | 2 | 1139.5524 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 41.06648838 | 8 | <0.0001 | | | | | |
| 2 | 22.02542948 | 4 | 0.0001981 | | | | | |
| 3 | 2.11571657 | 3 | 0.54873974 | | | | | |
| 4 | 0.896044122 | 1 | 0.34384465 | | | | | |

Figure 1-100. Details Regarding the Selected Model (Hill) for Decreased Maternal Absolute Weight Gain in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage GD 6 to 20

1.1.2.2.3 Renal Effects

Data sets for changes in absolute and relative kidney weight in male and female rats orally exposed for 13 weeks were identified for BMD modeling ([NTP, 1991](#)).

Modeled results were not presented for the absolute or relative kidney weight data in female rats because neither the constant nor nonconstant variance models provided adequate fit to the variance data.

1.1.2.2.3.1 Absolute Kidney Weight in F344 Male Rats – 13-Week Gavage

Absolute kidney weight was significantly increased in male rats exposed to 1,2-dichloroethane by gavage for 13 weeks (5 days per week) ([NTP, 1991](#)). The administered doses were duration adjusted to estimate an equivalent oral dose for animals exposed for 7 days per week. The dose and response data used for the modeling are presented in Table 1-73. Continuous models were used to fit dose-response data.

A BMR of one SD was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). A BMR of 10% RD was also selected because EPA considers a 10% change in kidney weight to be biologically significant.

Table 1-73. Absolute Kidney Weight in Male Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study

| Adjusted Dose (mg/kg-day) | Number of Animals | Mean (mg) | SD (mg) |
|---------------------------|-------------------|-----------|---------|
| 0 | 10 | 1324 | 92 |
| 21 | 10 | 1441 | 82 |
| 43 | 10 | 1600 | 171 |
| 86 | 10 | 1653 | 149 |

The BMD modeling results for increased absolute kidney weight in male rats are summarized in Table 1-74. Both the constant and nonconstant variance models provide adequate fit to the variance data. With the constant variance model applied, none of the models provided adequate fit to the means (test 4 p-value < 0.1). With the nonconstant variance model applied, only the Exponential 5 model provided adequate fit to the means (test 4 p-value > 0.1); therefore, this model was selected.

Table 1-74. Summary of BMD Modeling Results for Increased Absolute Kidney Weight in Male Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks (Nonconstant Variance)^a

| Model | Goodness of Fit | | BMD 1SD (mg/kg-day) | BMDL 1SD (mg/kg-day) | BMD 10%RD (mg/kg-day) | BMDL 10%RD (mg/kg-day) | Basis for Model Selection |
|----------------------|-----------------|--------------|---------------------|----------------------|-----------------------|------------------------|---|
| | test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.0388 | 507.1 | 27 | 17 | 35 | 27 | Only the Exponential 5 model provided adequate fit to the means (test 4 p-value > 0.1); therefore, this model was selected. |
| Exponential 5 | 0.3660 | 503.5 | 19 | 12 | 23 | 14 | |
| Hill | NA | 505.5 | 19 | 13 | 23 | 19 | |
| Polynomial Degree 3 | 0.0268 | 507.9 | - | - | 31 | 0 | |

| Model | Goodness of Fit | | BMD 1SD (mg/kg- day) | BMDL 1SD (mg/kg- day) | BMD 10%RD (mg/kg- day) | BMDL 10%RD (mg/kg- day) | Basis for Model Selection |
|------------------------|--------------------|-------|-------------------------------|--------------------------------|---------------------------------|----------------------------------|---------------------------|
| | test 4 p- value | AIC | | | | | |
| Polynomial Degree 2 | 0.0268 | 507.9 | - | - | 31 | 0 | |
| Power | 0.0268 | 507.9 | 23 | 14 | 31 | 24 | |
| Linear | 0.0268 | 507.9 | - | - | 31 | 24 | |

^a Selected model in bold.

Plots of the Exponential 5 model with BMRs of one SD and 10% RD are shown in Figure 1-101 and Figure 1-102, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-103 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-101. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 5) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 1SD (Nonconstant Variance)

Frequentist Exponential Degree 5 Model with BMR of 0.1 Relative Deviation for the BMD and 0.95 Lower Confidence Limit for the BMDL




Figure 1-102. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 5) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%RD (Nonconstant Variance)

| Model Results | | | | | | | | |
|-------------------------|-------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 18.89503404 | | | | | | | |
| BMDL | 11.64401146 | | | | | | | |
| BMDU | 25.54732985 | | | | | | | |
| AIC | 503.4998688 | | | | | | | |
| Test 4 P-value | 0.366027487 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 6 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| a | 1322.943197 | 28.52751242 | 1267.0303 | 1378.85609 | | | | |
| b | 0.031943071 | 6.44E-03 | 0.01931208 | 0.04457407 | | | | |
| c | 1.250041712 | 3.38E-02 | 1.18379403 | 1.31628939 | | | | |
| d | 2.234728895 | 9.38E-01 | 0.39703395 | 4.07242384 | | | | |
| rho | 3.760841847 | 3.12E-02 | 3.69972526 | 3.82195844 | | | | |
| log-alpha | Bounded | NA | NA | NA | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 10 | 1322.943197 | 1324 | 1324 | 91.4452272 | 92 | 92 | 0.036545426 |
| 21 | 10 | 1434.143559 | 1441 | 1441 | 106.432064 | 82 | 82 | 0.203716537 |
| 43 | 10 | 1610.401275 | 1600 | 1600 | 132.353569 | 171 | 171 | -0.248514042 |
| 86 | 10 | 1653.711018 | 1653 | 1653 | 139.126056 | 149 | 149 | -0.016161145 |

| Likelihoods of Interest | | | |
|-------------------------|-----------------|-----------------|------------|
| Model | Log Likelihood* | # of Parameters | AIC |
| A1 | -249.0626479 | 5 | 508.125296 |
| A2 | -245.3915079 | 8 | 506.783016 |
| A3 | -246.3413811 | 6 | 504.682762 |
| fitted | -246.7499344 | 5 | 503.499869 |
| R | -264.1885774 | 2 | 532.377155 |

| Tests of Interest | | | |
|-------------------|--------------------------|---------|------------|
| Test | -2*Log(Likelihood Ratio) | Test df | p-value |
| 1 | 37.59413899 | 6 | <0.0001 |
| 2 | 7.342280042 | 3 | 0.06175249 |
| 3 | 1.899746393 | 2 | 0.38679007 |
| 4 | 0.817106605 | 1 | 0.36602749 |

Figure 1-103. Details Regarding the Selected Model (Exponential 5) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks

1.1.2.2.3.2 Relative Kidney Weight in F344 Male Rats – 13-Week Gavage

Relative kidney weight was significantly increased in male rats exposed to 1,2-dichloroethane by gavage for 13 weeks (5 days per week) (NTP, 1991). The administered doses were duration adjusted to estimate an equivalent oral dose for animals exposed for 7 days per week. The dose and response data used for the modeling are presented in Table 1-75. Continuous models were used to fit dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* (U.S. EPA, 2012). A BMR of 10% RD was also selected because EPA considers a 10% change in relative kidney weight to be biologically significant.

Table 1-75. Relative Kidney Weight in Male Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study

| Adjusted Dose (mg/kg-day) | Number of Animals | Mean (Organ weight to body weight) | SD (Organ weight to body weight) |
|---------------------------|-------------------|------------------------------------|----------------------------------|
| 0 | 10 | 3.9 | 0.19 |
| 21 | 10 | 4.1 | 0.32 |
| 43 | 10 | 4.5 | 0.25 |
| 86 | 10 | 4.9 | 0.22 |

The BMD modeling results for increased relative kidney weight in male rats are summarized in Table 1-76. The constant variance model provided an adequate fit to the variance data. With the constant variance model applied, the goodness-of-fit p-values for the means (test 4) could not be derived for the Exponential 5 and Hill models because the models were saturated (degrees of freedom = 0). The remaining models provided adequate fit to the means (test 4 p-value > 0.1). The BMDLs for the fit

models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Power) was selected.

Table 1-76. Summary of BMD Modeling Results for Increased Relative Kidney Weight in Male Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks (Constant Variance)^a

| Model | Goodness of Fit | | BMD 1SD (mg/kg- day) | BMDL 1SD (mg/kg- day) | BMD 10%R D (mg/kg- day) | BMDL 10%R D (mg/kg- day) | Basis for Model Selection |
|------------------------|-----------------------|---------------|-------------------------------|--------------------------------|-------------------------------------|--------------------------------------|--|
| | test 4 p- value | AIC | | | | | |
| Exponential 3 | 0.2945 | 6.746 | 23 | 18 | 36 | 30 | Goodness-of-fit p-values could not be derived for the Exponential 5 and Hill models because the models were saturated (degrees of freedom = 0). The remaining models provided adequate fit to the means (test 4 p-value > 0.1). BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | 8.301 | 23 | 13 | 32 | 21 | |
| Hill | NA | 8.301 | 23 | 13 | 31 | 21 | |
| Polynomial Degree 3 | 0.3915 | 6.1763 | 20 | 16 | 33 | 27 | |
| Polynomial Degree 2 | 0.3908 | 6.180 | 20 | 16 | 33 | 27 | |
| Power | 0.3916 | 6.1758 | 20 | 16 | 33 | 27 | |
| Linear | 0.3916 | 6.1758 | 20 | 16 | 33 | 27 | |

^a Selected model in bold.

Plots of the Power model with BMRs of one SD and 10% RD are shown in Figure 1-104 and Figure 1-105, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-106 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-104. Plot of Response by Dose with Fitted Curve for the Selected Model (Power) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 1SD (Constant Variance)




Figure 1-105. Plot of Response by Dose with Fitted Curve for the Selected Model (Power) for Increased Absolute Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%RD (Constant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 20.30524804 | | | | | | | |
| BMDL | 16.22947938 | | | | | | | |
| BMDU | 31.96250624 | | | | | | | |
| AIC | 6.175774284 | | | | | | | |
| Test 4 P-value | 0.391649386 | | | | | | | |
| D.O.F. | 2 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 4 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| g | 3.902142333 | 0.059216216 | 3.78608068 | 4.01820398 | | | | |
| v | 0.011942871 | 1.20E-03 | 0.00958432 | 0.01430142 | | | | |
| n | Bounded | NA | NA | NA | | | | |
| alpha | 0.058807687 | 7.73E-04 | 0.05729203 | 0.06032334 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 10 | 3.902142333 | 3.9 | 3.9 | 0.24250296 | 0.19 | 0.19 | -0.027936363 |
| 21 | 10 | 4.152942628 | 4.1 | 4.1 | 0.24250296 | 0.32 | 0.32 | -0.690380391 |
| 43 | 10 | 4.415685794 | 4.5 | 4.5 | 0.24250296 | 0.25 | 0.25 | 1.099470812 |
| 86 | 10 | 4.929229256 | 4.9 | 4.9 | 0.24250296 | 0.22 | 0.22 | -0.381154207 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | 0.849501122 | 5 | 8.30099776 | | | | | |
| A2 | 2.355544823 | 8 | 11.2889104 | | | | | |
| A3 | 0.849501122 | 5 | 8.30099776 | | | | | |
| fitted | -0.087887142 | 3 | 6.17577428 | | | | | |
| R | -24.92705487 | 2 | 53.8541097 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 54.56519938 | 6 | <0.0001 | | | | | |
| 2 | 3.012087401 | 3 | 0.38976529 | | | | | |
| 3 | 3.012087401 | 3 | 0.38976529 | | | | | |
| 4 | 1.874776528 | 2 | 0.39164939 | | | | | |

Figure 1-106. Details Regarding the Selected Model (Power) for Increased Relative Kidney Weight in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks

1.1.2.2.4 Immune Effects

Data sets identified for BMD modeling for immune effects in a 14-day gavage study ([Munson et al., 1982](#)) include changes in leukocyte count, antibody-forming cells/spleen, and antibody-forming

cells/10⁶ cells in male mice. Data sets for the incidence of thymus necrosis in male and female rats in a 13-week gavage study ([NTP, 1991](#)) were also identified for BMD modeling.

Modeled results were not presented for the antibody-forming cells/spleen or antibody-forming cells/10⁶ cells in male mice data sets ([Munson et al., 1982](#)) because none of the models provided adequate fits to the means (test 4 p-value < 0.1) either assuming constant or nonconstant variance.

1.1.2.2.4.1 Leukocyte Count in CD-1 Male Mice – 14-day Gavage Study

Leukocyte counts were significantly decreased in male mice exposed to 1,2-dichloroethane by gavage daily for 14 days ([Munson et al., 1982](#)). The dose and response data used for the modeling are presented in Table 1-77. Continuous models were used to fit the dose-response data.

A BMR of one SD was chosen according to EPA’s *BMD Technical Guidance* ([U.S. EPA, 2012](#)). A BMR of 10% RD was also selected because EPA considers a 10% change in leukocyte counts to be biologically relevant.

Table 1-77. Leukocytes in Male Mice and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 14-Day Oral Exposure Study

| Dose (mg/kg-day) | Number of Animals | Mean | SD |
|------------------|-------------------|------|------|
| 0 | 12 | 8.24 | 3.26 |
| 4.89 | 10 | 7.60 | 1.64 |
| 48.9 | 10 | 5.76 | 1.55 |

The BMD modeling results for decreased leukocyte count in male mice are summarized in Table 1-78. The constant variance model did not provide adequate fit to the variance data, but the nonconstant variance model did. With the nonconstant variance model applied, all models except the Exponential 5 and Hill models provided adequate fit to the means (test 4 p-value < 0.1). The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC (Exponential 3) was selected.

Table 1-78. Summary of BMD Modeling Results for Decreased Leukocytes in Male Mice Following Oral Exposure to 1,2-Dichloroethane for 14 Days (Nonconstant Variance)^a

| Model | Goodness of Fit | | BMD 1SD (mg/kg-day) | BMDL 1SD (mg/kg-day) | BMD 10%RD (mg/kg-day) | BMDL 10%RD (mg/kg-day) | Basis for Model Selection |
|---------------------|-----------------|-------|---------------------|----------------------|-----------------------|------------------------|---|
| | test 4 p-value | AIC | | | | | |
| Exponential 3 | 0.1556 | 146.8 | 56.1 | 30.9 | 15.2 | 9.75 | All models except the Exponential 5 and Hill models provided adequate fit to the means (test 4 p-value < 0.1). BMDLs for the fit models differed by < 3-fold; therefore, EPA chose the model with the lowest AIC. |
| Exponential 5 | NA | 148.8 | - | - | 3.69 | 0.980 | |
| Hill | NA | 148.8 | - | - | 3.14 | 0.480 | |
| Polynomial Degree 2 | 0.1435 | 147.0 | 55.4 | 34.0 | 17.2 | 12.0 | |
| Power | 0.1435 | 147.0 | 55.8 | 34.0 | 17.3 | 12.0 | |

| Model | Goodness of Fit | | BMD 1SD (mg/kg -day) | BMDL 1SD (mg/kg -day) | BMD 10%RD (mg/kg- day) | BMDL 10%RD (mg/kg- day) | Basis for Model Selection |
|--------|-----------------------|-------|-------------------------------|--------------------------------|---------------------------------|----------------------------------|---------------------------|
| | test 4 p- value | AIC | | | | | |
| Linear | 0.1435 | 147.0 | 55.4 | 34.0 | 17.2 | 12.0 | |

^a Selected model in bold.

Plots of the Exponential 3 model with BMRs of one SD and 10% RD are shown in Figure 1-107 and Figure 1-108, respectively. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-109 (BMD and BMDL shown are for BMR of one SD; the rest is applicable to both BMRs).




Figure 1-107. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Leukocytes in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage for 14 Days and BMR of 1SD (Nonconstant Variance)




Figure 1-108. Plot of Response by Dose with Fitted Curve for the Selected Model (Exponential 3) for Decreased Leukocytes in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage for 14 Days and BMR of 10%RD (Nonconstant Variance)

| Model Results | | | | | | | | |
|--------------------------------|--------------------------|------------------|---------------|---------------|--------------|-----------|-------------|-----------------|
| Benchmark Dose | | | | | | | | |
| BMD | 56.08674721 | | | | | | | |
| BMDL | 30.86793476 | | | | | | | |
| BMDU | 149.2332182 | | | | | | | |
| AIC | 146.8344073 | | | | | | | |
| Test 4 P-value | 0.155640847 | | | | | | | |
| D.O.F. | 1 | | | | | | | |
| Model Parameters | | | | | | | | |
| # of Parameters | 5 | | | | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | | | | |
| a | 8.073279139 | 0.575470972 | 6.94537675 | 9.20118153 | | | | |
| b | 0.006937773 | 2.22E-03 | 0.00257845 | 0.0112971 | | | | |
| d | Bounded | NA | NA | NA | | | | |
| rho | 3.534193135 | 1.97E+00 | -0.3223061 | 7.39069233 | | | | |
| log-alpha | -5.468519023 | 3.87E+00 | -13.048931 | 2.11189326 | | | | |
| Goodness of Fit | | | | | | | | |
| Dose | Size | Estimated Median | Calc'd Median | Observed Mean | Estimated SD | Calc'd SD | Observed SD | Scaled Residual |
| 0 | 12 | 8.073279139 | 8.24 | 8.24 | 2.60238213 | 3.26 | 3.26 | 0.221926671 |
| 4.89 | 10 | 7.803981293 | 7.6 | 7.6 | 2.45095372 | 1.64 | 1.64 | -0.263181422 |
| 48.9 | 10 | 5.750590964 | 5.76 | 5.76 | 1.42893177 | 1.55 | 1.55 | 0.020822538 |
| Likelihoods of Interest | | | | | | | | |
| Model | Log Likelihood* | # of Parameters | AIC | | | | | |
| A1 | -71.42798259 | 4 | 150.855965 | | | | | |
| A2 | -67.34059772 | 6 | 146.681195 | | | | | |
| A3 | -68.40916542 | 5 | 146.818331 | | | | | |
| fitted | -69.41720367 | 4 | 146.834407 | | | | | |
| R | -74.55945884 | 2 | 153.118918 | | | | | |
| Tests of Interest | | | | | | | | |
| Test | -2*Log(Likelihood Ratio) | Test df | p-value | | | | | |
| 1 | 14.43772225 | 4 | 0.00602144 | | | | | |
| 2 | 8.174769753 | 2 | 0.01678307 | | | | | |
| 3 | 2.137135415 | 1 | 0.14377013 | | | | | |
| 4 | 2.016076497 | 1 | 0.15564085 | | | | | |

Figure 1-109. Details Regarding the Selected Model (Exponential 3) for Decreased Leukocytes in Male Mice Exposed to 1,2-Dichloroethane Via Oral Gavage for 14 Days

1.1.2.2.4.2 Thymus Necrosis in F344 Male Rats – One Time per Day, 5 days per Week

There was an increased incidence of thymus necrosis in male rats exposed to 1,2-dichloroethane by gavage for 13 weeks (5 days per week) (NTP, 1991). The administered doses were duration adjusted to estimate an equivalent oral dose for animals exposed for 7 days per week. The dose and response data used for the modeling are presented in Table 1-79. Dichotomous models were used to fit dose-response data.

A BMR of 10% ER was chosen according to *BMD Technical Guidance* (U.S. EPA, 2012).

Table 1-79. Incidence of Thymus Necrosis in Male Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study

| Adjusted Dose (mg/kg-day) | Number of Animals | Incidence |
|---------------------------|-------------------|-----------|
| 0 | 10 | 0 |
| 86 | 10 | 0 |
| 171 | 10 | 4 |
| 343 | 10 | 10 |

The BMD modeling results for increased incidence of thymus necrosis in male rats are summarized in Table 1-80. All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Multistage 1-degree/Quantal Linear model. The BMD computation failed for the Weibull model because the lower limit included zero; therefore, this model was unusable. BMDLs for the remaining models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree).

Table 1-80. Summary of BMD Modeling Results for Increased Incidence of Thymus Necrosis in Male Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks^a

| Model | Goodness of Fit | | BMD 10%ER (mg/kg- day) | BMDL 10%ER (mg/kg-day) | Basis for Model Selection |
|---------------------|-----------------|--------------|---------------------------------|------------------------------|---|
| | p- value | AIC | | | |
| Dichotomous Hill | 1.000 | 15.46 | 160 | 100 | All models provided adequate fit to the data (chi-square p-value > 0.1) except for the Multistage 1-degree/Quantal Linear model. The BMD computation failed for the Weibull model and was unusable. BMDLs differed by < 3-fold; therefore, EPA chose the model with the lowest AIC (Multistage Degree 3). |
| Gamma | 0.9979 | 15.54 | 130 | 92 | |
| Log-Logistic | 1.000 | 15.46 | 160 | 100 | |
| Multistage 3 | 0.9335 | 15.09 | 100 | 59 | |
| Multistage 2 | 0.4338 | 20.47 | 71 | 44 | |
| Multistage 1 | 0.0490 | 28.56 | 27 | 17 | |
| Weibull | 0.9988 | 15.52 | 140 | 0 | |
| Logistic | 1.000 | 15.46 | 150 | 93 | |
| Log-Probit | 1.000 | 17.46 | 160 | 97 | |
| Probit | 1.000 | 17.46 | 150 | 87 | |
| Quantal Linear | 0.0490 | 28.56 | 27 | 17 | |

^a Selected model in bold.

A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-110. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-111.




Figure 1-110. Plot of Response by Dose with Fitted Curve for the Selected Model (Multistage 3-Degree) for Increased Incidence of Thymus Necrosis in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 100.8873865 | | | | |
| BMDL | 59.30171288 | | | | |
| BMDU | 124.622876 | | | | |
| AIC | 15.08642199 | | | | |
| P-value | 0.933531247 | | | | |
| D.O.F. | 4 | | | | |
| Chi ² | 0.836191799 | | | | |
| Slope Factor | 0.001686292 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | Bounded | NA | NA | NA | |
| b2 | Bounded | NA | NA | NA | |
| b3 | Bounded | NA | NA | NA | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 86 | 0.063178343 | 0.631783434 | 0 | 10 | -0.821213 |
| 171 | 0.401330243 | 4.013302431 | 4 | 10 | -0.008582 |
| 343 | 0.984084626 | 9.840846256 | 10 | 10 | 0.4021538 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -6.73011667 | 4 | - | - | NA |
| Fitted Model | -7.543210994 | 0 | 1.62618865 | 4 | 0.8040777 |
| Reduced Model | -25.89786556 | 1 | 38.3354978 | 3 | <0.0001 |

Figure 1-111. Details Regarding the Selected Model (Multistage 3-Degree) for Increased Incidence of Thymus Necrosis in Male Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks

1.1.2.2.4.3 Thymus Necrosis in F344 Female Rats – 13-Week Gavage

There was an increased incidence of thymus necrosis in female rats exposed to 1,2-dichloroethane by gavage for 13 weeks (5 days per week) (NTP, 1991). The administered doses were duration adjusted to estimate an equivalent oral dose for animals exposed for 7 days per week. The dose and response data used for the modeling are presented in Table 1-81. Dichotomous models were used to fit the dose-response data.

A BMR of 10% ER was chosen according to *BMD Technical Guidance* (U.S. EPA, 2012).

Table 1-81. Increased Incidence of Thymus Necrosis in Female Rats and Associated Doses Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 13-Week Oral Exposure Study

| Adjusted Dose (mg/kg-day) | Number of Animals | Incidence |
|---------------------------|-------------------|-----------|
| 0 | 10 | 0 |
| 107 | 10 | 0 |
| 214 | 10 | 5 |

The BMD modeling results for increased incidence of thymus necrosis in female rats are summarized in Table 1-82. All models provided adequate fit to the data (chi-square p-value > 0.1). BMDLs were not sufficiently close (differed by > 3-fold); therefore, the model with the lowest BMDL was selected (Multistage 1-degree).

This data set is not well suited for BMD modeling; there is a single non-zero data point and no data to inform the shape of the curve at the region of interest, as defined by the BMR. As a result, the different models provide a broad range of BMD and BMDL estimates. Selection of the low end of the range to represent the BMDL is not among the more realistic possibilities and in this case involves extrapolation below the range of observation to generate a BMD estimate well below the lowest tested dose.

Table 1-82. Summary of BMD Modeling Results for Increased Incidence of Thymus Necrosis in Female Rats Following Oral Exposure to 1,2-Dichloroethane for 13 Weeks^a

| Model | Goodness of Fit | | BMD 10%ER (mg/kg-day) | BMDL 10%ER (mg/kg-day) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------------|------------------------|--|
| | p-value | AIC | | | |
| Dichotomous Hill | 1.000 | 15.86 | 189 | 97.1 | All models provided adequate fit to the data (chi-square p-value > 0.1). BMDLs differed by > 3-fold; therefore, EPA chose the model with the lowest BMDL. NOTE: This data set is not well suited for BMD modeling and the results should be interpreted with caution. |
| Gamma | 0.9780 | 15.95 | 156 | 95.7 | |
| Log-Logistic | 1.000 | 15.86 | 189 | 97.1 | |
| Multistage 2 | 0.4109 | 18.83 | 97.2 | 41.9 | |
| Multistage 1 | 0.1740 | 21.10 | 55.6 | 28.7 | |
| Weibull | 1.000 | 15.86 | 193 | 96.1 | |
| Logistic | 0.9994 | 15.87 | 188 | 107 | |
| Log-Probit | 0.9996 | 17.86 | 199 | 98.2 | |
| Probit | 1.000 | 17.86 | 193 | 102 | |
| Quantal Linear | 0.1740 | 21.10 | 55.6 | 28.7 | |

^a Selected model in bold.

A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-112. Additional modeling details, including model parameters, goodness of fit at each dose, and log likelihood are shown in Figure 1-113.




Figure 1-112. Plot of Response by Dose with Fitted Curve for the Selected Model (Quantal Linear) for Increased Incidence of Thymus Necrosis in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks and BMR of 10%ER

| Model Results | | | | | |
|-----------------------------|-----------------------|-----------------|------------|------------|-----------------|
| Benchmark Dose | | | | | |
| BMD | 55.60812223 | | | | |
| BMDL | 28.68887964 | | | | |
| BMDU | 129.2673929 | | | | |
| AIC | 21.09542551 | | | | |
| P-value | 0.173995734 | | | | |
| D.O.F. | 2 | | | | |
| Chi ² | 3.49744899 | | | | |
| Slope Factor | 0.003485671 | | | | |
| Model Parameters | | | | | |
| # of Parameters | 2 | | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf | |
| g | Bounded | NA | NA | NA | |
| b1 | 0.001894697 | 0.182567022 | -0.3559301 | 0.35971949 | |
| Goodness of Fit | | | | | |
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 1.523E-07 | 0 | 10 | -0.00039 |
| 107 | 0.183503423 | 1.835034229 | 0 | 10 | -1.499149 |
| 214 | 0.333333329 | 3.333333294 | 5 | 10 | 1.118034 |
| Analysis of Deviance | | | | | |
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -6.931471806 | 3 | - | - | NA |
| Fitted Model | -9.547712753 | 1 | 5.23248189 | 2 | 0.073077 |
| Reduced Model | -13.51683627 | 1 | 13.1707289 | 2 | 0.0013804 |

Figure 1-113. Details Regarding the Selected Model (Quantal Linear) for Increased Incidence of Thymus Necrosis in Female Rats Exposed to 1,2-Dichloroethane Via Oral Gavage for 13 Weeks

1.2 Cancer Endpoints

The inhalation unit risk for 1,2-dichloroethane was based on an inhalation study for 1,2-dichloroethane by [Nagano et al. \(2006\)](#). EPA conducted BMD modeling on these data as described below.

The BMD modeling of cancer incidence data was conducted with the EPA's BMD software (BMDS, version 3.3). Modeled concentrations were in units of ppm. For these data, the Multistage model was fit to the incidence data using a BMR of 10% ER. The Multistage cancer model was run for all polynomial degrees up to $n-1$ (where n is the number of dose groups including control). Adequacy of model fit was judged based on the chi-square goodness-of-fit p-value ($p > 0.1$), magnitude of scaled residuals in the vicinity of the BMR, and visual inspection of the model fit. Among all models providing adequate fit, the BMDL from the model with the lowest AIC was selected if the BMDLs were sufficiently close (< 3 -fold); if the BMDLs were not sufficiently close (> 3 -fold), model-dependence is indicated, and the model with the lowest reliable BMDL was selected.

Where applicable, the MS Combo model was used to evaluate the combined cancer risk of tumors observed in multiple tissues in a test group, assuming that the tumors in the different tissues occurred independently. MS Combo was run using the incidence data for the individual tumors and the polydegrees identified in the model runs for the individual tumors.

1.2.1 Rat Data

1.2.1.1 Tumor Incidence in Male Rats

1.2.1.1.1 Subcutaneous Fibromas in Male Rats

Male rats exhibited a significantly increased trend for the incidence of subcutaneous fibromas in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). The concentration and response data used for the modeling are presented in Table 1-83.

Table 1-83. Increased Incidence of Subcutaneous Fibromas in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 6 |
| 2 | 50 | 9 |
| 7 | 50 | 12 |
| 29 | 50 | 15 |

The BMD modeling results for subcutaneous fibromas in male rats are summarized in Table 1-84. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The Multistage 2- and 3-degree models converged on the 1-degree model; therefore, the 1-degree Multistage model was selected as the more parsimonious choice. A plot of the Multistage 1 model with a BMR of 10% ER is shown in Figure 1-114. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-115.

Table 1-84. Summary of BMD Modeling Results for Increased Incidence of Subcutaneous Fibromas in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.6229 | 205.2 | 14 | 7.3 | 0.014 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The 2- and 3-degree models converged on the 1-degree model; |
| Multistage 2 | 0.6229 | 205.2 | 14 | 7.3 | 0.014 | |
| Multistage 1 | 0.6229 | 205.2 | 14 | 7.3 | 0.014 | |

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|--------------------------------------|-----------------|-----|-----------------------|------------------------|------------------------------|---|
| | p-value | AIC | | | | |
| | | | | | | therefore, EPA chose the 1-degree Multistage model. |
| ^a Selected model in bold. | | | | | | |




Figure 1-114. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Subcutaneous Fibromas in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 14.31033134 |
| BMDL | 7.324332663 |
| BMDU | Infinity |
| AIC | 205.197232 |
| P-value | 0.622942744 |
| D.O.F. | 2 |
| Chi ² | 0.946601335 |
| Slope Factor | 0.013653121 |

| Model Parameters | | | | |
|------------------|-------------|-------------|--------------|-------------|
| # of Parameters | 2 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.155527448 | 1.70E-02 | 0.122295876 | 0.188759019 |
| b1 | 0.007362549 | 0.108083178 | -0.204476589 | 0.219201687 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.155527448 | 7.776372388 | 6 | 50 | -0.637008884 |
| 2 | 0.167871284 | 8.393564185 | 9 | 50 | 0.209320438 |
| 7 | 0.19794724 | 9.89736201 | 12 | 50 | 0.668351272 |
| 29 | 0.317884967 | 15.89424837 | 15 | 50 | -0.224304589 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -100.0131354 | 4 | - | - | NA |
| Fitted Model | -100.598616 | 2 | 1.170961199 | 2 | 0.556838181 |
| Reduced Model | -102.7913341 | 1 | 4.385436226 | 3 | 0.222739593 |

Figure 1-115. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Subcutaneous Fibromas in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.1.2 Mammary Gland Fibroadenomas in Male Rats

Male rats exhibited significantly increased incidences of mammary gland fibroadenomas in a 2-year inhalation toxicity study of 1,2-dichloroethane ([Nagano et al., 2006](#)). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-85.

Table 1-85. Increased Incidence of Mammary Gland Fibroadenomas in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 0 |
| 2 | 50 | 0 |
| 7 | 50 | 1 |
| 29 | 50 | 5 |

The BMD modeling results for mammary gland fibroadenomas in male rats are summarized in Table 1-86. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The Multistage 3-degree model converged on the 2-degree model. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 1-degree). A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-116. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-117.

Table 1-86. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Fibroadenomas in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|-------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.8739 | 46.8 | 29 | 18 | 0.0056 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The 3-degree model converged on the 2-degree model. The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 1-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.8739 | 46.8 | 29 | 18 | 0.0056 | |
| Multistage 1 | 0.9425 | 45.0 | 32 | 17 | 0.0057 | |

^a Selected model in bold.




Figure 1-116. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Fibroadenomas in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 32.00911617 |
| BMDL | 17.49222195 |
| BMDU | 68.22514663 |
| AIC | 45.03571882 |
| P-value | 0.942486739 |
| D.O.F. | 3 |
| Chi ² | 0.389090832 |
| Slope Factor | 0.005716827 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 2 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | Bounded | NA | NA | NA |
| b1 | 0.003291578 | 3.90E-02 | -0.073102918 | 0.079686075 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-07 | 0 | 50 | -0.001 |
| 2 | 0.00656155 | 0.328077513 | 0 | 50 | -0.573 |
| 7 | 0.022777645 | 1.138882259 | 1 | 50 | -0.130 |
| 29 | 0.091041451 | 4.55207255 | 5 | 50 | 0.210 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -21.15610433 | 4 | - | - | NA |
| Fitted Model | -21.51785941 | 1 | 0.723510151 | 3 | 0.867660684 |
| Reduced Model | -26.94843364 | 1 | 10.86114845 | 3 | 0.012500888 |

Figure 1-117. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Fibroadenomas in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.1.3 Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats

Male rats exhibited significantly increased incidences of mammary gland adenomas and fibroadenomas (combined) in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-87.

Table 1-87. Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 1 |
| 2 | 50 | 2 |
| 7 | 50 | 1 |
| 29 | 50 | 7 |

The BMD modeling results for mammary gland adenomas and fibroadenomas (combined) in male rats are summarized in Table 1-88. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree). A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-118. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-119.

Table 1-88. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.7617 | 81.43 | 27 | 15 | 0.0065 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 3-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.7116 | 81.58 | 27 | 15 | 0.0066 | |
| Multistage 1 | 0.5722 | 82.24 | 27 | 14 | 0.0072 | |

^a Selected model in bold.




Figure 1-118. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 27.49900097 |
| BMDL | 15.31882395 |
| BMDU | 61.22780582 |
| AIC | 81.42793799 |
| P-value | 0.761735667 |
| D.O.F. | 2 |
| Chi ² | 0.544311354 |
| Slope Factor | 0.006527916 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 4 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.026277252 | 3.23E-02 | -0.036979536 | 0.089534039 |
| b1 | Bounded | NA | NA | NA |
| b2 | Bounded | NA | NA | NA |
| b3 | 5.06672E-06 | 5.86E-02 | -0.114805797 | 0.11481593 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.026277252 | 1.313862577 | 1 | 50 | -0.273819621 |
| 2 | 0.026316719 | 1.31583597 | 2 | 50 | 0.596429835 |
| 7 | 0.027968001 | 1.398400056 | 1 | 50 | -0.336902066 |
| 29 | 0.13946499 | 6.973249521 | 7 | 50 | 0.010130105 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -38.44929297 | 4 | - | - | NA |
| Fitted Model | -38.71396899 | 2 | 0.529352051 | 2 | 0.767454546 |
| Reduced Model | -42.59643946 | 1 | 7.764940939 | 3 | 0.051127864 |

Figure 1-119. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.1.4 Peritoneal Mesothelioma in Male Rats

Male rats exhibited a significantly increased trend for incidence of peritoneal mesothelioma in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). The concentrations and response data used for the modeling are presented in Table 1-89.

Table 1-89. Increased Incidence of Peritoneal Mesothelioma in Male Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 1 |
| 2 | 50 | 1 |
| 7 | 50 | 1 |
| 29 | 50 | 5 |

The BMD modeling results for peritoneal mesothelioma in male rats are summarized in Table 1-90. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree). A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-120. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-121.

Table 1-90. Summary of BMD Modeling Results for Increased Incidence of Peritoneal Mesothelioma in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------------|------------------------|------------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.9989 | 65.92 | 31 | 19 | 0.0052 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 3-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.9830 | 65.96 | 32 | 19 | 0.0052 | |
| Multistage 1 | 0.8132 | 66.39 | 38 | 18 | 0.0055 | |

^a Selected model in bold.




Figure 1-120. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Peritoneal Mesothelioma in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 31.09224433 |
| BMDL | 19.32721096 |
| BMDU | Infinity |
| AIC | 65.92230974 |
| P-value | 0.998892256 |
| D.O.F. | 2 |
| Chi ² | 0.002216716 |
| Slope Factor | 0.005174052 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 4 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.019616475 | 3.73E-02 | -0.053527977 | 0.092760928 |
| b1 | Bounded | NA | NA | NA |
| b2 | Bounded | NA | NA | NA |
| b3 | 3.50527E-06 | 4.87E-02 | -0.095482039 | 0.095489049 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.019616475 | 0.980823761 | 1 | 50 | 0.01936279 |
| 2 | 0.019643967 | 0.982198346 | 1 | 50 | 0.017962251 |
| 7 | 0.02079449 | 1.039724502 | 1 | 50 | -0.038958239 |
| 29 | 0.099946886 | 4.997344318 | 5 | 50 | 0.001187972 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -30.96001566 | 4 | - | - | NA |
| Fitted Model | -30.96115487 | 2 | 0.002278419 | 2 | 0.998861439 |
| Reduced Model | -33.58882955 | 1 | 5.255349352 | 3 | 0.154026085 |

Figure 1-121. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Peritoneal Mesothelioma in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.1.5 Combined Mammary Gland, Subcutaneous, and Peritoneal Tumors in Male Rats

Male rats exhibited significantly increased incidences of subcutaneous fibromas, mammary gland adenomas and fibroadenomas, and peritoneal mesothelioma in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). The MS Combo model was applied to the incidence data for the individual tumors and polydegrees identified for the individual tumor models, shown above (Sections 1.2.1.1.1 through 1.2.1.1.4), using a BMR of 10% ER.

The BMD Multistage Cancer/Multi-tumor modeling results for combined subcutaneous fibromas, mammary gland adenomas and fibroadenomas, and peritoneal mesothelioma in male rats are summarized in Table 1-91. A plot of the dose-response curves for the incidence of individual tumor types with a BMR of 10% ER is shown in Figure 1-122. Additional modeling details, including the cancer slope factor and log likelihood are shown in Figure 1-123.

Table 1-91. Summary of BMD Multi-Tumor (MS Combo) Modeling Results for Increased Incidence of Subcutaneous Fibromas, Mammary Gland Adenomas and Fibroadenomas, and Peritoneal Mesothelioma (Combined) in Male Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay

| Model | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) |
|---|-----------------|------------------|------------------------|
| Multi-tumor (MS Combo) | 12 | 5.3 | 0.019 |
| Subcutaneous fibroma Multistage 1 | 14 | 7.3 | 0.014 |
| Mammary gland adenoma and fibroadenoma Multistage 3 | 27 | 15 | 0.0065 |
| Peritoneal mesothelioma Multistage 3 | 31 | 19 | 0.0052 |




Figure 1-122. Plot of Response by Concentration with Fitted Curve for Selected Models used for the Multi-Tumor (MS Combo) Model for Increased Incidence of Subcutaneous Fibromas, Mammary Gland Adenomas and Fibroadenomas, and Peritoneal Mesothelioma (Combined) in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)




Figure 1-123. Details Regarding the Multi-Tumor (MS Combo) Model for the Increased Incidence of Subcutaneous Fibromas, Mammary Gland Adenomas and Fibroadenomas, and Peritoneal Mesothelioma (Combined) in Male Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.2 Tumor Incidence in Female Rats

1.2.1.2.1 Subcutaneous Fibromas in Female Rats

Female rats exhibited significantly increased incidences of subcutaneous fibromas in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). The concentration and response data used for the modeling are presented in Table 1-92.

Table 1-92. Increased Incidence of Subcutaneous Fibromas in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 0 |
| 2 | 50 | 0 |
| 7 | 50 | 1 |
| 29 | 50 | 5 |

The BMD modeling results for subcutaneous fibromas in female rats are summarized in Table 1-93. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The Multistage 3-degree model converged on the 2-degree model. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 1-degree). A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-124. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-125.

Table 1-93. Summary of BMD Modeling Results for Increased Incidence of Subcutaneous Fibromas in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------------|------------------------|------------------------------|--|
| | p-value | AIC | | | | |
| Multistage 3 | 0.8739 | 46.75 | 29 | 18 | 0.0056 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The 3-degree model converged on the 2-degree model. The BMDs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 1-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.8739 | 46.75 | 29 | 18 | 0.0056 | |
| Multistage 1 | 0.9425 | 45.04 | 32 | 17 | 0.0057 | |

^a Selected model in bold.




Figure 1-124. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Subcutaneous Fibromas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 32.00911617 |
| BMDL | 17.49222195 |
| BMDU | 68.22514663 |
| AIC | 45.03571882 |
| P-value | 0.942486739 |
| D.O.F. | 3 |
| Chi ² | 0.389090832 |
| Slope Factor | 0.005716827 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 2 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | Bounded | NA | NA | NA |
| b1 | 0.003291578 | 3.90E-02 | -0.073102918 | 0.079686075 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 1.523E-08 | 7.61499E-07 | 0 | 50 | -0.000872639 |
| 2 | 0.00656155 | 0.328077513 | 0 | 50 | -0.572780511 |
| 7 | 0.022777645 | 1.138882259 | 1 | 50 | -0.130138968 |
| 29 | 0.091041451 | 4.55207255 | 5 | 50 | 0.209943818 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -21.15610433 | 4 | - | - | NA |
| Fitted Model | -21.51785941 | 1 | 0.723510151 | 3 | 0.867660684 |
| Reduced Model | -26.94843364 | 1 | 10.86114845 | 3 | 0.012500888 |

Figure 1-125. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Subcutaneous Fibromas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.2.2 Mammary Gland Adenomas in Female Rats

Female rats exhibited significantly increased incidences of mammary gland adenomas in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). The concentration and response data used for the modeling are presented in Table 1-94.

Table 1-94. Increased Incidence of Mammary Gland Adenomas in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 3 |
| 2 | 50 | 5 |
| 7 | 50 | 5 |
| 29 | 50 | 11 |

The BMD modeling results for mammary gland adenomas in female rats are summarized in Table 1-95. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The Multistage 3-degree model converged on the 1-degree model. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected. The 1-degree Multistage model was selected as the more parsimonious choice. A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-126. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-127.

Table 1-95. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenomas in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|--|
| | p-value | AIC | | | | |
| Multistage 3 | 0.8523 | 144.7 | 18 | 9.4 | 0.011 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The 3-degree model converged on the 1-degree model. The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 1-degree Multistage model, which had the lowest AIC and was the more parsimonious choice. |
| Multistage 2 | 0.5709 | 146.7 | 18 | 9.4 | 0.011 | |
| Multistage 1 | 0.8516 | 144.7 | 18 | 9.4 | 0.011 | |

^a Selected model in bold.




Figure 1-126. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 17.53354728 |
| BMDL | 9.407667481 |
| BMDU | 58.29776202 |
| AIC | 144.7453231 |
| P-value | 0.851617435 |
| D.O.F. | 2 |
| Chi ² | 0.321235746 |
| Slope Factor | 0.010629627 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 2 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.070489692 | 2.44E-02 | 0.022654769 | 0.118324615 |
| b1 | 0.006009082 | 7.59E-02 | -0.142761504 | 0.154779667 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.070489692 | 3.524484591 | 3 | 50 | -0.279373318 |
| 2 | 0.081593839 | 4.079691958 | 5 | 50 | 0.455637574 |
| 7 | 0.108777313 | 5.438865655 | 5 | 50 | -0.188181718 |
| 29 | 0.219141052 | 10.95705262 | 11 | 50 | 0.012974475 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -70.20207154 | 4 | - | - | NA |
| Fitted Model | -70.37266157 | 2 | 0.341180063 | 2 | 0.843167175 |
| Reduced Model | -73.38499825 | 1 | 6.024673375 | 3 | 0.110415963 |

Figure 1-127. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.2.3 Mammary Gland Fibroadenomas in Female Rats

Female rats exhibited significantly increased incidences of mammary gland fibroadenomas in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). The concentration and response data used for the modeling are presented in Table 1-96.

Table 1-96. Increased Incidence of Mammary Gland Fibroadenomas in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 4 |
| 2 | 50 | 1 |
| 7 | 50 | 6 |
| 29 | 50 | 13 |

The BMD modeling results for mammary gland fibroadenomas in female rats are summarized in Table 1-97. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The Multistage 3-degree model converged on the 2-degree model. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 1-degree). A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-128. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-129.

Table 1-97. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Fibroadenomas in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.1157 | 140.8 | 15 | 7.8 | 0.013 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The 3-degree model converged on the 2-degree model. The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 1-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.1157 | 140.8 | 15 | 7.8 | 0.013 | |
| Multistage 1 | 0.2797 | 138.9 | 13 | 7.7 | 0.013 | |

^a Selected model in bold.




Figure 1-128. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Fibroadenomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 12.58447242 |
| BMDL | 7.741278443 |
| BMDU | 26.11089254 |
| AIC | 138.9247592 |
| P-value | 0.279716095 |
| D.O.F. | 2 |
| Chi ² | 2.547960275 |
| Slope Factor | 0.012917763 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 2 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.051585929 | 2.83E-02 | -0.003958966 | 0.107130823 |
| b1 | 0.008372264 | 7.82E-02 | -0.144863436 | 0.161607964 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.051585929 | 2.579296433 | 4 | 50 | 0.884612032 |
| 2 | 0.067334456 | 3.3667228 | 1 | 50 | -1.289863384 |
| 7 | 0.105571164 | 5.278558204 | 6 | 50 | 0.314010118 |
| 29 | 0.256033853 | 12.80169265 | 13 | 50 | 0.055424925 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -65.83951967 | 4 | - | - | NA |
| Fitted Model | -67.46237958 | 2 | 3.245719818 | 2 | 0.197333535 |
| Reduced Model | -73.38499825 | 1 | 11.84523735 | 3 | 0.007932559 |

Figure 1-129. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Fibroadenomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.2.4 Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats

Female rats exhibited significantly increased incidences of mammary gland adenomas and fibroadenomas (combined) in a 2-year inhalation toxicity study of 1,2-dichloroethane ([Nagano et al., 2006](#)). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-98.

Table 1-98. Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 7 |
| 2 | 50 | 6 |
| 7 | 50 | 11 |
| 29 | 50 | 22 |

The BMD modeling results for mammary gland adenomas and fibroadenomas (combined) in female rats are summarized in Table 1-99. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The Multistage 3-degree model converged on the 2-degree model. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 1-degree). A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-130. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-131.

Table 1-99. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.5222 | 205.0 | 7.5 | 4.5 | 0.022 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The 3-degree model converged on the 2-degree model. The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 1-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.5222 | 205.0 | 7.5 | 4.5 | 0.022 | |
| Multistage 1 | 0.8084 | 203.0 | 7.5 | 4.5 | 0.022 | |

^a Selected model in bold.




Figure 1-130. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 6.836211264 |
| BMDL | 4.501066127 |
| BMDU | 12.30383058 |
| AIC | 202.9818151 |
| P-value | 0.808367209 |
| D.O.F. | 2 |
| Chi ² | 0.425477713 |
| Slope Factor | 0.022216959 |

| Model Parameters | | | | |
|------------------|-------------|-------------|--------------|-------------|
| # of Parameters | 2 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.121858132 | 1.93E-02 | 0.083957729 | 0.159758535 |
| b1 | 0.015412121 | 0.125543163 | -0.230647959 | 0.261472202 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.121858132 | 6.092906583 | 7 | 50 | 0.367485117 |
| 2 | 0.148513267 | 7.425663362 | 6 | 50 | -0.523177864 |
| 7 | 0.211664854 | 10.5832427 | 11 | 50 | 0.12810729 |
| 29 | 0.438362491 | 21.91812455 | 22 | 50 | 0.017488478 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -99.2363119 | 4 | - | - | NA |
| Fitted Model | -99.49090755 | 2 | 0.509191289 | 2 | 0.775229903 |
| Reduced Model | -107.8552683 | 1 | 16.72872151 | 3 | 0.000803582 |

Figure 1-131. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas and Fibroadenomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.2.5 Mammary Gland Adenocarcinomas in Female Rats

Female rats exhibited a significantly increased trend for the incidence of mammary gland adenocarcinomas in a 2-year inhalation toxicity study of 1,2-dichloroethane ([Nagano et al., 2006](#)). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation dose for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-100.

Table 1-100. Increased Incidence of Mammary Gland Adenocarcinomas in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 1 |
| 2 | 50 | 2 |
| 7 | 50 | 0 |
| 29 | 50 | 5 |

The BMD modeling results for mammary gland adenocarcinomas in female rats are summarized in Table 1-101. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree). A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-132. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-133.

Table 1-101. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenocarcinomas in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.3593 | 66.04 | 31 | 23 | 0.0043 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 3-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.3371 | 66.36 | 33 | 22 | 0.0045 | |
| Multistage 1 | 0.2854 | 67.32 | 44 | 20 | 0.0050 | |

^a Selected model in bold.




Figure 1-132. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenocarcinomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 31.32763273 |
| BMDL | 23.02143452 |
| BMDU | Infinity |
| AIC | 66.03592535 |
| P-value | 0.35930595 |
| D.O.F. | 2 |
| Chi ² | 2.04716205 |
| Slope Factor | 0.004343778 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 4 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.020074114 | 3.65E-02 | -0.051434256 | 0.091582483 |
| b1 | Bounded | NA | NA | NA |
| b2 | Bounded | NA | NA | NA |
| b3 | 3.42685E-06 | 4.79E-02 | -0.093905886 | 0.09391274 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.020074114 | 1.003705676 | 1 | 50 | -0.003698829 |
| 2 | 0.020100978 | 1.005048882 | 2 | 50 | 0.992448894 |
| 7 | 0.021225251 | 1.061262562 | 0 | 50 | -1.030175986 |
| 29 | 0.098644728 | 4.932236411 | 5 | 50 | 0.030512266 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -29.55331172 | 4 | - | - | NA |
| Fitted Model | -31.01796267 | 2 | 2.929301905 | 2 | 0.231158663 |
| Reduced Model | -33.58882955 | 1 | 5.141733748 | 3 | 0.161708073 |

Figure 1-133. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenocarcinomas in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.2.6 Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats

Female rats exhibited significantly increased incidences of mammary gland adenomas, fibroadenomas, and adenocarcinomas (combined) in a 2-year inhalation toxicity study of 1,2-dichloroethane ([Nagano et al., 2006](#)). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation dose for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). The concentration and response data used for the modeling are presented in Table 1-102.

Table 1-102. Increased Incidence of Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 50 | 8 |
| 2 | 50 | 8 |
| 7 | 50 | 11 |
| 29 | 50 | 25 |

The BMD modeling results for mammary gland adenomas, fibroadenomas, and adenocarcinomas (combined) in female rats are summarized in Table 1-103. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The Multistage 3-degree model converged on the 2-degree model. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 1-degree). A plot of the Multistage 1-degree model with a BMR of 10% ER is shown in Figure 1-134. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-135.

Table 1-103. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------------|------------------------|------------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.8383 | 216.0 | 9.2 | 4.1 | 0.025 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The 3-degree model converged on the 2-degree model. The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 1-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.8383 | 216.0 | 9.2 | 4.1 | 0.025 | |
| Multistage 1 | 0.8714 | 214.3 | 5.9 | 4.0 | 0.025 | |

^a Selected model in bold.




Figure 1-134. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 5.931093454 |
| BMDL | 3.976641889 |
| BMDU | 10.25361658 |
| AIC | 214.2892161 |
| P-value | 0.871412464 |
| D.O.F. | 2 |
| Chi ² | 0.275279724 |
| Slope Factor | 0.025146846 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 2 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.141490568 | 1.79E-02 | 0.106338916 | 0.176642219 |
| b1 | 0.017764096 | 0.1380138 | -0.252737984 | 0.288266175 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.141490568 | 7.074528384 | 8 | 50 | 0.34794801 |
| 2 | 0.171456388 | 8.572819379 | 8 | 50 | -0.195639194 |
| 7 | 0.241874456 | 12.09372282 | 11 | 50 | -0.314504792 |
| 29 | 0.487121544 | 24.35607719 | 25 | 50 | 0.130475851 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -104.969745 | 4 | - | - | NA |
| Fitted Model | -105.1446081 | 2 | 0.349726056 | 2 | 0.839572011 |
| Reduced Model | -114.6113834 | 1 | 18.93355072 | 3 | 0.000282186 |

Figure 1-135. Details Regarding the Selected Model (Multistage 1-Degree) for the Increased Incidence of Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.1.2.7 Combined Mammary Gland and Subcutaneous Tumors in Female Rats

Female rats exhibited significantly increased incidences of subcutaneous fibromas and mammary gland adenomas, fibroadenomas, and adenocarcinomas in a 2-year inhalation toxicity study of 1,2-dichloroethane ([Nagano et al., 2006](#)). The MS Combo model was applied to the incidence data for the individual tumors and polydegrees identified for the individual tumor models, shown above (Sections 1.2.1.2.1 through 1.2.1.2.6), using a BMR of 10% ER.

The BMD Multistage Cancer/Multi-tumor modeling results for combined subcutaneous fibromas and mammary gland adenomas, fibroadenomas, and adenocarcinomas in female rats are summarized in

Table 1-104. A plot of the dose-response curves for the incidence of individual tumor types with a BMR of 10% ER is shown in Figure 1-136. Additional modeling details, including the cancer slope factor and log likelihood are shown in Figure 1-137.

Table 1-104. Summary of BMD Multi-Tumor (MS Combo) Modeling Results for Increased Incidence of Subcutaneous Fibromas and Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay

| Model | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) |
|--|-----------------|------------------|------------------------|
| Multi-tumor (MS Combo) | 5.0 | 3.5 | 0.029 |
| Subcutaneous fibroma Multistage 1 | 32 | 17 | 0.0057 |
| Mammary gland adenoma, fibroadenoma, and adenocarcinoma Multistage 1 | 5.9 | 4.0 | 0.025 |




Figure 1-136. Plot of Response by Concentration with Fitted Curve for Selected Models Used for the Multi-Tumor (MS Combo) Model for Increased Incidence of Subcutaneous Fibromas and Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)




Figure 1-137. Details Regarding the Multi-Tumor (MS Combo) Model for the Increased Incidence of Subcutaneous Fibromas and Mammary Gland Adenomas, Fibroadenomas, and Adenocarcinomas (Combined) in Female Rats Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2 Mouse Data

1.2.2.1 Bronchiolo-Alveolar Adenomas in Female Mice

Female mice exhibited a significantly increased trend for the incidence of bronchiolo-alveolar adenomas in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* (U.S. EPA, 2012). The concentration and response data used for the modeling are presented in Table 1-105.

Table 1-105. Increased Incidence of Bronchiolo-Alveolar Adenomas in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 49 | 4 |
| 2 | 50 | 1 |
| 5 | 50 | 3 |
| 16 | 50 | 8 |

The BMD modeling results for bronchiolo-alveolar adenomas in female mice are summarized in Table 1-106. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree). A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-138. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-139.

Table 1-106. Summary of BMD Modeling Results for Increased Incidence of Bronchiolo-Alveolar Adenomas in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------------|------------------------|------------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.4013 | 110.3 | 15 | 9.4 | 0.011 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 3-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.3754 | 110.4 | 15 | 9.3 | 0.011 | |
| Multistage 1 | 0.2359 | 111.5 | 17 | 8.1 | 0.012 | |

^a Selected model in bold.




Figure 1-138. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Bronchiolo-Alveolar Adenomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 15.28951263 |
| BMDL | 9.41161009 |
| BMDU | Infinity |
| AIC | 110.3012141 |
| P-value | 0.401340961 |
| D.O.F. | 2 |
| Chi ² | 1.825887871 |
| Slope Factor | 0.010625175 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 4 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.052388509 | 2.35E-02 | 0.0063456 | 0.098431418 |
| b1 | Bounded | NA | NA | NA |
| b2 | Bounded | NA | NA | NA |
| b3 | 2.94779E-05 | 6.54E-02 | -0.128136264 | 0.12819522 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.052388509 | 2.567036922 | 4 | 49 | 0.894373541 |
| 2 | 0.052611951 | 2.630597564 | 1 | 50 | -1.005355325 |
| 5 | 0.055873787 | 2.793689339 | 3 | 50 | 0.123433378 |
| 16 | 0.160167058 | 8.00835292 | 8 | 50 | -0.002951663 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -52.08803088 | 4 | - | - | NA |
| Fitted Model | -53.15060707 | 2 | 2.125152373 | 2 | 0.345564424 |
| Reduced Model | -55.67027459 | 1 | 5.039335051 | 3 | 0.168939394 |

Figure 1-139. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Bronchiolo-Alveolar Adenomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2.2 Bronchiolo-Alveolar Carcinomas in Female Mice

Female mice exhibited a significantly increased trend for the incidence of bronchiolo-alveolar carcinomas in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-107.

Table 1-107. Increased Incidence of Bronchiolo-Alveolar Carcinomas in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 49 | 1 |
| 2 | 50 | 0 |
| 5 | 50 | 1 |
| 16 | 50 | 3 |

The BMD modeling results for bronchiolo-alveolar carcinomas in female mice are summarized in Table 1-108. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The Multistage 3-degree model converged on the 2-degree model. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected. The 2-degree Multistage model was selected as the more parsimonious choice. A plot of the Multistage 2-degree model with a BMR of 10% ER is shown in Figure 1-140. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-141.

Table 1-108. Summary of BMD Modeling Results for Increased Incidence of Bronchiolo-Alveolar Carcinomas in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.6059 | 47.82 | 23 | 14 | 0.0069 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The Multistage 3-degree model converged on the 2-degree model. The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 2-degree Multistage model, which had the lowest AIC and was the more parsimonious choice. |
| Multistage 2 | 0.6059 | 47.82 | 23 | 14 | 0.0070 | |
| Multistage 1 | 0.5056 | 48.31 | 40 | 16 | 0.0062 | |

^a Selected model in bold.




Figure 1-140. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 2-Degree) for the Increased Incidence of Bronchiolo-Alveolar Carcinomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 23.02764702 |
| BMDL | 14.38723029 |
| BMDU | Infinity |
| AIC | 47.81567877 |
| P-value | 0.605883382 |
| D.O.F. | 2 |
| Chi ² | 1.002135499 |
| Slope Factor | 0.006950608 |

| Model Parameters | | | | |
|------------------|-------------|-------------|--------------|-------------|
| # of Parameters | 3 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.0113772 | 0.053529952 | -0.093539578 | 0.116293979 |
| b1 | Bounded | NA | NA | NA |
| b2 | 0.000198691 | 3.67E-02 | -0.071651227 | 0.072048609 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.0113772 | 0.557482822 | 1 | 49 | 0.592671971 |
| 2 | 0.012162611 | 0.608130555 | 0 | 50 | -0.77982726 |
| 5 | 0.016275792 | 0.813789577 | 1 | 50 | 0.206418171 |
| 16 | 0.060405966 | 3.020298307 | 3 | 50 | -0.011679786 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -21.13187787 | 4 | - | - | NA |
| Fitted Model | -21.90783939 | 2 | 1.551923026 | 2 | 0.460261021 |
| Reduced Model | -23.3559877 | 1 | 2.896296628 | 3 | 0.407892098 |

Figure 1-141. Details Regarding the Selected Model (Multistage 2-Degree) for the Increased Incidence of Bronchiolo-Alveolar Carcinomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2.3 Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice

Female mice exhibited a significantly increased trend for the incidence of bronchiolo-alveolar adenomas and carcinomas (combined) in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-109.

Table 1-109. Increased Incidence of Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 49 | 5 |
| 2 | 50 | 1 |
| 5 | 50 | 4 |
| 16 | 50 | 11 |

The BMD modeling results for bronchiolo-alveolar adenomas and carcinomas (combined) in female mice are summarized in Table 1-110. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 2-degree). A plot of the Multistage 2-degree model with a BMR of 10% ER is shown in Figure 1-142. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-143.

Table 1-110. Summary of BMD Modeling Results for Increased Incidence of Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.1016 | 132.0 | 13 | 7.6 | 0.013 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 2-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.2413 | 130.0 | 12 | 7.5 | 0.013 | |
| Multistage 1 | 0.1217 | 131.7 | 11 | 6.1 | 0.016 | |

^a Selected model in bold.




Figure 1-142. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 2-Degree) for the Increased Incidence of Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 12.14137745 |
| BMDL | 7.47918985 |
| BMDU | 19.79664615 |
| AIC | 130.0361342 |
| P-value | 0.241329764 |
| D.O.F. | 2 |
| Chi ² | 2.843181935 |
| Slope Factor | 0.013370432 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 3 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.061269856 | 2.31E-02 | 0.016075343 | 0.10646437 |
| b1 | Bounded | NA | NA | NA |
| b2 | 0.00071473 | 7.85E-02 | -0.153059287 | 0.154488747 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.061269856 | 3.002222954 | 5 | 49 | 1.15299002 |
| 2 | 0.063949778 | 3.197488882 | 1 | 50 | -1.228915905 |
| 5 | 0.077894352 | 3.894717584 | 4 | 50 | 0.053347965 |
| 16 | 0.218232649 | 10.91163246 | 11 | 50 | 0.026751484 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -61.33348348 | 4 | - | - | NA |
| Fitted Model | -63.01806711 | 2 | 3.369167271 | 2 | 0.185521661 |
| Reduced Model | -67.07521698 | 1 | 8.114299727 | 3 | 0.043707604 |

Figure 1-143. Details Regarding the Selected Model (Multistage 2-Degree) for the Increased Incidence of Bronchiolo-Alveolar Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2.4 Endometrial Stromal Polyps in Female Mice

Female mice exhibited a significantly increased trend for the incidence of endometrial stromal polyps in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-111.

Table 1-111. Increased Incidence of Endometrial Stromal Polyps in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 49 | 2 |
| 2 | 50 | 0 |
| 5 | 50 | 1 |
| 16 | 50 | 6 |

The BMD modeling results for endometrial stromal polyps in female mice are summarized in Table 1-112. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree). A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-144. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-145.

Table 1-112. Summary of BMD Modeling Results for Increased Incidence of Endometrial Stromal Polyps in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.3370 | 70.10 | 16 | 12 | 0.0086 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 3-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.2839 | 70.39 | 16 | 11 | 0.0087 | |
| Multistage 1 | 0.1521 | 72.23 | 21 | 11 | 0.0094 | |

^a Selected model in bold.




Figure 1-144. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Endometrial Stromal Polyps in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 15.88183212 |
| BMDL | 11.62829155 |
| BMDU | 33.54474972 |
| AIC | 70.10389143 |
| P-value | 0.336969894 |
| D.O.F. | 2 |
| Chi ² | 2.175523375 |
| Slope Factor | 0.008599716 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 4 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.019204564 | 3.85E-02 | -0.056279724 | 0.094688853 |
| b1 | Bounded | NA | NA | NA |
| b2 | Bounded | NA | NA | NA |
| b3 | 2.63012E-05 | 5.35E-02 | -0.10487753 | 0.104930133 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.019204564 | 0.941023655 | 2 | 49 | 1.091656458 |
| 2 | 0.019410912 | 0.970545585 | 0 | 50 | -0.98516272 |
| 5 | 0.022423786 | 1.121189281 | 1 | 50 | -0.114452357 |
| 16 | 0.119373076 | 5.968653792 | 6 | 50 | 0.012830596 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -31.60416819 | 4 | - | - | NA |
| Fitted Model | -33.05194571 | 2 | 2.895555043 | 2 | 0.235092195 |
| Reduced Model | -36.65806521 | 1 | 7.212238997 | 3 | 0.065432012 |

Figure 1-145. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Endometrial Stromal Polyps in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2.5 Mammary Gland Adenocarcinomas in Female Mice

Female mice exhibited a significantly increased trend for the incidence of mammary gland adenocarcinomas in a 2-year inhalation toxicity study of 1,2-dichloroethane ([Nagano et al., 2006](#)). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-113.

Table 1-113. Increased Incidence of Mammary Gland Adenocarcinomas in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 49 | 1 |
| 2 | 50 | 2 |
| 5 | 50 | 1 |
| 16 | 50 | 6 |

The BMD modeling results for mammary gland adenocarcinomas in female mice are summarized in Table 1-114. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree). A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-146. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-147.

Table 1-114. Summary of BMD Modeling Results for Increased Incidence of Mammary Gland Adenocarcinomas in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.7560 | 77.60 | 16 | 9.9 | 0.010 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 3-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.7044 | 77.77 | 17 | 9.7 | 0.010 | |
| Multistage 1 | 0.5949 | 78.34 | 18 | 9.1 | 0.011 | |

^a Selected model in bold.




Figure 1-146. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenocarcinomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 16.27040863 |
| BMDL | 9.941820292 |
| BMDU | Infinity |
| AIC | 77.60267152 |
| P-value | 0.755966155 |
| D.O.F. | 2 |
| Chi ² | 0.559517345 |
| Slope Factor | 0.01005852 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 4 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.026137909 | 3.28E-02 | -0.038234568 | 0.090510386 |
| b1 | Bounded | NA | NA | NA |
| b2 | Bounded | NA | NA | NA |
| b3 | 2.44615E-05 | 5.38E-02 | -0.105461386 | 0.105510309 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.026137909 | 1.280757549 | 1 | 49 | -0.248083557 |
| 2 | 0.026328467 | 1.316423363 | 2 | 50 | 0.595784802 |
| 5 | 0.029111123 | 1.455556151 | 1 | 50 | -0.377596042 |
| 16 | 0.118984218 | 5.949210887 | 6 | 50 | 0.020822887 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -36.5269587 | 4 | - | - | NA |
| Fitted Model | -36.80133576 | 2 | 0.548754125 | 2 | 0.760045437 |
| Reduced Model | -39.65162334 | 1 | 5.700575167 | 3 | 0.127122222 |

Figure 1-147. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Mammary Gland Adenocarcinomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2.6 Hepatocellular Adenomas in Female Mice

Female mice exhibited a significantly increased trend for the incidence of hepatocellular adenomas in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation dose for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-115.

Table 1-115. Increased Incidence of Hepatocellular Adenomas in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 49 | 1 |
| 2 | 50 | 1 |
| 5 | 50 | 1 |
| 16 | 50 | 6 |

The BMD modeling results for hepatocellular adenomas in female mice are summarized in Table 1-116. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 3-degree). A plot of the Multistage 3-degree model with a BMR of 10% ER is shown in Figure 1-148. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-149.

Table 1-116. Summary of BMD Modeling Results for Increased Incidence of Hepatocellular Adenomas in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.9911 | 70.08 | 16 | 11 | 0.0093 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 3-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.9337 | 70.21 | 16 | 11 | 0.0095 | |
| Multistage 1 | 0.6159 | 71.18 | 18 | 9.5 | 0.011 | |

^a Selected model in bold.




Figure 1-148. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 3-Degree) for the Increased Incidence of Hepatocellular Adenomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 15.86314678 |
| BMDL | 10.72443119 |
| BMDU | Infinity |
| AIC | 70.08200509 |
| P-value | 0.991114758 |
| D.O.F. | 2 |
| Chi ² | 0.017849902 |
| Slope Factor | 0.009324504 |

| Model Parameters | | | | |
|------------------|-------------|-------------|--------------|-------------|
| # of Parameters | 4 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.019111852 | 3.87E-02 | -0.056762546 | 0.09498625 |
| b1 | Bounded | NA | NA | NA |
| b2 | Bounded | NA | NA | NA |
| b3 | 2.63943E-05 | 0.053688515 | -0.105201162 | 0.105253951 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.019111852 | 0.936480755 | 1 | 49 | 0.065638085 |
| 2 | 0.019318949 | 0.96594745 | 1 | 50 | 0.034647579 |
| 5 | 0.022342749 | 1.117137467 | 1 | 50 | -0.11082622 |
| 16 | 0.119625443 | 5.981272148 | 6 | 50 | 0.007657574 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -33.03170698 | 4 | - | - | NA |
| Fitted Model | -33.04100255 | 2 | 0.018591143 | 2 | 0.990747499 |
| Reduced Model | -36.65806521 | 1 | 7.234125334 | 3 | 0.064798208 |

Figure 1-149. Details Regarding the Selected Model (Multistage 3-Degree) for the Increased Incidence of Hepatocellular Adenomas in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2.7 Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice

Female mice exhibited a significantly increased trend for the incidence of hepatocellular adenomas and carcinomas (combined) in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). For BMD modeling, the exposure concentrations were first duration adjusted to estimate an equivalent inhalation concentration for animals exposed for 24 hours per day, 7 days per week rather than 6 hours per day, 5 days per week. Then, the Multistage cancer models were fit to the dose-response data.

A BMR of 10% ER was chosen according to EPA's *BMD Technical Guidance* ([U.S. EPA, 2012](#)). The concentration and response data used for the modeling are presented in Table 1-117.

Table 1-117. Increased Incidence of Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice and Associated Concentrations Selected for Dose-Response Modeling for 1,2-Dichloroethane from a 2-Year Chronic Bioassay

| Adjusted Concentration (ppm) | Number of Animals | Incidence |
|------------------------------|-------------------|-----------|
| 0 | 49 | 2 |
| 2 | 50 | 1 |
| 5 | 50 | 2 |
| 16 | 50 | 6 |

The BMD modeling results for hepatocellular adenomas and carcinomas (combined) in female mice are summarized in Table 1-118. All Multistage models provided an adequate fit (chi-square p-value > 0.1) to the data. The BMDLs for the fit models were sufficiently close (differed by < 3-fold); therefore, the model with the lowest AIC was selected (Multistage 2-degree). A plot of the Multistage 2-degree model with a BMR of 10% ER is shown in Figure 1-150. Additional modeling details, including model parameters, goodness of fit at each concentration, and log likelihood are shown in Figure 1-151.

Table 1-118. Summary of BMD Modeling Results for Increased Incidence of Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay^a

| Model | Goodness of Fit | | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) | Basis for Model Selection |
|---------------------|-----------------|--------------|-----------------|------------------|------------------------|---|
| | p-value | AIC | | | | |
| Multistage 3 | 0.5369 | 86.42 | 17 | 10 | 0.0097 | All the Multistage models provided adequate fit (chi-square p-value > 0.1). The BMDLs were sufficiently close (differed by < 3-fold); therefore, EPA chose the 2-degree Multistage model, which had the lowest AIC. |
| Multistage 2 | 0.8193 | 84.43 | 17 | 10 | 0.0097 | |
| Multistage 1 | 0.6051 | 85.07 | 20 | 9.5 | 0.011 | |

^a Selected model in bold.




Figure 1-150. Plot of Response by Concentration with Fitted Curve for the Selected Model (Multistage 2-Degree) for the Increased Incidence of Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| Benchmark Dose | |
|------------------|-------------|
| BMD | 16.704916 |
| BMDL | 10.26208393 |
| BMDU | Infinity |
| AIC | 84.42822854 |
| P-value | 0.81925885 |
| D.O.F. | 2 |
| Chi ² | 0.398710378 |
| Slope Factor | 0.009744609 |

| Model Parameters | | | | |
|------------------|-------------|-----------|--------------|-------------|
| # of Parameters | 3 | | | |
| Variable | Estimate | Std Error | Lower Conf | Upper Conf |
| g | 0.030173657 | 3.26E-02 | -0.03365823 | 0.094005545 |
| b1 | Bounded | NA | NA | NA |
| b2 | 0.000377563 | 5.45E-02 | -0.106466417 | 0.107221543 |

| Goodness of Fit | | | | | |
|-----------------|-----------------------|-------------|----------|------|-----------------|
| Dose | Estimated Probability | Expected | Observed | Size | Scaled Residual |
| 0 | 0.030173657 | 1.478509213 | 2 | 49 | 0.428878843 |
| 2 | 0.031637234 | 1.581861679 | 1 | 50 | -0.462631823 |
| 5 | 0.039284849 | 1.964242466 | 2 | 50 | 0.025513498 |
| 16 | 0.119525528 | 5.976276377 | 6 | 50 | 0.009704333 |

| Analysis of Deviance | | | | | |
|----------------------|----------------|-----------------|-------------|-----------|-------------|
| Model | Log Likelihood | # of Parameters | Deviance | Test d.f. | P Value |
| Full Model | -40.00137558 | 4 | - | - | NA |
| Fitted Model | -40.21411427 | 2 | 0.42547738 | 2 | 0.808367344 |
| Reduced Model | -42.53972311 | 1 | 4.651217674 | 3 | 0.199192151 |

Figure 1-151. Details Regarding the Selected Model (Multistage 2-Degree) for the Increased Incidence of Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2.8 Combined Lung, Uterine, Mammary Gland, and Liver Tumors in Female Mice

Female mice exhibited significantly increased incidences of bronchiolo-alveolar adenomas and carcinomas, uterine endometrial stromal polyps, mammary gland adenocarcinomas, and hepatocellular adenomas and carcinomas in a 2-year inhalation toxicity study of 1,2-dichloroethane ([Nagano et al., 2006](#)). The MS Combo model was applied to the incidence data for the individual tumors and polydegrees identified for the individual tumor models, shown above (Sections 1.2.2.1 through 1.2.2.7), using a BMR of 10% ER.

The BMD Multistage Cancer/Multi-tumor modeling results for combined bronchiolo-alveolar adenomas and carcinomas, uterine endometrial stromal polyps, mammary gland adenocarcinomas, and hepatocellular adenomas and carcinomas in female mice are summarized in Table 1-119. A plot of the dose-response curves for the incidence of individual tumor types with a BMR of 10% ER is shown in Figure 1-152. Additional modeling details, including the cancer slope factor and log likelihood are shown in Figure 1-153.

Table 1-119. Summary of BMD Multi-Tumor (MS Combo) Modeling Results for Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Uterine Endometrial Stromal Polyps, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay

| Model | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) |
|--|-----------------------|------------------------|---------------------------|
| Multi-tumor (MS Combo) | 8.3 | 4.6 | 0.022 |
| Bronchiolo-alveolar carcinomas and adenomas in the lungs Multistage 2 | 12 | 7.5 | 0.013 |
| Endometrial stromal polyp in the uterus Multistage 3 | 16 | 12 | 0.0086 |
| Adenocarcinoma in the mammary gland Multistage 3 | 16 | 9.9 | 0.010 |
| Hepatocellular adenomas and carcinomas in the liver Multistage 2 | 17 | 10 | 0.0097 |




Figure 1-152. Plot of Response by Concentration with Fitted Curve for Selected Models used for the Multi-Tumor (MS Combo) Model for Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Uterine Endometrial Stromal Polyps, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

| User Input | | Model Results | | | | | | | | | | | | | | | | | | | |
|--|------------------------------|---------------|--|-----------|------------------------------|---|-----|------------------|------|------------|-------------|------|-------------|------|-------------|--------------|-------------|-------------------------|--------------|----------------------------------|-------------|
| <table border="1"> <thead> <tr> <th colspan="2">Info</th> </tr> </thead> <tbody> <tr> <td>Model</td> <td>frequentist Multi-tumor v1.0</td> </tr> </tbody> </table> | | Info | | Model | frequentist Multi-tumor v1.0 | <table border="1"> <thead> <tr> <th colspan="2">Benchmark Dose</th> </tr> </thead> <tbody> <tr> <td>BMD</td> <td>8.337858446</td> </tr> <tr> <td>BMDL</td> <td>4.613136813</td> </tr> <tr> <td>BMDU</td> <td>10.36198116</td> </tr> <tr> <td>Slope Factor</td> <td>0.021677224</td> </tr> <tr> <td>Combined Log-Likelihood</td> <td>-173.0854629</td> </tr> <tr> <td>Combined Log-Likelihood Constant</td> <td>148.7966046</td> </tr> </tbody> </table> | | Benchmark Dose | | BMD | 8.337858446 | BMDL | 4.613136813 | BMDU | 10.36198116 | Slope Factor | 0.021677224 | Combined Log-Likelihood | -173.0854629 | Combined Log-Likelihood Constant | 148.7966046 |
| Info | | | | | | | | | | | | | | | | | | | | | |
| Model | frequentist Multi-tumor v1.0 | | | | | | | | | | | | | | | | | | | | |
| Benchmark Dose | | | | | | | | | | | | | | | | | | | | | |
| BMD | 8.337858446 | | | | | | | | | | | | | | | | | | | | |
| BMDL | 4.613136813 | | | | | | | | | | | | | | | | | | | | |
| BMDU | 10.36198116 | | | | | | | | | | | | | | | | | | | | |
| Slope Factor | 0.021677224 | | | | | | | | | | | | | | | | | | | | |
| Combined Log-Likelihood | -173.0854629 | | | | | | | | | | | | | | | | | | | | |
| Combined Log-Likelihood Constant | 148.7966046 | | | | | | | | | | | | | | | | | | | | |
| <table border="1"> <thead> <tr> <th colspan="2">Model Options</th> </tr> </thead> <tbody> <tr> <td>Risk Type</td> <td>Extra Risk</td> </tr> <tr> <td>BMR</td> <td>0.1</td> </tr> <tr> <td>Confidence Level</td> <td>0.95</td> </tr> <tr> <td>Background</td> <td></td> </tr> </tbody> </table> | | Model Options | | Risk Type | Extra Risk | BMR | 0.1 | Confidence Level | 0.95 | Background | | | | | | | | | | | |
| Model Options | | | | | | | | | | | | | | | | | | | | | |
| Risk Type | Extra Risk | | | | | | | | | | | | | | | | | | | | |
| BMR | 0.1 | | | | | | | | | | | | | | | | | | | | |
| Confidence Level | 0.95 | | | | | | | | | | | | | | | | | | | | |
| Background | | | | | | | | | | | | | | | | | | | | | |

Figure 1-153. Details Regarding the Multi-Tumor (MS Combo) Model for the Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Uterine Endometrial Stromal Polyps, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

1.2.2.9 Combined Lung, Mammary Gland, and Liver Tumors in Female Mice (Alternate Analysis of Combined Tumors with Uterine Polyps Excluded)

This section presents an alternate analysis of combined tumors in female mice that excludes uterine polyps because these lesions are considered preneoplastic by some researchers. Female mice exhibited significantly increased incidences of bronchiolo-alveolar adenomas and carcinomas, mammary gland adenocarcinomas, and hepatocellular adenomas and carcinomas in a 2-year inhalation toxicity study of 1,2-dichloroethane (Nagano et al., 2006). The MS Combo model was applied to the incidence data for the individual tumors and polydegrees identified for the individual tumor models, shown above (Sections 1.2.2.1 through 1.2.2.7), using a BMR of 10% ER.

The BMD Multistage Cancer/Multi-tumor modeling results for combined bronchiolo-alveolar adenomas and carcinomas, mammary gland adenocarcinomas, and hepatocellular adenomas and carcinomas in female mice are summarized in Table 1-120. A plot of the dose-response curves for the incidence of individual tumor types with a BMR of 10% ER is shown in Figure 1-154. Additional modeling details, including the cancer slope factor and log likelihood are shown in Figure 1-155.

Table 1-120. Summary of BMD Multi-Tumor (MS Combo) Modeling Results for Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Following Inhalation Exposure to 1,2-Dichloroethane in a 2-Year Chronic Bioassay

| Model | BMD 10%ER (ppm) | BMDL 10%ER (ppm) | Slope Factor (per ppm) |
|---|-----------------|------------------|------------------------|
| Multi-tumor (MS Combo) | 9.0 | 4.8 | 0.021 |
| Bronchiolo-alveolar carcinomas and adenomas in the lungs Multistage 2 | 12 | 7.5 | 0.013 |
| Adenocarcinoma in the mammary gland Multistage 3 | 16 | 9.9 | 0.010 |
| Hepatocellular adenomas and carcinomas in the liver Multistage 2 | 17 | 10 | 0.0097 |




Figure 1-154. Plot of Response by Concentration with Fitted Curve for Selected Models Used for the Multi-Tumor (MS Combo) Model for Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)




Figure 1-155. Details Regarding the Multi-Tumor (MS Combo) Model for the Increased Incidence of Bronchiolo-alveolar Adenomas and Carcinomas, Mammary Gland Adenocarcinomas, and Hepatocellular Adenomas and Carcinomas (Combined) in Female Mice Exposed to 1,2-Dichloroethane Via Inhalation (2-Year Study)

REFERENCES

- Aafjes, JH; Vels, JM; Schenck, E. (1980). Fertility of rats with artificial oligozoospermia. *J Reprod Fertil* 58: 345-351. <https://hero.epa.gov/reference/1332483/>
- Dow Chemical. (2006). Re: Testing consent order for ethylene dichloride; final report (docket no . OPPT-2003-0010) [TSCA Submission]. (Study ID No. 041115. 40060000065). HAP Task Force for Ethylene Dichloride. <https://hero.epa.gov/reference/6570013/>
- Igwe, OJ; Que Hee, SS; Wagner, WD. (1986). Interaction between 1,2-dichloroethane and disulfiram. I. Toxicologic effects. *Fundam Appl Toxicol* 6: 733-746. <https://hero.epa.gov/reference/200386/>
- IRFMN. (1978). Clinical chemistry results in adult rats exposed to ethylene dichloride by inhalation for 12 months [TSCA Submission]. (OTS0515737. 86-870001661). Shell Oil Company. <https://hero.epa.gov/reference/5447364/>
- Meistrich, ML; Finch, M; da Cunha, MF; Hacker, U; Au, WW. (1982). Damaging effects of fourteen chemotherapeutic drugs on mouse testis cells. *Cancer Res.* <https://hero.epa.gov/reference/7565892/>
- Munson, AE; Sanders, VM; Douglas, KA; Sain, LE; Kauffmann, BM; White Jr, KL. (1982). In vivo assessment of immunotoxicity. *Environ Health Perspect* 43: 41-52. <https://hero.epa.gov/reference/62637/>
- Nagano, K; Umeda, Y; Senoh, H; Gotoh, K; Arito, H; Yamamoto, S; Matsushima, T. (2006). Carcinogenicity and chronic toxicity in rats and mice exposed by inhalation to 1,2-dichloroethane for two years. *J Occup Health* 48: 424-436. <https://hero.epa.gov/reference/200497/>
- NTP. (1991). Toxicity studies of 1,2-dichloroethane (ethylene bichloride) (CAS No. 107-06-2) in F344/N rats, Sprague Dawley rats, Osborne-Mendel rats, and B6C3F1 mice (drinking water and gavage studies). In Toxicity Report Series, vol 4. (NTP TOX 4; NIH Publication No. 91-3123). Research Triangle Park, NC. <https://hero.epa.gov/reference/1772371/>
- Payan, JP; Saillenfait, AM; Bonnet, P; Fabry, JP; Langonne, I; Sabate, JP. (1995). Assessment of the developmental toxicity and placental transfer of 1,2-dichloroethane in rats. *Toxicol Sci* 28: 187-198. <https://hero.epa.gov/reference/12099/>
- Rao, KS; Murray, JS; Deacon, MM; John, JA; Calhoun, LL; Young, JT. (1980). Teratogenicity and reproduction studies in animals inhaling ethylene dichloride. In B Ames; P Infante; R Reitz (Eds.), (pp. P149-P166). Cold Spring Harbor, NY: Cold Spring Harbor Laboratory. <https://hero.epa.gov/reference/5453539/>
- Robaire, B; Smith, S; Hales, BF. (1984). Suppression of spermatogenesis by testosterone in adult male rats: effect on fertility, pregnancy outcome and progeny. *Biol Reprod* 31: 221-230. <https://hero.epa.gov/reference/4992207/>
- Storer, RD; Jackson, NM; Conolly, RB. (1984). In vivo genotoxicity and acute hepatotoxicity of 1,2-dichloroethane in mice: Comparison of oral, intraperitoneal, and inhalation routes of exposure. *Cancer Res* 44: 4267-4271. <https://hero.epa.gov/reference/200614/>
- U.S. EPA. (2012). Benchmark dose technical guidance [EPA Report]. (EPA100R12001). Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <https://hero.epa.gov/reference/1239433/>
- Working, PK. (1988). Male reproductive toxicology: comparison of the human to animal models [Review]. *Environ Health Perspect* 77: 37-44. <https://hero.epa.gov/reference/12974611/>
- Zeng, N; Jiang, H; Fan, Q; Wang, T; Rong, W; Li, G; Li, R; Xu, D; Guo, T; Wang, F; Zeng, L; Huang, M; Zheng, J; Lu, F; Chen, W; Hu, Q; Huang, Z; Wang, Q. (2018). Aberrant expression of miR-451a contributes to 1,2-dichloroethane-induced hepatic glycerol gluconeogenesis disorder by inhibiting glycerol kinase expression in NIH Swiss mice. *J Appl Toxicol* 38: 292-303. <https://hero.epa.gov/reference/5555689/>

Zhang, Y; Li, G; Zhong, Y; Huang, M; Wu, J; Zheng, J; Rong, W; Zeng, L; Yin, X; Lu, F; Xie, Z; Xu, D; Fan, Q; Jia, X; Wang, T; Hu, Q; Chen, W; Wang, Q; Huang, Z. (2017). 1,2-dichloroethane induces reproductive toxicity mediated by the CREM/CREB signaling pathway in male NIH Swiss mice. Toxicol Sci 160: 299-314. <https://hero.epa.gov/reference/4453049/>