

Integration and Synthesis Summary for Fish

This Integration and Synthesis Summary includes our jeopardy analysis for fish species that we or EPA determined would “likely be adversely affected” by the proposed action. Our jeopardy analysis of the proposed action’s impacts to listed species is split into three major factors: vulnerability, exposure, and toxicity. The tables below contain summaries of vulnerability, exposure, and toxicity. Data and information used for each species include environmental baselines, cumulative effects, exposure information, and expected toxic effects for all species, and a template worksheet to show how species were assessed are in Appendix E. Status of the Species for each species can be found in Appendix B.

Most of these species have low exposure to atrazine due to the factors described in the tables or individual rationales below, in combination with reductions in atrazine spray drift and runoff resulting from implementation of conservation measures added to the product label (including those developed during this consultation through the Herbicide Strategy^{Error! Bookmark not defined.}; see Conservation Measures section below). We anticipate that these measures will reduce exposure from agricultural uses to a level where no more than low level adverse effects are anticipated for many listed fish species.

Vulnerability

For the fish species that we or EPA determined are “likely to be adversely affected” by the proposed action, we considered several factors for each listed fish to determine the current vulnerability of that species to additional stressors. This effort allows us to consider whether a species’ current condition is stable, moving toward recovery, or moving toward further decline. In general, we expect the species’ vulnerability to additional stressors to be higher if they are near extinction, far from recovery, or moving toward further decline than if their condition is stable or improving. We also identify which species are most (and least) susceptible to additional stressors in general based on information from species listing and recovery documents, or other sources as cited and considered in the Status of the Species and Critical Habitat section of this Opinion (Appendix B).

Our assessment of vulnerability focuses on six factors (as currently understood and available): (1) the species listing status and recent 5-year status review recommendation (if available), (2) distribution, (3) number of populations¹, (4) species population trends, (5) if pesticides have been noted as a threat, and (6) current and projected future impacts from activities associated with environmental baseline and cumulative effects. We obtained the information to create the vulnerability summary from the Status of the Species accounts (Appendix B), overarching

¹ The number will vary in value and importance by species and in some cases is unknown. In general, species with a greater number of populations have greater representation, will be more resilient, and when distributed geographically, will have greater redundancy. Conversely, species with fewer populations, in general, have less representation, are less resilient, and have less redundancy.

Environmental Baseline section of the Opinion, five-year species status reviews, species recovery plans, species status assessments, range and critical habitat information from our ECOS² repository, and other sources containing the best available scientific information for the species.

We scored each of the six vulnerability components with high, medium, or low scores. We assigned a high vulnerability ranking to a species if all vulnerability components were scored as high, a mixture of medium and high, or if a threatened species was recommended for uplisting to endangered status in the most recent 5-year status review or proposed rule. We assigned a medium vulnerability ranking if a species' scores were all medium, a mix of high, medium, and low, or a mix of high and low (unless the species has been recommended for uplisting or delisting). We assigned a low vulnerability ranking to species with only low scores, a mixture of low and medium scores, or if the species was recommended for delisting. Considerations regarding specific aspects of the species' vulnerability or beyond what was included in the vulnerability ranking were applicable in our jeopardy analyses for some species depending on unique aspects of their vulnerability factors, recovery needs, or life history. This information is reflected in the rationales for conclusion below.

Exposure

We anticipate listed fish species will be exposed to atrazine primarily through direct contact in water. Atrazine is moderately mobile in water and is relatively persistent in the environment relative to other pesticides on the market, indicating that off-site transport, particularly through runoff, may result in exposure to listed fish species in areas far from agricultural and non-agricultural use sites.

Exposure to Agricultural Uses

Atrazine has several registered agricultural uses (see Appendix 1-4 of EPA's Biological Evaluation) in the conterminous United States. We characterize the expected level of exposure using overlaps between the species' ranges and agricultural areas where atrazine is registered for use (i.e., overlap data), past atrazine usage data (when available; the amount and location where atrazine has been used in the past), any species-specific considerations such as life history information (e.g., habitat preferences, dietary needs, dispersal behavior), and existing protections or conservation actions (e.g., existing label measures, conservation measures from the action agency). Instead of using the species' range, the EPA uses the HUC-12 watersheds that contain the species range to calculate the extent of overlap and past atrazine usage. Given that we typically do not anticipate the specific waterbodies required for fully aquatic listed species occur directly in atrazine use sites and that off-site exposure is inherently included when overlap and

² <https://ecos.fws.gov/ecp/>

usage are calculated at a watershed scale, we use the watershed overlap and usage data to characterize potential exposure to agricultural uses of atrazine.

Species with greater than 10% overlap between their range and agricultural atrazine use sites are assigned a high overlap score, species with 5-10% overlap are assigned a medium overlap score, and species with less than 5% overlap are assigned a low overlap score. In addition to range overlaps with use sites, we considered past usage data within a species' range to determine how much of a species' range we expect to be treated with atrazine each year of the proposed action. Except where otherwise noted, usage data is provided by EPA applying data from their National and State Summary Use and Usage Matrix, as described in the *Usage Analysis* section of this Opinion. Species with usage data that indicate a large portion of their range (>10%) is treated with atrazine each year are assigned a high usage score. Species that have a medium portion of their range (5-10%) treated with atrazine each year are assigned a medium usage score, and species where data indicate a low portion of their range (<5%) is treated with atrazine each year are assigned a low usage score.

We determine the agricultural exposure ranking by qualitatively considering both the total overlap and total usage, as well as any additional exposure considerations that might modify the level of exposure likely to occur. When overlap and usage scores are the same, we assign the agricultural exposure ranking the same score (e.g., if both overlap and usage is high, the agricultural exposure ranking is high). In cases where overlap is high and usage is medium or when overlap is medium and usage is low, we use the overlap score as the agricultural exposure ranking to maintain conservative exposure assumptions. As usage is a subset of overlap, the overlap score will always be greater than the usage score. In cases where overlap is high, but usage is low, we anticipate a moderate portion of the range may be treated over the duration of the proposed action even if only a small portion of the range is treated in any given year (particularly if the areas treated occur in different locations each year), leading to an agricultural exposure ranking of medium. For species where there are additional exposure considerations, we adjust the agricultural exposure ranking to reflect this additional information, as appropriate.

Agricultural uses of atrazine include labeled uses for corn, vegetables and ground fruit (i.e., sweet corn), sod, orchards (i.e., guava and macadamia nut), other grains (including sugarcane and sorghum), and fallow fields only within the conterminous United States.

Exposure to Non-Agricultural Uses

In addition to agricultural uses, atrazine is registered for use on non-agricultural turf, including residential lawns and golf course turf. UDLs for non-agricultural uses sites that represent turf tend to be less defined than those for agricultural UDLs and are less likely to accurately represent the actual footprint of these use sites on the landscape. As such, we assess exposure of species to all non-agricultural uses of atrazine in a qualitative manner, considering the life history of species, methods of application, atrazine usage, and any existing conservation measures to reduce drift and runoff or otherwise limit exposure to species. To facilitate this analysis, for

every species in this Appendix, we reviewed species' documents (e.g., Status of the Species (Appendix B), 5-year reviews, Species Status Assessments, recovery plans, listing rules) to determine if the species could occur near non-agricultural atrazine use sites (i.e., residential areas where lawns or golf courses are likely present) and the manner in which they may rely on these sites.

Depending on region, cool-season, warm-season, or a combination of turf grass species are managed on golf courses and lawns. Cool-season grasses grow best in cooler conditions, and warm-season grasses thrive in hot, dry weather (USDA, 2004); there is a transition zone across the U.S. where either category of turf grasses may be planted based on microclimate conditions. Exposure to triazines will kill cool-season grasses, but warm-season grasses can tolerate exposure to atrazine. As such, EPA estimated where in the U.S. only cool-season grasses are exclusively used in turf based on the U.S. Department of Agriculture's plant hardiness zone map as atrazine use is not expected in these areas (USDA, 2023). Because hardiness zones will change over time with environmental conditions, EPA created a static map based on the hardiness zones where they expect warm- and cool-season grasses are grown based on the most recent data mapped (i.e., 1991-2020). EPA determined zones 1a-6a represent cool-season grasses (i.e., white areas) and zones 6b-13b may include warm-season grasses (i.e., black areas) (Figure 1). We expect the cool- and warm-season grass assessment to apply to all turf, including residential, commercial, and golf course turf. We refer to EPA's cool-season map in species assessments where relevant, particularly if a species occurs exclusively in the cool-season zone where we expect atrazine will not be used on turf and no exposure will occur from this use.




Figure 1. Map showing where cool-season grasses (white areas) and warm-season grasses (black areas) are used on turf across the continental U.S.

Particularly for residential and commercial turf uses, qualitative usage information obtained by EPA from the National Association of Landscape Professionals (NALP) indicate that atrazine is no longer commonly used on residential or commercial turf due to preferential use of newer herbicides. If atrazine were used on residential or commercial turf, it would be applied during the fall and spring as a pre-emergent. In addition, commercial and residential applicators typically apply herbicides with hand-held equipment that release coarse droplets, limiting the potential for spray drift.

Particularly for golf course turf uses, we obtained qualitative usage information directly from the Golf Course Superintendents Association of America (GCSAA) and an academic turf scientist that indicate that atrazine is used to control winter annual broadleaf and annual bluegrass weeds on golf courses. They are applied as a pre-emergent in early fall and early winter to fairways and roughs, which make up approximately 30% of a golf course's acreage. Triazines are not applied to tee boxes or greens, which make up an additional 6% of golf course acreage. Most applications are made at rates lower than what is on the label (i.e., 1-1.5 lbs a.i./acre). These applications are made only once or twice a year, 45-60 days apart. In general, golf courses typically apply herbicides using dedicated ground equipment with a low boom height (as per the label), and golf course superintendents make use of several tools to monitor soil moisture before any applications are made to help ensure turf and soil conditions do not lead to off-target movement of herbicides. In addition, riparian buffer zones are often used on golf courses between all water features to reduce off target movement (Golf Course Superintendents Association of America [GCSAA], pers. comm., 2025). The no-till methodology and continuous cover of a turf grass area inherent in managing golf course turf are equivalent to additional run-off mitigations (i.e., equivalent to six points on EPA's mitigation menu), and we considered them in our assessment.

For most species in this Appendix, we anticipate that non-agricultural uses will not meaningfully add to the overall level of anticipated exposure considered in our analysis of agricultural uses. Due to runoff and spray drift considerations described above, off-site exposure is not expected to result in effects to most species in this Appendix. In addition, we expect most listed species' habitat requirements preclude them from occupying non-agricultural use sites where atrazine may be used. For species whose habitat is known or presumed to occur in non-agricultural use sites of atrazine, we consider, individually and qualitatively, the extent and manner of non-agricultural atrazine usage within the species' range to generally determine whether a small, moderate, or large number of individuals are likely to be exposed and the expected level of adverse effects from non-agricultural exposure of atrazine.

References:

GCSAA (Golf Course Superintendents Association of America). 2025. Personal communication with USFWS HQ staff.

USDA. 2004. Comparing warm-season and cool-season grasses for erosion control, water quality, and wildlife habitat. Natural Resources Conservation Service, U.S. Department of Agriculture. 5 pp.

USDA. 2023. Plant Hardiness Zone Map. Agricultural Research Service, U.S. Department of Agriculture. Accessed from <https://planthardiness.ars.usda.gov/> on August 20, 2025.

Toxicity

We characterize the expected toxic effect to species based on the anticipated level of direct and indirect³ adverse effects to individuals. Our analysis of toxicity assumes individuals are exposed to atrazine at levels estimated by EPA's environmental exposure modeling and is focused on determining the level of adverse effect expected to occur once exposure has taken place. Direct effects are based on the anticipated level of mortality and sublethal effects (e.g., reduced growth) likely to occur in exposed individuals. Indirect effects are based on the impact a listed species is likely to experience when the organisms they rely on, such as those that act as food sources, are exposed to atrazine and experience adverse effects.

We consider in our analysis if EECs exceed the threshold for sublethal effects on growth to the fish, or reproductive effects to the aquatic invertebrate as fish prey. Data indicate reduced reproduction can occur to fish at EECs that may be observed in concentrations in lower flow or lower volume water bodies within the range for some fishes. We consider the NOAEC a conservative threshold for qualitatively estimating anticipated sublethal effects to listed fishes.

Because mortality from atrazine exposure at estimated environmental concentrations to fish is not anticipated (see section *Effects to Fish and Aquatic-phase Amphibians* in the main body of the Opinion), we focus our assessment on sublethal effects to fish, and indirect effects to the fish from effects to fish prey (e.g., other fish, aquatic invertebrates) and plant related food resources. Concentrations of atrazine can vary greatly among different regions and aquatic habitat types. Where atrazine enters smaller streams or static waters (e.g., low flow/low volume waterbodies) from runoff or spray drift, we generally anticipate high levels of sublethal effects to individual fish where exposure occurs. In larger waterbodies (e.g., where concentrations may be lower due to dilution or other factors as described in the *Effects of the Action* section of the Biological Opinion), we expect lower levels of sublethal effects fish. We determine the agricultural toxicity

³ While our Opinion considers all consequences of the proposed action (per the definition of effects of the action at 50 CFR Part 402.02), the terms "direct" and "indirect" effects were used in EPA's BE, and are used in environmental risk assessment terminology in general, and do not have the same meaning as used in ESA regulations. As used in the effects analysis section, direct effects to species are those caused by the pesticide itself through dietary, dermal, or inhalation routes of exposure. Indirect effects occur when the pesticide acts on elements of the ecosystem that are required by the species, such as alterations to prey or shelter. Thus, in the effects analysis section, we may sometimes continue to use these terms to link back to the analysis in EPA's BE.

ranking for fish by qualitatively assessing the expected levels of adverse effects (i.e., sublethal effects to the fish).

Fishes can consume other fish, amphibians, arthropod prey such as aquatic insects, and plant-based and microbial resources (e.g., algae, detritus). We expect some impacts to fish or aquatic amphibian prey, and we anticipate some effects to aquatic arthropod prey but not to the extent it will reduce the listed fish species' ability to forage as atrazine exposure is not expected to eliminate these prey items completely in the aquatic habitats where they are found. We anticipate some fish dietary items (e.g., phytoplankton, algae) in low flow or low volume waterbodies will be impacted by atrazine applications but we do not anticipate that it will eliminate all algae or phytoplankton within a water body and that these resources will replenish over time in any dynamic aquatic system (flowing or non-flowing) based on several mesocosm and microcosm studies discussed in the main body of the Opinion. We don't expect impacts to detritus from atrazine exposure. Therefore, we do not anticipate significant reductions in plant-base food availability for fish. As such, impacts to growth to fish are the primary driver in determining the toxicity ranking for listed fish species.

Experimental Populations, Nonessential

We considered the following nonessential experimental populations for fish species in this section of the consultation: The boulder darter (Entity ID 8921), bull trout (Entity ID 10037), Colorado pikeminnow (Entity ID 2142), duskytail darter (Entity ID 9502, 6503), Rio Grande silvery minnow (Entity ID 10052), slender chub (Entity ID 9504), smoky madtom (Entity ID 5981), spotfin chub (Entity IDs 1934, 9061 and 9505), Topeka shiner (Entity ID 10910), yellowfin madtom (Entity IDs 2956, 4496, 9506), and woundfin (Entity ID 2599). We do not provide separate analyses and jeopardy determinations for these populations. Rather, we treat all populations of the species (including populations designated as experimental) as a single listed entity when making jeopardy determinations or for other analyses in a section 7 consultation. An "essential experimental population" is a reintroduced population whose loss would be likely to appreciably reduce the likelihood of the survival of the species in the wild. However, there are no "essential experimental populations" in this consultation. A "nonessential experimental population" is a reintroduced population whose loss would not be likely to appreciably reduce the likelihood of survival of the species in the wild. By definition, a "nonessential experimental population" is not essential to the continued existence of the species. Therefore, no proposed action impacting a population so designated could lead to a jeopardy determination for the entire species. In cases where our assessment of the listed entity (i.e., the non-experimental population(s) of the species) leads to a "not likely to jeopardize" determination, we generally assume any added effects to the nonessential experimental population will not change these determinations. However, we consider the role of the experimental population in the survival and recovery of the species and consider this information in our jeopardy analyses as appropriate.

Conservation Measures

The technical registrants have previously agreed to substantial conservation measures that were incorporated into EPA's 2021 BE. These conservation measures include the following:

- Prohibit use in Hawaii, Alaska, and the Territories,
- Prohibit use on roadsides, shelterbelts, Conservation Reserve Program (CRP) land, conifers (including Christmas tree plantings), timber and forestry, and miscanthus and other perennial bioenergy crops,
- Prohibit application via mechanically pressurized handguns to macadamia nuts, sweet corn, and guava,
- Restrict "fallow" uses on all labels to the following scenarios and geographies only:
 - Wheat-corn-fallow and wheat-fallow-wheat in CO, KS, ND, NE, SD, and WY,
 - Wheat-sorghum-fallow in AR, CO, GA, IL, KS, LA, MS, MO, NE, NM, NC, OK, SD, and TX
- Reduce the single maximum application rate of turf, granular formulations to 2.0 lbs. AI/A, and reduce the single maximum application rate of turf, sprays to 1.0 lb. AI/A,
- Restrict applications made by backpack-spray to landscape turf to spot treatments only,
- Restrict applicators from applying atrazine products to the same sorghum acre,
- Require all applications to use coarse or coarser droplet sizes,
- Require an in-field downwind buffer of 15-ft for all ground applications (from the edge of all streams and rivers as well as the high-tide line for all estuarine/marine environments, and from threatened and endangered species critical habitat and/or species locations)
- Prohibit all ground applications when wind speeds exceed 10 miles per hour at the application site,
- For ground boom applications, only apply with the release height recommended by the manufacturer, but no more than 4-ft above the ground or crop canopy,
- Require an in-field downwind buffer of 150-ft for all aerial applications (from the edge of all streams and rivers as well as the high-tide line for all estuarine/marine environments, and from threatened and endangered species critical habitat and/or species locations),
- If the windspeed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the field,
- If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters,
- Prohibit all aerial applications when wind speeds exceed 15 miles per hour at the application site,

- Restrict aerial applications from releasing spray at a height greater than 10-ft above the ground or vegetative canopy unless a greater application height is necessary for pilot safety,
- Prohibit aerial applications of non-liquid formulations,
- Prohibit all applications during temperature inversions.

While these conservation measures are impactful and contribute to reducing the level of exposure and adverse effects to listed species, EPA and the Service anticipate substantial risk of adverse effects to many listed species remain after incorporating these measures into the proposed action.

Herbicide Strategy Conservation Measures

As part of the atrazine ESA consultation with the Service, EPA is implementing the final Herbicide Strategy to inform and identify any necessary conservation measures where EPA's analysis indicated there was a risk of population level effects to listed species. The measures identified by EPA, and committed to by the technical registrants, include:

- a standard 170-foot wind-directional spray drift buffer for aerial applications⁴ (not in addition to the buffers the technical registrants committed to previously), and
- a minimum of three runoff mitigation points⁵ necessary in all areas where atrazine is used, as well as additional runoff mitigation points (i.e., six points total) for certain atrazine uses limited to certain geographic areas when required to protect specific listed species.

In addition to the conservation measures identified through EPA's Herbicide Strategy, in the course of this consultation the technical registrants have also committed to additional measures for specific registered uses of atrazine to reduce exposure to listed species, including:

- Reduce the maximum annual application rate for field corn from 2.5 lbs. AI/A to 2.0 lbs. AI/A,
- For sweet corn uses, adopt one of the following:
 - Do not apply atrazine to sweet corn from August 15th to November 1st; when applied during other times of the year, use as a pre-emergent up to 2.0 lbs ai/acre.
 - With no timing restrictions for use, use as pre-emergent up to 1.25 lbs ai/acre followed by post-emergent 0.75 lbs ai/acre.

⁴ Note: The 170-foot aerial buffer replaces the 150-foot aerial buffer agreed to before implementation of the Herbicide Strategy.

⁵ Ecological Mitigation Support Document to Support Endangered Species Strategies

- Restrict “corn” in wheat-corn-fallow rotations to “field corn” meaning “wheat-field corn-fallow rotations”,
- Off-label all uses in California except for Imperial County, and
- Add the restriction “Do not apply atrazine products during rain or when soils are saturated or above field capacity” to all formulations.

The spray drift buffers will be placed on the general label and will apply to all uses of atrazine. EPA’s Herbicide Strategy provides applicators with options to reduce the distance of this buffer by using other spray drift reduction strategies that we anticipate will result in an equivalent reduction in spray drift entering non-target habitats as stated buffers. These measures and the degree to which applicators can reduce buffers by employing them are described in EPA’s Herbicide Strategy and EPA’s Ecological Mitigation Support Document to Support Endangered Species Strategies. These documents are provided in Appendix A-1.

Based on EPA’s analyses, the required spray drift conservation measures described above (from the current label, those from implementation of the Herbicide Strategy, and additional measures committed to through consultation for specific registered atrazine uses) will reduce spray drift from entering species’ habitats by >95%. The Service anticipates that this reduction will minimize off-site transport of atrazine from spray drift to a level where no more than low levels of effects are likely to occur to most species.

As stated above, all agricultural labels will include a requirement for applicators to achieve three points of runoff mitigation, as described in the Herbicide Strategy, for all agricultural uses. EPA’s Herbicide Strategy provides applicators with various options to reduce runoff and erosion and assigns points to each option based on its effectiveness. Applicators must implement sufficient mitigation points to meet the label requirement. Applicators can achieve the required points using the conservation measures identified on EPA’s Mitigation Menu website⁶. The menu provides a suite of options, including relief points for certain field characteristics and likelihood for pesticide transport.

We expect implementation of the required runoff and erosion reduction measures to minimize off-site transport of atrazine to habitats of listed species. EPA’s analyses indicated that the general label requirement of three runoff mitigation points will reduce estimated environmental concentrations of atrazine in runoff by up to an order of magnitude (i.e., up to 90% reduction, in other words reduce pesticide loading to one-tenth of pre-runoff mitigation levels).

In cases where EPA has identified additional runoff measures are needed, additional points (up to six points total) will be required. EPA will communicate where additional runoff mitigation points are needed and for what specific atrazine uses through their Bulletins Live! Two online platform, which all applicators are required to check before making pesticide applications. In

⁶ Mitigation Menu website: <https://www.epa.gov/pesticides/mitigation-menu>

areas requiring up to six runoff mitigation points total, EPA expects estimated environmental concentrations of atrazine will decrease by up to two orders of magnitude (i.e., reduce pesticide loading to one-one hundredth of pre runoff mitigation levels; 99% reduction).

For all the species in this document, we expect that the runoff and mitigation measures will reduce exposure concentrations to within one order of magnitude of the exposure level where fish exposed to atrazine in areas off-site will not accumulate more than low levels of atrazine and are not likely to experience more than low levels of sublethal adverse effects to growth or reproduction (if any). Additionally, we anticipate these agricultural measures will reduce exposure to plant species, resulting in no more than low levels of adverse effects to plants that provide food or habitat features for listed fish species.

Summary of Conclusions for Fish

After reviewing the current status of the species, the environmental baseline for the action area, the effects of the proposed action, and the cumulative effects, it is our biological opinion that the registration of simazine, as proposed, is not likely to jeopardize the continued existence of at least 100 of the 109 fish species in this Appendix. For the remaining nine fish in this appendix, we plan to continue coordination with EPA and the technical registrants to further assess these species.

In our analysis, some species that had the same or very similar rationales for their conclusions were grouped together, to increase efficiency and avoid repetition. Relevant information and data unique to each individual species was considered when assigning species to groups and incorporated into the rationales as appropriate. Species-specific information (e.g., environmental baseline, cumulative effects, status of the species, exposure, and toxicity) was considered for all species, including those species in the grouped analyses, and are presented in full in Appendices B and E. This approach allowed us to streamline our discussion in this Opinion by avoiding repeating our findings when species in the respective groupings would be expected to be affected similarly. The use of these groupings, therefore, does not mean that our evaluation failed to evaluate each individual species. On the contrary, our detailed process for each species-specific analysis remained the same, including for species for which we summarized our findings in tables below.

Species proposed for delisting

Species with low exposure informed by low overlap with agriculture and low likelihood of non-agricultural exposure

Species with low agricultural exposure achieved through conservation measures and low likelihood of non-agricultural exposure

Species requiring further analysis

In our draft Biological Opinion, we focused our analyses on 1) species with low expected exposure to atrazine (due to low overlap, usage, or conservation measures adopted prior to consultation), and 2) species with more than low levels of exposure that benefited from conservation measures identified through the Herbicide Strategy that aimed to reduce off-site transport of atrazine (i.e., listed plants and listed animals that depend on plant resources). For the species in Table 4, we identified the need for further coordination. We expect Herbicide Strategy conservation measures to reduce pesticide loading into aquatic habitats by up to 90% (i.e., one order of magnitude) compared to unmitigated runoff. We anticipate that this reduction will minimize off-site transport and reduce the likelihood, magnitude, and frequency of exposure of atrazine to a level where no more than low levels of adverse effects are likely to occur to listed mammals through this exposure route. While the conservation measures are expected to reduce the extent of off-field exposure and reduce exposure concentrations, we anticipate atrazine residues could remain at levels high enough to cause greater than low levels of adverse direct and/or indirect effects to these fish. We intend to continue coordinating with EPA and atrazine registrants between the release of this draft Opinion and the transmission of the final Opinion to gain information regarding the exposure and effects of each species to atrazine. As such, we have not yet made determinations for these species.

Table 4. Species requiring further analysis

| Common Name | Scientific Name | Vulnerability Ranking | Agricultural Exposure Ranking | Toxicity Ranking |
|-----------------------|-----------------------------------|-----------------------|-------------------------------|------------------|
| Arkansas River shiner | <i>Notropis girardi</i> | High | High | High |
| Carolina madtom | <i>Noturus furiosus</i> | High | High | High |
| Fountain darter | <i>Etheostoma fonticola</i> | High | High | High |
| June sucker | <i>Chasmistes liorus</i> | Medium | High | High |
| Pallid sturgeon | <i>Scaphirhynchus albus</i> | Medium | High | High |
| Peppered chub | <i>Macrhybopsis tetranema</i> | High | High | High |
| Topeka shiner | <i>Notropis topeka</i> (=tristis) | Medium | High | High |
| Yaqui catfish | <i>Ictalurus pricei</i> | High | High | High |
| Yaqui chub | <i>Gila purpurea</i> | Medium | High | High |